

Comparative modulation of lncRNAs in wild-type and *rag1*-heterozygous mutant zebrafish exposed to immune challenge with spring viraemia of carp virus (SVCV)

Valentina Valenzuela-Muñoz^{1*}, Patricia Pereiro^{1,2*}, Margarita Álvarez-Rodríguez², Cristian Gallardo-Escárate¹, Antonio Figueras², Beatriz Novoa²

¹Laboratory of Biotechnology and Aquatic Genomics, Interdisciplinary Center for Aquaculture Research (INCAR), University of Concepción, Concepción, Chile.

²Instituto de Investigaciones Marinas (IIM), CSIC, Vigo, Spain.

*: these authors have contributed equally to this work

Corresponding Authors

Beatriz Novoa, Instituto de Investigaciones Marinas (IIM), CSIC, Eduardo Cabello 6, 36208, Vigo, Spain
(beatriznovoa@iim.csic.es)

C Gallardo-Escárate, Universidad de Concepción, P.O. Box 160-C, Concepción 4030000, Chile (e-mail:
crisgallardo@udec.cl)

Supplementary figures

Figure S1. Survival curve of WT and *rag1^{+/−}* zebrafish infected with SVCV.




Figure S2. GO enrichment analysis of exclusive coding RNAs differently expressed in WT and *rag1^{+/−}* zebrafish during SVCV infection.

