

**SUPPLEMENTARY TABLE 2** Primer sequences used for reverse transcription-PCR.

<b>Gene name</b>	<b>NCBI Reference Sequence</b>	<b>Sequence (5' - 3')</b>
<i>NR3C1</i> <i>nuclear receptor subfamily</i> <i>3 group C member 1</i>	NM_001204264.1	TGAAATGGGCAAAGGCGATACC GGTCATACATGCAGGGTAGAGT
<i>NR3C2</i> <i>nuclear receptor subfamily</i> <i>3 group C member 2</i>	NM_000901.4	GTGTCCCAACAATTCTGGGCAG GACTCCACCTTGGGCAGCT
<i>UBE3A</i> <i>ubiquitin protein ligase E3A</i>	NM_130838.1	GGCTTTTCGGAGAGGTTTTCAT CAGAGTCCCTGGTATAGCCACC
<i>GILZ</i> <i>TSC22 domain family</i> <i>member 3 (TSC22D3)</i>	NM_198057.2	CAAGATTGAGCAGGCCATGGA TGGCTCTTCAGGGCTCAGC
<i>BDNF</i> <i>brain-derived neurotrophic</i> <i>factor</i>	NM_170735.5	TGAGTCTCCAGGACAGCAAAGC GTGGACGTTTACTTCTTTCATGG
<i>β-actin</i>	NM_001101.3	TTGCTGACAGGATGCAGAAG ACATCTGCTGGAAGGTGGAC
<i>GAPDH</i>	NM_002046.5	GTGGTCTCCTCTGACTTCAACA ACCACCCTGTTGCTGTAGCC