

SUPPLEMENTARY TABLE S3. SOMATIC FINDINGS AT FIRST CLINICAL EVALUATION AT THE CLINIC, DIFFERENTIATING AUTOIMMUNE DISEASE AND NONAUTOIMMUNE DISEASE GROUPS

<i>Somatic sign onset/presentation</i>	<i>Total cohort (n=45)</i>	<i>AD (n=26)</i>	<i>Non-AD (n=19)</i>	<i>Chi-square</i>	<i>Significance (p-value)</i>
Abnormalities in somatic assessment of:					
Heart	1/28 (4%) (n=28)	0/17 (0%) (n=17)	1/11 (9%) (n=10)	—	—
Lungs	0/27 (0%) (n=27)	0/18 (0%) (n=18)	0/9 (0%) (n=9)	—	—
Stomach	0/13 (0%) (n=13)	0/7 (0%) (n=7)	0/6 (0%) (n=6)	—	—
Joints	0/16 (0%) (n=16)	0/11 (0%) (n=11)	0/5 (0%) (n=5)	—	—
Ear, nose, and throat infection signs	12/26 (46%) (n=26)	8/17 (47%) (n=17)	4/9 (44%) (n=9)	0.02	0.899
Palatal petechiae	2/22 (10%) (n=22)	1/14 (7%) (n=14)	1/8 (13%) (n=8)	0.17	0.678
Skin abnormalities (dermatographia, livedo, severe dryness, redness, rash, or eczema)	11/18 (61%) (n=18)	6/11 (55%) (n=11)	5/7 (71%) (n=7)	0.52	0.477
Neurological abnormalities	7/31 (23%) (n=31)	3/20 (15%) (n=20)	4/11 (36%) (n=11)	1.79	0.185
Abnormalities in motor function	7/31 (23%) (n=31)	4/19 (21%) (n=19)	3/11 (27%) (n=11)	0.15	0.699
Chorea	1/41 (2%) (n=41)	1/24 (4%) (n=24)	0/17 (0%) (n=17)	—	—
Choreiform movements	6/26 (23%) (n=26)	3/17 (18%) (n=17)	3/9 (33%) (n=9)	0.79	0.373

AD, autoimmune disease; AD group, diagnosis of AD in the patient or first-degree relative; non-AD group, neither the patient nor first-degree relative has a diagnosis of AD.