

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Ambient Temperature and Markers of Fetal Growth: A Retrospective Observational Study of 29 Million U.S. Singleton Births

Shengzhi Sun, Keith R. Spangler, Kate R. Weinberger, Jeff D. Yanosky, Joseph M. Braun, and Gregory A. Wellenius

Table of Contents

Table S1. Summary statistics of average temperature during the entire pregnancy by county.

Table S2. Summary statistics of average temperature during the entire pregnancy, overall and by geographic region and climate zone.

Table S3. Pearson correlation between overall and trimester-specific temperature and fine particulate matter.

Figure S1. Location of the 403 US counties, 6 National Climate Assessment geographic regions, and 7 US Building America climate zones.

Figure S2. Flow diagram of birth exclusion in this study.

Figure S3. An illustration of how the gridded PRISM data intersect with census tracts in Providence county, RI.

Figure S4. Exposure-response curves for the associations between average temperatures during the entire pregnancy and small for gestational age (A) and birth weight z-score (B) among 31,921,046 births with gestational age ranging from 22 to 44 weeks in 403 US counties between 1989 and 2002. Models were adjusted for maternal age, race, marital status, years of education, smoking or drinking during pregnancy, parity, chronic hypertension, and year and month of conception.