

Table S5. Gene-specific primers used in qRT-PCR experiments:

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
β -actin	GGTACCACCATGTACCCAGG	GAAAGGGTGTAACGCAGC
CXCL13	CAGAATGAGGCTCAGCACAGC	CAGAATACCGTGGCCTGGAG
MADCAM-1	GAGCAAGAAGAGGAGATACAAGAG	TGGTGACCTGGCAGTGAAG
CCL21	AGACTCAGGAGCCCAAAGCA	GTTGAAGCAGGGCAAGGGT
CCL19	ACCAAGGACCAGAAGAGAGGA	ACTCACATCGACTCTCTAGG
Nfkb2	AATCTGGGTGTCCTGCATGTAA	AAGGAAAGCTGAGAAGCGTAGC
TNF	CATCTTCTCAAATTCGAGTGACAA	CCAGCTGCTCCTCCACTTG
MIP-2	AAAGTTTGCCTTGACCCTGA	TCCAGGTCAGTTAGCCTTGC
VCAM-1	TCGGGCGAAAAATAGTCCTT	CCGGCATATACGAGTGTGAA
OTUB1	GTAGCGACTCCGAAGGTGTT	ACCAGAGGATTCTGCACAGC