

| Phenotype | Figure | Sample Type | Treatment | Host Genotype (DGRP line) | | | | | |
|------------------------|--------|------------------------------------|--------------|---------------------------|-----|-----|-----|-----|-----|
| | | | | 437 | 491 | 566 | 589 | 748 | 892 |
| Larval encapsulation | 1A | Single larva dissection | G486 | 37 | 64 | 56 | 73 | 73 | 61 |
| | | | NSRef | 41 | 53 | 56 | 45 | 45 | 58 |
| Oil melanization | 1B | Single larva dissection | No Infection | 39 | 43 | 65 | 46 | 54 | 56 |
| | | | G486 | 38 | 27 | 50 | 21 | 52 | 30 |
| | | | NSRef | 15 | 9 | 41 | 16 | 40 | 34 |
| Hemocyte counts | 2 | Hemolymph pooled from 10-12 larvae | No Infection | 17 | 11 | 14 | 15 | 14 | 17 |
| | | | G486 | 11 | 5 | 6 | 8 | 10 | 8 |
| | | | NSRef | 13 | 5 | 10 | 11 | 10 | 7 |
| <i>Tep1</i> expression | 3A | RNA pooled from 10 larvae | No Infection | 8 | 8 | 8 | 8 | 8 | 8 |
| | | | G486 | 8 | 8 | 8 | 8 | 8 | 8 |
| | | | NSRef | 8 | 8 | 8 | 8 | 8 | 8 |
| PO activity | 3B | Hemolymph pooled from 20-25 larvae | No Infection | 12 | 10 | 12 | 8 | 12 | 11 |
| | | | G486 | 12 | 9 | 12 | 7 | 12 | 10 |
| | | | NSRef | 12 | 8 | 12 | 10 | 12 | 10 |
| Adult encapsulation | S1 | Groups of 40 animals | G486 | 8 | 5 | 8 | 8 | 5 | 3 |
| | | | NSRef | 8 | 5 | 8 | 8 | 5 | 4 |