

**Supplementary Data 1:** Results from the missing data analysis. Patients completing the survey in full, were more likely to have undergone surgery than those who did not ( $p = 0.04$ ). Data is presented as mean +/- standard deviation, unless specified as a percentage. Numbers within brackets indicate data points for the respective variable with incomplete data.

	<b>INCOMPLETE SURVEYS (N&lt;178)</b>	<b>COMPLETED SURVEYS (N=481)</b>	<b>P</b>
<b>%MALE</b>	16.3% (7/43)	29.1% (140/481)	.076
<b>AGE</b>	55.1 +/-10.9 (43)	53.6 +/- 9.8	.344
<b>%SURGERY</b>	35.9% (46/128)	45.9% (221/481)	.043
<b>LENGTH OF SYMPTOMS (YRS)</b>			.304
<b>0-1</b>	18.0% (23)	15.0% (72)	
<b>1-3</b>	21.9% (28)	29.1% (140)	
<b>3-10</b>	35.2% (45)	37.6% (181)	
<b>10-25</b>	18.8% (24)	15.4% (74)	
<b>25+</b>	6.3% (8)	2.9% (14)	
<b>LIMB PAIN VAS</b>	3.6 +/- 1.9 (17)	3.1 +/- 2.6	.854
<b>MJOA</b>	12.2 +/- 3.2 (47)	11.9 +/- 3.0	.364