

Table A.6.1. Estimated odds ratio, standard error, 95% confidence interval, and P-values for effects of spatial scale and publication year on tick-borne disease risk, assuming neighborhood extends 750 m from property boundaries.

<b>spatial scale</b>	<b>odds ratio</b>	<b>SE</b>	<b>CI</b>	<b>P</b>	<b>number of studies</b>	<b>number of records</b>
<b>Yard</b>	2.62	1.16	[1.97, 3.48]	<0.0001	13	59
<b>neighborhood</b>	4.1	1.29	[2.5, 6.72]	<0.0001	4	5
<b>outside neighborhood</b>	2.05	1.13	[1.61, 2.62]	<0.0001	8	33
<b>Year</b>	0.97	1.01	[0.96, 0.99]	<0.0001	NA	NA

Table A.6.2. Estimated odds ratio, standard error, 95% confidence interval, and P-values for effects of spatial scale and publication year on tick-borne disease risk, assuming neighborhood extends 250 m from property boundaries.

<b>spatial scale</b>	<b>odds ratio</b>	<b>SE</b>	<b>CI</b>	<b>P</b>	<b>number of studies</b>	<b>number of records</b>
<b>yard</b>	2.61	1.16	[1.96, 3.47]	<0.0001	13	59
<b>neighborhood</b>	4.08	1.29	[2.49, 6.69]	<0.0001	4	5
<b>outside neighborhood</b>	2.04	1.13	[1.6, 2.6]	<0.0001	9	34
<b>year</b>	0.97	1.01	[0.96, 0.99]	<0.0001	NA	NA