

Supplemental Table 4. aPL, APS and past thrombotic event prevalences according to the different RP subsets

Characteristics	No RP N=5	Non complicated RP N=96	Pitting scars N=19	Tip ulceration N=44	Gangrene N=4
LA, n (%)	1 (20)	7 (7)	2 (11)	6 (14)	1 (25)
aCL overall positivity, n (%)	0 (0)	8 (9)	3 (16)	2 (5)	1 (25)
IgG	-	7 (7)	1 (5)	1 (2)	1 (25)
IgM	-	1 (1)	1 (5)	1 (2)	0 (0)
Anti-β2GP1 overall positivity	0 (0)	4 (4)	1 (5)	7 (16)	2 (50)
IgG	-	1 (1)	0 (0)	5 (11)	2 (50)
IgM	-	3 (3)	1 (5)	3 (7)	1 (25)
APS, n (%)	0 (0)	3 (3)	1 (5)	1 (2)	2 (50)
Overall arterial thrombosis history, n (%)	0 (0)	2 (2)	1 (5)	1 (2)	1 (25)
Overall venous thrombosis history, n (%)	0 (0)	10 (10)	2 (11)	0 (0)	1 (25)

RP: Raynaud's phenomenon; LA: lupus anticoagulant; aCL: anticardiolipin; β2GP1: Beta2glycoprotein 1; APS: antiphospholipid syndrome