Procyanidin A1 Alleviates Inflammatory Response induced by LPS through NF-κB, MAPK, and Nrf2/HO-1 Pathways in RAW264.7 cells

Shan Han^{1,2#}, Hongwei Gao^{1,2#}, Shaoru Chen³, Qinqin Wang^{1,2}, Xinxing Li^{1,2}, Li-Jun Du⁴, Jun Li⁵, Ying-Ying Luo⁵, Jun-Xiu Li^{1,2}, Li-Chun Zhao^{1,2*}, Jianfang Feng^{1,2*}, Shilin Yang^{1,2} ¹ College of Pharmacy, Guangxi University of Chinese Medicine, Nanning 530000, China; ² Guangxi Engineering Technology Research Center of Advantage Chinese Patent Drug and Ethnic Drug Development, Nanning, 530020; ³ Department of Integrative Medical Sciences, Northeast Ohio Medical University, Rootstown, Ohio 44272, USA ⁴ School of Life Sciences, Tsinghua University, Beijing 100084, China; ⁵State Key Laboratory of Innovative Drug and Efficient Energy-Saving Pharmaceutical Equipment, Jiangxi University of Traditional Chinese Medicine, Nanchang 330004, China;

Running Title: Procyanidin A1 shows anti-inflammatory activity

Shan Han and Hongwei Gao equally contributed to this work.

*Co-corresponding Authors:

Dr. Li-Chun Zhao; Email: hyzlc@126.com; Address: College of Pharmacy, Guangxi University of Chinese Medicine, Nanning 530000, China; Tel: +86-771-4953513 Fax: +86-771-4953513 Dr. Jian-Fang Feng; Email:fengjianfang@vip.163.com; Address: College of Pharmacy, Guangxi University of Chinese Medicine, Nanning 530000, China; Tel: +86-771-4953513 Fax: +86-771-495351 Fax: +86-771-495351 Fax: +86-771-495351 Fax: +86-771-495351 Fax: +86-771-495351 Fax: +86-771 Fax: +86-771-495351 Fax: +86-771-49535







































Cytoplasm GAPDH





