

Supplemental Material

Data S1.

Supplemental Methods: Search Strategies

Database: MEDLINE (PubMed) (From 1946 to December 31st, 2018)

Search Strategy:

1. Stroke[MeSH] OR "brain ischemia"[MesH] OR stroke[tiab] OR "cerebrovascular accident"[tiab] (298198)
2. Thrombophilia[MeSH] OR thrombophili* [tiab] OR "inherited thrombophilia"[tiab] (28458)
3. Factor V"[MeSH] OR "Factor V Leiden"[tiab] OR "activated protein C resistance"[MeSH](8387)
4. Prothrombin[MeSH] OR prothrombin mutation [tiab] (10665)
5. Protein C deficiency[MesH] OR protein C deficiency[tiab] (1966)
6. Protein S deficiency[MeSH] OR protein S deficiency[tiab] (1704)
7. Antithrombin III Deficiency[MeSH] OR antithrombin deficiency[tiab] (1565)
8. 2 OR 3 OR 4 OR 5 OR 6 OR 7 (56664)
9. 1 AND 8 (2073)
10. Filters: Publication date to 2018/12/31; Humans; English; Adolescent: 13-18 years; Adult: 19+ year (1115)

MEDLINE (PubMed) Query:

```
((((("thrombophilia"[MeSH Terms] OR (thrombophilia[tiab] OR thrombophilia'[tiab] OR thrombophilia's[tiab] OR thrombophilic[tiab] OR thrombophilic[tiab] OR thrombophilicity[tiab] OR thrombophilics[tiab] OR thrombophilie[tiab] OR thrombophilies[tiab] OR thrombophilis[tiab]) OR "inherited thrombophilia"[tiab] OR "genetic polymorphisms"[tiab]) OR ("Factor V"[MeSH] OR "Factor V Leiden"[tiab] OR "activated protein C resistance"[MeSH])) OR ("prothrombin"[MeSH Terms] OR prothrombin mutation[tiab]) OR ("protein c deficiency"[MeSH Terms] OR "Protein C deficiency"[tiab])) OR ("protein s deficiency"[MeSH Terms] OR protein S deficiency[tiab])) OR ("antithrombin iii deficiency"[MeSH Terms] OR antithrombin deficiency[tiab])) AND ("stroke"[MeSH Terms] OR "brain ischemia"[MesH] OR stroke[tiab] OR "cerebrovascular accident"[tiab]) AND (("0001/01/01"[PDAT] : "2018/12/31"[PDAT]) AND "humans"[MeSH Terms] AND English[lang] AND ("adolescent"[MeSH Terms] OR "adult"[MeSH Terms]))
```

Database: EMBASE (From inception to December 31st, 2018)

Search Strategy:

1. 'cerebrovascular accident'/exp OR 'cerebrovascular accident':ti,ab OR 'brain ischemia'/exp OR 'brain ischemia':ti,ab OR 'stroke':ti,ab (513281)
2. 'thrombophilia'/exp OR 'thrombophili*':ti,ab OR 'inherited thrombophilia'/exp OR 'inherited thrombophilia':ti,ab OR 'genetic polymorphis*':ti,ab (45868)
3. 'blood clotting factor 5'/exp OR 'blood clotting factor 5 leiden'/exp OR 'factor v leiden':ti,ab OR 'activated protein c resistance'/exp OR 'activated protein c resistance':ti,ab (15829)
4. 'prothrombin'/exp OR (prothrombin NEAR/2 mutation):ti,ab (21764)
5. 'protein c deficiency'/exp OR 'protein c deficiency':ti,ab (3310)
6. 'protein s deficiency'/exp OR 'protein s deficiency':ti,ab (3066)
7. 'antithrombin deficiency'/exp OR 'antithrombin deficiency':ti,ab (2951)
8. 2 OR 3 OR 4 OR 5 OR 6 OR 7 (75736)
9. 1 AND 8 (4356)
10. #9 AND ([adolescent]/lim OR [adult]/lim OR [aged]/lim OR [middle aged]/lim OR [very elderly]/lim OR [young adult]/lim) AND [<1966-2018]/py AND [english]/lim NOT 'case report'/de (1300)

EMBASE Query:

('cerebrovascular accident'/exp OR 'cerebrovascular accident':ti,ab OR 'brain ischemia'/exp OR 'stroke':ti,ab) AND ('thrombophilia'/exp OR 'thrombophili*':ti,ab OR 'inherited thrombophilia'/exp OR 'inherited thrombophilia':ti,ab OR 'genetic polymorphis*':ti,ab OR 'blood clotting factor 5'/exp OR 'blood clotting factor 5 leiden'/exp OR 'factor v leiden':ti,ab OR 'activated protein c resistance'/exp OR 'activated protein c resistance':ti,ab OR 'prothrombin'/exp OR ((prothrombin NEAR/2 mutation):ti,ab) OR 'antithrombin deficiency'/exp OR 'antithrombin deficiency':ti,ab OR 'protein c deficiency'/exp OR 'protein c deficiency':ti,ab OR 'protein s deficiency'/exp OR 'protein s deficiency':ti,ab) AND ([adolescent]/lim OR [adult]/lim OR [aged]/lim OR [middle aged]/lim OR [very elderly]/lim OR [young adult]/lim) AND [<1966-2018]/py AND [english]/lim NOT 'case report'/de

Database: Cochrane Library (From 1946 to December 31st, 2018)**Search Strategy:**

1. Stroke[MeSH] OR "brain ischemia"[MesH] OR stroke:ti,ab,kw OR "cerebrovascular accident":ti,ab,kw (52117)
2. Thrombophilia[MeSH] OR thrombophili*:ti,ab,kw OR "inherited thrombophilia*":ti,ab,kw (700)
3. Factor V"[MeSH] OR "Factor V Leiden":ti,ab,kw OR "activated protein C resistance"[MeSH](178)
4. Prothrombin[MeSH] OR "prothrombin mutation":ti,ab,kw (461)
5. Protein C deficiency[MesH] OR "protein C deficiency":ti,ab,kw (44)
6. Protein S deficiency[MeSH] OR "protein S deficiency":ti,ab,kw (30)
7. Antithrombin III Deficiency[MeSH] OR "antithrombin deficiency":ti,ab,kw (48)
8. 2 OR 3 OR 4 OR 5 OR 6 OR 7 (1189)
9. 1 AND 8 (70)
10. 9 limit to December 2018 (54)

Table S1. Characteristics of Included Studies: Types of Thrombophilias, Numbers of Participants, and Study Population.

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Anadure 2017(1)	India	2010-2014	+					120/120	Hospitalized stroke ward patients, National Institute of Mental Health and Neuro Sciences, Bangalore Age 15-45 Partial or complete occlusion of common carotid, internal carotid and vertebral arteries/ anterior circulation strokes/ posterior circulation stroke/ occlusive disease of the large vessels of the brain on MRA or DSA Exclude: Hemorrhagic stroke; stroke due to trauma, infection, or tumors; vessel dissection; CNS vasculitis; malignancy or blood dyscrasia; aortoarteritis; nephrotic syndrome; cardioembolic stroke; vascular malformations and aneurysms; immunocompromised patients	Unrelated healthy subjects	CT/MRI/MRA	Age, sex
Aznar 2004(2)	Spain	NR	+	+				49/294	Age 18-50 Unit of thrombophilia Cryptogenic stroke without signs of atherosclerosis, heart disease, foramen ovale or vessel occlusive disease	Healthy subjects from the same geographic area and ethnic background	CT/MRI	Age, geographic area, ethnicity
Belvis 2006(3)	Spain	2001-2004	+	+				89/150	Stroke Unit, Hospital de la Santa Creu I Sant Pau, Barcelona First-ever cryptogenic stroke by TOAST criteria Exclude: Pregnancy and puerperal stage; TIA; strokes in other TOAST criteria	Previously published study in Barcelona population	CT/MRI	-
Bentolila 1997(4)	France	1993-1995		+				125/134	Age < 45 Non-transient arterial cerebral ischemia	Young healthy white men and women without history of thrombosis	CT	-
Biswas 2009(5)	India	NR	+	+	+	+		120/120	Outpatient Departments and Wards of the Departments of Neurosciences and Hematology, All India Institute of Medical Sciences, New Delhi Age <40 with acute ischemic stroke Stroke of non-cardioembolic origin Present within 4 weeks of onset Of Northern Indian origin Exclude: Cardioembolic stroke; past history of cardiovascular disease; on oral anticoagulants during the first sample collection; DM, hyperlipoproteinemia, cancer, sickle cell anemia, and liver disease	Apparently healthy hospital staffs and their relatives or from unrelated attendants of the patients Exclude: Taking medication; history of surgery or trauma in the past 30 days; History of bleeding, thrombotic, or cardiac disorders; pregnancy	CT/MRI	Age, sex

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Bolaman 2009(6)	Turkey	2003-2004	+	+				24/53	Department of Neurology, Adnan Menderes University, Aydin Stroke from cerebral infarct	Healthy subjects	CT/MRI	-
Buyru 2005(7)	Turkey	NR	+					29/20	Exclude: cerebral venous thrombosis; previous thromboembolic events; cerebrovascular thromboembolic and hemorrhagic event; TIA, drug use to influence the coagulation pathways; age <45 Neurology Clinic, Haydarpaşa Numune Hospital, Istanbul Ischemic stroke patients	Healthy subjects	CT/MRI	-
Catto 1995(8)	United Kingdom	NR	+					386/247	Four acute-care hospitals in Leeds Acute ischemic stroke by WHO definition	Leeds blood transfusion service and general practitioners	WHO definition	-
Celiker 2009(9)	Turkey	2000-2003	+	+				162/285	Neurology Department of Baskent University Hospital Acute ischemic stroke	Previously published study	MRI	-
Chatterjee 2013(10)	India	NR		+	+	+	+	52/52	Stroke Clinic, Department of Neurology, All India Institute of Medical Sciences, New Delhi Age <45 with non-embolic arterial ischemic stroke	Healthy individuals	CT/MRI	Age, sex
Chen 2003(11)	Taiwan	NR			+	+	+	104/35	Exclude: Cardioembolic, hemorrhagic stroke; systemic diseases (cancer, sickle cell anemia, vasculitis) Non-cardiac cerebral ischemia	Hospital employees, family of patients, or individuals asking for neurologic examination free of medical disorders	CT/MRI	-
Cushman 1998(12)	USA	1989-1990	+					149/482	Exclude: SLE; APS; collagen diseases; taking anticoagulants and hormonal drugs; TIA Cardiovascular Health Study (Random sample of Medicare eligibility list) Age ≥65 Development of stroke in participants free of baseline history of stroke	Cardiovascular Health Study	Review by committee	-
D'Amico 1998(13)	Italy	1996-1997	+		+	+		31/124	C. Besta Neurological Institute, Milan and Ospedale L. Mandic Age <45 with ischemic stroke	Healthy volunteer from staff Age <45	CT/MRI	-
De Lucia 1999(14)	Italy	1994-1995	+		+	+	+	50/100	Ischemic stroke	Healthy subjects	WHO definition	Age, sex
Djordjevic 2012(15)	Serbia	NR	+	+				73/120	Young adults having cerebral infarcts	Healthy blood donors	MRI	-
Egan 2000(16)	USA	1997-1998		+				42/635	Oregon Health Sciences University Hospital Age ≤55 with an arterial stroke	Normal healthy individuals from Portland	CT/MRI	-
Erten 2015(17)	Turkey	2007-2009	+	+				212/238	Research and Training Hospital Neurology Clinic, Süleyman Demirel University Ischemic stroke patients	Individuals without history of stroke	NR	Age, Sex

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Eterovic 2007(18)	Croatia	1999-2003	+	+				120/120	Department of Neurology, Clinical Hospital Split First-time acute ischemic stroke Exclude: Age >65, secondary hypercoagulability status, DM type 1, significant obstruction of carotid arteries	Persons attending regular checkups, blood donors, volunteer staffs without cerebrovascular disease	CT/MRI	Age, Sex, cardiovascular risk factors
Fan 2010(19)	USA	1991-1994	+	+				156/5817	Second phase of The Third National Health and Nutrition Examination Survey (NHANES III) Age ≥17 with self-reported stroke	Same population without self-reported stroke	Interview	-
Favaretto 2012(20)	Italy	2008-2011	+	+	+	+	+	340/272	Angiology Unit, S. Orsola-Malpighi University Hospital Stroke of unknown cause by TOAST criteria Exclude: Large vessel, small vessel, cardioembolic stroke; stroke from multiple etiologies; neoplasia, dementia	Out-patients referred for varicose veins and/or early venous insufficiency in the legs without clinical indication of stroke Exclude: ABI≤0.9; previous coronary event; neoplasia; DVT	CT/MRI	-
Go 2003(21)	USA	1996-1997	+					137/214	Cohort from Anticoagulation and Risk Factors in Atrial Fibrillation (ATRIA) study Ambulatory adults with nonvalvular AF Patients Ischemic stroke	Same cohort with no prior hospitalization for ischemic stroke	ICD-9 with record review	Follow-up time
Haeusler 2012(22)	Germany	NR	+	+				44/282	Stroke outpatient clinic Cryptogenic stroke by TOAST criteria Exclude: Age ≥55	Healthy blood donors without vascular diseases	CT/MRI	-
Halbmayer 1998(23)	Austria	NR	+	+				20/20	Unexplained juvenile stroke	Healthy subjects	NR	Age, sex
Hamedani 2013(24)	USA	1992-1996 2001-2003 2003-2007	+					830/907	Genetics of Early Onset Stroke (GEOS) study First ischemic stroke Age 15-49 Exclude: trauma; procedure; hemorrhage; CVST; infection; vasculitis	Baltimore-Washington area Participants without history of stroke	Record review	Age, region of residence
Hankey 2001(25)	Australia	1996-1998	+	+	+	+	+	219/205	A university teaching hospital in Western Australia First-ever ischemic stroke	Randomly selected from the Western Australian electoral roll	CT/MRI	Age, sex, postal code
Jerrard-Dunne 2003(26)	England	NR			+	+	+	130/130	Stroke services in South London Age ≤65 Acute ischemic stroke	Sampling of primary care lists in the same geographic area Stroke-free individuals	CT/MRI	Age, sex, ethnicity
Jiang 2014(27)	USA	1992-1996 2001-2003 2003-2007		+				397/426	Genetics of Early Onset Stroke (GEOS) study First ischemic stroke Age 15-49 Exclude: trauma; procedure; hemorrhage; CVST; infection; vasculitis	Baltimore-Washington area Participants without history of stroke	Record review	Age, region of residence

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Juul 2002(28)	Denmark	1991-1997	+					641/7907	The Copenhagen City Heart Study and Copenhagen University Hospital Age 20-95 Ischemic stroke	Control subjects from the Copenhagen City Heart Study free of MI, IS, and non-MI IHD	CT	-
Kamberi 2016(29)	Macedonia	2008-2010	+	+				39/102	Neurology Department, Clinical Hospital, Tetovo Age 18-90 First-ever ischemic stroke Exclude: Renal failure; MI; recent inflammatory or immunologic disease.	Healthy subjects from local residents	CT/MRI	-
Karakus 2005(30)	Turkey	NR	+	+	+	+	+	21/81	Age < 50 Cerebral infarction	Volunteer physicians and laboratory staff	CT/MRI	-
Karttunen 2003(31)	Finland	1991-1998	+	+	+	+	+	58/104	Age 15-60 Ischemic brain infarction of undetermined cause and PFO	Spouse, friend, or randomly selected controls from the population register of the hospital catchment area	CT	Age, sex
Kholodkova 2015(32)	Ukraine	NR	+	+				122/40	Neurology Unit, Kyiv City Hospital Acute ischemic stroke	Healthy donors without previous history of stroke	CT/MRI	-
Krajcoviechova 2015(33)	Czech Republic	2009-2012	+	+				423/614	Thomayer Hospital or Charles University Hospital Age 18-81 First-ever acute ischemic stroke	Participants of the Czech post-MONICA study residing in Prague East and Pilsen districts Age 50-75 Free of vascular diseases	CT/MRI	-
Kumar 2017(34)	India	NR	+					250/250	All India Institute of Medical Sciences, New Delhi Age 18-85, North Indian Ischemic stroke within three years before recruitment	Spouses, relatives or patients attending neurology department for treatment other than stroke Age 18-85, North Indian No prior stroke	CT	Age, sex
Linnemann 2008(35)	Germany	2000-2006	+	+	+	+	+	41/993	Patients with history of VTE registered in the MAISTHRO Database Age 17-90 with ischemic stroke	Same cohort without stroke	CT/MRI	-
Longstreth 1998(36)	USA	1991-1995	+	+				41/382	King, Pierce, Snohomish counties, Washington Women age 18-44 Diagnosed of first stroke	Random-digit dialing Same area Healthy women age 18-44	CT/MRI	Age
Lopaciuk 2001(37)	Poland	1996-1999	+	+				100/238	Age ≤45 History of ischemic stroke without a cardiac embolic source	Healthy blood donor and hospital staff	CT/MRI	-
Martinelli 2006(38)	Italy	1994-2005	+	+				105/293	Referred for thrombophilia screening at Thrombosis Center, University of Milan and IRCCS Maggiore Hospital Caucasian women of fertile age	Partners and friends of patients	CT/MRI	-

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Mayer 1993(39)	USA	1990-1991				+		94/94	First ischemic stroke Columbia-Presbyterian Medical Center North Manhattan Stroke Study Age >39 Residing in the Zip code of northern Manhattan	Caucasian women of fertile age without previous thrombosis Patients admitted to the medicine and neurology services Age >39 in same area Exclude: Thrombosis; bleeding; renal/liver failure; malignancy; infection	NR	-
Mochan 2005(40)	South Africa	NR				+		33/33	Chris Hani Baragwanath Hospital in Soweto HIV-infected Stroke with cerebral infarction	Inpatients of same hospital HIV-infected No cerebral infarction	CT	Age, sex, CD4 count
Moskau 2010(41)	Germany	1999-2000	+	+				167/500	Of German descent Ischemic stroke	Blood donors of German descent	CT/MRI	-
Nagayama 1996(42)	Japan	NR	+					106/37	Chronic ischemic stroke	Patients with neurological diseases without vascular involvement	NR	-
Pahus 2016(43)	Denmark	2004-2012	+	+	+	+	+	377/64631	Center of Hemophilia and Thrombosis, Aarhus University Hospital Age 18 - 50 referred to thrombophilia investigation Ischemic stroke	Previously published data of general western population	WHO and record review	-
Pestana 2009(44)	Venezuela	2005-2007	+					54/134	Banco Municipal de Sangre, Caracas Stroke	Randomly selected, unrelated, and apparently healthy subjects without personal and family history of vascular, arterial, or thromboembolic diseases	CT	-
Petrovic 2003(45)	Slovenia	NR	+					96/115	Acute cerebral infarction	General population in the same region Exclude: History of CVD, CHD, PAD; carotid bruit	CT/MRI	-
Pezzini 2005(46)	Italy	1997-2002	+	+				163/158	Department of Neurology, University of Brescia, Brescia Age <45 Admitted patients with first-ever ischemic stroke	Staff of same hospital No vascular diseases	CT/MRI	Age, sex
Pezzini 2007(47)	Italy	NR	+	+				108/216	Department of Neurology, University of Brescia, Brescia Age <45 Unselected, unrelated women admitted first-ever acute ischemic stroke	Women from the staff of same hospital	CT/MRI	Age, sex

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Press 1996(48)	USA	NR	+					161/367	Wards and clinics at Portland Veterans Affairs Medical Center and Oregon Health Sciences University Acute ischemic stroke within 7 days of enrollment	Healthy elderly (N=54) Stroke risk group (N=116) Blood donors (N=197)	NR	-
Pullmann 2004(49)	Slovakia	NR	+	+				23/71	White SLE patients with thromboembolic stroke Exclude: Hemorrhage; vasculitis	SLE patients without CVA/CHD	CT/MRI	-
Ranellou 2015(50)	Greece	NR	+	+				51/70	Evangelismos General Hospital Age 18-50 Ischemic stroke within 24 hour of onset Native Greek Exclude: Major systemic diseases; coagulopathy; anticoagulants; trauma	Healthy blood donors from the same area without history of stroke/thrombosis	CT/MRI	Age
Ridker 1995(51)	USA	NR	+					209/704	Physicians' Health Study Apparently healthy US male physicians Age 40-84 Developed stroke during 10-year follow-up	Randomly selected participants from the same study No cardiovascular disease at the time of event in cases	CT and record review	Age, smoking status, time since randomization in study
Ridker 1999(52)	USA	NR		+				259/1774	Physicians' Health Study Apparently healthy US male physicians Age 40-84 Developed stroke during 10-year follow-up	Randomly selected participants from the same study No cardiovascular disease during follow up	CT and record review	Age, smoking status
Ripoll 1997(53)	French	NR	+					321/428	Age >65 Ischemic cerebrovascular events	Age >65 No personal or familial history of CVA/CHD	NR	-
Romdhane 2011(54)	Tunisia	NR			+	+	+	20/54	Age <50 First non-cardioembolic ischemic stroke	NR	NR	Age
Rubattu 2005(55)	Italy	1998-2003	+	+				294/286	Neurological Department, University of Sassari, Sardinia Ischemic stroke	Randomly selected patients admitted to the same hospital with vascular risk factors or history of cardiovascular disease Exclude: Current or previous CVA	CT/MRI	-
Rubattu 2005(56)	Italy	NR	+	+				115/180	Neurological Institute, University La Sapienza, Rome Age 15-45 First-ever ischemic stroke within the 8 weeks preceding the admission into the hospital	Healthy blood donors from the same center No drug/OCP use No family history of stroke	CT/MRI	Age
Sastry 2006(57)	United Kingdom	1993-1998	+	+	+	+	+	101/101	Hospitals in the North West and Mersey Regions, Manchester Age 16-39 at the time of stroke First ischemic stroke identified by ICD-9	Age 16-39 Resides in Manchester Exclude: History of cardiovascular	WHO and record review	Age, sex

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Shi 2008(58)	China	2006-2007	+					97/99	Exclude: Surgery/injury within 10 weeks; malignancy Neurological department, Beijing Tiantan Hospital Age 18-45 Acute ischemic stroke	disease; stroke; arterial thrombosis Other departments of the Tiantan Hospital Age 18-45 No cardiovascular or cerebrovascular diseases	CT/MRI	Age, sex
Slooter 2005(59)	Netherlands	1990-2001	+	+				193/767	Exclude: AF; hemorrhage Nine participating Dutch hospitals University Medical Center Utrecht Women age 18-49 Hospitalized for a first ischemic stroke	Random-digit dialing Women age 18-49 No history of CHD/CVD/PAD	CT/MRI	Age, residence, year of stroke
Smiles 2002(60)	USA	NR		+				182/453	Exclude: TIA; hemorrhagic stroke; CSVT; dissection; history of CHD/CVD; terminal illness; aphasia or cognitive impairment Cardiovascular Health Study (Random sample of Medicare eligibility list) Age ≥65 (free of clinical CVD at baseline) Had stroke during 6-year follow up	Cardiovascular Health Study Free of subclinical CVD group African-American group Population-based controls	CT	-
Supanc 2014(61)	Croatia	2009-2012	+	+				155/150	Department of Neurology, Sestre milosrdnice University Hospital Center, Zagreb Ischemic stroke Age <55	Subjects treated at Pain Clinic of same center No history of vascular or thromboembolic disease	CT/MRI	Age, sex
Szolnoki 2003(62)	Hungary	1998-2002	+	+				867/743	Department of Neurology and Neurophysiology, Pandy Kalman County Hospital Consecutive Hungarian patients First acute ischemic stroke	Randomly selected from local GP registers Healthy Caucasian Hungarian No evidence of stroke on CT/MRI	MRI	Age, sex
Tatarskyy 2010(63)	Ukraine	2008-2009	+	+				183/188	Randomly selected from different regions in Ukraine Survivor of ischemic stroke referred for rehabilitation	I: General population of Ukraine II: Healthy individuals age >65 without history of ischemic stroke	NR	-
They-They 2012(64)	Morocco	2008-2009	+	+				91/182	University Hospital Center, Casablanca Admitted for ischemic stroke	Presumably healthy blood donors	CT/MRI	Age, sex, ethnicity
Tupitsyna 2013(65)	Russia	NR	+	+				1450/817	Stroke patients from Russian and Ukrainian population	Russian and Ukrainian population	NR	-
Voetsch 2000(66)	Brazil	1996-1998	+	+				153/225	University Hospitals of the State University of Campinas and the University of Sao Paulo First cerebral ischemic event occurring at age 15-45 Absence of systemic disease or cancer	Randomly selected hospital staffs	CT/MRI	Age, sex

Reference	Country	Study period	Types of thrombophilia tested					Number of cases/controls	Study population	Control Identification	Stroke diagnosis	Case-control matching for
			FVL	PTM	PCD	PSD	ATD					
Wypasek 2009(67)	Poland	NR	+	+				100/107	Exclude: PCD, PSD, ATD; Antiphospholipid antibodies PFO patients Age <65 History of first-ever cryptogenic ischaemic stroke after a median of 8 (3–19) months before enrollment	Apparently healthy white individuals	CT/MRI	Age, sex
Zimba 2017(68)	Zambia	2014-2015			+	+		52/52	Exclude: Inflammatory states; cancer; AF; DM; valvular heart disease; CAD; carotid artery stenosis In-patients and Out-patients at University Teaching Hospital, Lusaka Age ≥18 HIV positive Ischemic stroke	Same center HIV positive patients without ischemic stroke	CT/MRI	Age, sex, ethnicity

NR, not reported; FVL, Factor V Leiden; PTM, Prothrombin G20210A Mutation; PCD, Protein C Deficiency; PSD, Protein S Deficiency; ATD, Antithrombin Deficiency; CT, computed tomography; MRI, Magnetic Resonance Imaging; MRA Magnetic Resonance Angiogram; TIA, transient ischemic attack; DM, Diabetes Mellitus; WHO, World Health Organization; SLE, systemic lupus erythematosus; APS, antiphospholipid syndrome; PFO, patent foramen ovale; VTE, venous thromboembolism; CVD, cerebrovascular disease; CHD, coronary heart disease; CVA, cerebrovascular accidents; AF, atrial fibrillation; OCP, oral contraceptive pills; PAD; peripheral arterial disease; CSVt, cerebral venous sinus thrombosis; HIV, human immunodeficiency virus.

Table S2. Characteristics of Included Studies: Demographic Data of Cases and Controls.

References	Mean Age, year Cases/Controls	Male, % Cases/Controls	Cryptogenic stroke in cases, %	Ethnicity, % Cases/Controls	Diabetes, % Cases/Controls	Hypertension, % Cases/Controls	Dyslipidemia, % Cases/Controls	Hormonal Drug Use, % Cases/Controls	Smoking, % Cases/Controls
Anadure 2017(1)	33/35	93/93	NR	NR/NR	NR/NR	13/NR	33/NR	NR/NR	63/43*
Aznar 2004(2)	18-50 (Range)/NR	NR/NR	100	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Belvis 2006(3)	57/48	60/45	100	NR/NR	17/2.7	36/13	38/14	6/NR	51/37
Bentolila 1997(4)	41/34	58/50	44	White 100/100	NR/NR	NR/NR	NR/NR	25/NR	NR/NR
Biswas 2009(5)	NR/NR	NR/NR	NR	Indian 100/100	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Bolaman 2009(6)	64/59	67/58	NR	NR/NR	33/32	63/60	38/36	NR/NR	21/19
Buyru 2005(7)	67/61 (Median)	NR/NR	NR	NR/NR	21/NR	59/NR	NR/NR	NR/NR	10/NR
Catto 1995(8)	74/76 (Median)	NR/NR	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Celiker 2009(9)	70/NR	54/NR	30	NR/NR	27/NR	65/NR	39/NR	NR/NR	NR/NR
Chatterjee 2013(10)	23/NR	63/63	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Chen 2003(11)	62/61	68/69	NR	NR/NR	28/0	70/0	NR/NR	NR/NR	NR/NR
Cushman 1998(12)	76/72	48/37	NR	NR/NR	39/16	60/34	NR/NR	NR/NR	45/50
D'Amico 1998(13)	34/33	65/32	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
De Lucia 1999(14)	35/NR	76/NR	NR	NR/NR	NR/NR	NR/NR	54/NR	0/NR	NR/NR
Djordjevic 2012(15)	40/39	56/70	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Egan 2000(16)	43/41	52/49	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Erten 2015(17)	66/62*	52/45	51	NR/NR	30/24	61/43*	28/23	NR/NR	38/37
Eterovic 2007(18)	61/61 (Median)	64/64	NR	White 100/ 100	30/30	56/56	32/32	NR/NR	38/38
Fan 2010(19)	68/44*	44/48	NR	White 82/82 Black 15/12 Hispanic 3/6	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Favaretto 2012(20)	51/52.2	PFO 49 Non- PFO 46 /74	100	White 100/100	PFO 5 Non- PFO 3 /4	PFO 14 Non- PFO 13 /20	PFO 19 Non- PFO 23 /21.7	PFO 12 Non- PFO 12 /0	PFO 21 Non- PFO 24 /17
Go 2003(21)	74/69*	51/62*	NR	White 80/84 Black 6/2 Hispanic 4/2 Native American 1/0 Others 4/9	20/14	66/44*	NR/NR	NR/NR	NR/NR
Haeusler 2012(22)	36/39 (Median)	41/36	100	NR/NR	5/NR	223/NR	36/NR	NR/NR	40.9/NR
Halbmayer 1998(23)	39/39	50/50	100	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Hamedani 2013(24)	41/40*	57/55	50.4	White 56/60 Black 44/40	17/5*	43/19*	NR/NR	16/8*	41/28*
Hankey 2001(25)	66/67	64/64	20	NR/NR	25/11*	54/33*	24/22	NR/NR	33/18*
Jerrard-Dunne 2003(26)	White 53/54 Black Caribbean 57/56 Black African 53/55	62/50	NR	White 39/39 Black Caribbean 39/39 Black African 23/23	26/10	59/39	49/42	NR/NR	40/23
Jiang 2014(27)	41/39*	63/55*	NR	White 100/100	11/2*	32/16*	NR/NR	23/11*	43/24*
Juul 2002(28)	63/56*	61/43*	NR	White >99/>99	14/3*	46/17*	NR/NR	NR/NR	80/73*
Kamberi 2016(29)	63/49*	NR/NR	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Karakus 2005(30)	40/42	38/52	100	NR/NR	10/5	33/1*	43/43	NR/NR	38/22
Karttunen 2003(31)	44/45	55/57	100	NR/NR	2/4	16/13	NR/NR	27/27	29/24
Kholodkova 2015(32)	73/NR	NR/NR	0	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Krajcoviechova 2015(33)	66/61*	61/49*	36	NR/NR	30/10*	87/58*	93/86*	NR/NR	60/53*

References	Mean Age, year Cases/Controls	Male, % Cases/Controls	Cryptogenic stroke in cases, %	Ethnicity, % Cases/Controls	Diabetes, % Cases/Controls	Hypertension, % Cases/Controls	Dyslipidemia, % Cases/Controls	Hormonal Drug Use, % Cases/Controls	Smoking, % Cases/Controls
Kumar 2017(34)	51/53	81/81	NR	North Indian 100/100	32/10*	58/17*	23/6*	NR/NR	39/27*
Linnemann 2008(35)	66/48 (Median)*	56/38*	NR	NR/NR	15/6*	56/25*	46/19*	NR/NR	15/21
Longstreth 1998(36)	37/38	0/0	NR	White 83/90 Black 6/2 Others 11/8	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Lopaciuk 2001(37)	38/33	51/66	55	NR/NR	20/NR	42/NR	35/NR	NR/NR	61/NR
Martinelli 2006(38)	35/35	0/0	52	White 100/100	NR/NR	12/3*	10/0*	NR/NR	27/21
Mayer 1993(39)	68/68	48/40	17.1	White 14/21 Black 38/37 Hispanic 46/39 Others 2/2	30/18	75/46*	NR/NR	NR/NR	48/36
Mochan 2005(40)	NR/NR	NR/NR	NR	Black 100/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Moskau 2010(41)	55/33*	38/29*	17	White 100/100	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Nagayama 1996(42)	40/55	63/18	NR	Japanese 100/100	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Pahus 2016(43)	43/NR	51/NR	NR	NR/NR	8/NR	32/NR	37/NR	22/NR	50/NR
Pestana 2009(44)	39/39	39/39	NR	NR/NR	2/2	13/16	NR/NR	13/10	14/21
Petrovic 2003(45)	62/63	63/57	NR	White 100/100	31/4*	66/33*	NR/NR	NR/NR	44/25*
Pezzini 2005(46)	35/35	52/54	31	White 100/100	3/3	17/6*	28/20*	NR/NR	47/25*
Pezzini 2007(47)	34/35	0/0	38	White 100/100	0/2	12/4*	25/11*	40/14	41/13*
Press 1996(48)	64/68	91/80	NR	NR/NR	36/21	72/64	26/30	NR/NR	41/10
Pullmann 2004(49)	37/46	4/1	NR	White 100/100	18/16	41/35	51/37	NR/NR	24/25
Ranellou 2015(50)	37/38	49/54	NR	NR/NR	2/0	16/0*	10/3	NR/NR	35/36
Ridker 1995(51)	63/60	100/100	NR	Predominantly white/Predominantly white	13/4	36/17	12/9	NR/NR	60/58
Ridker 1999(52)	60/59	100/100	NR	Predominantly white/Predominantly white	7/3*	28/16*	12/9*	NR/NR	56/57
Ripoll 1997(53)	66/NR	NR/NR	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Romdhane 2011(54)	37/NR	55/NR	NR	NR/NR	NR/NR	<1/NR	NR/NR	<1/NR	<1/NR
Rubattu 2005(55)	75/73 (Median)*	60/57	NR	NR/NR	24/29	65/49*	22/13*	NR/NR	41/38
Rubattu 2005(56)	36/35	44/54	12	NR/NR	2/0	22/4*	18/18	44/Excluded	57/49
Sastry 2006(57)	33/33	43/43	NR	NR/NR	3/1	21/8*	NR/NR	16/16	47/41
Shi 2008(58)	39/39	81/81	NR	Asian 100/100	20/5*	50/13*	NR/NR	NR/NR	72/42*
Slooter 2005(59)	39/40	0/0	0	NR/NR	4/1*	32/6*	8/3*	52/36*	35/32
Smiles 2002(60)	76/72	41/39	NR	White 95/94 Black 5/5 Other 0/5	24/14	53/34	NR/NR	NR/NR	45/55
Supanc 2014(61)	NR/NR	NR/NR	36	White 100/100	4/3	37/19*	63/53	NR/NR	45/24*
Szolnoki 2003(62)	61/60	53/53	NR	White 100/100	32/6*	51/18*	NR/NR	NR/NR	33/11*
Tatarskyy 2010(63)	65/30*	52/45	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
They-They 2012(64)	49/46	51/52	6.6	NR/NR	18/9*	51/10*	NR/NR	NR/NR	22/7*
Tupitsyna 2013(65)	NR/NR	NR/NR	NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR	NR/NR
Voetsch 2000(66)	33/34	41/44	NR	White 75/53 Black 25/47	5/NR	39/NR	31/NR	NR/NR	52/NR
Wypasek 2009(67)	43/44	30/32	NR	White 100/100	NR/NR	NR/NR	NR/NR	NR/NR	19/34

References	Mean Age, year Cases/Controls	Male, % Cases/Controls	Cryptogenic stroke in cases, %	Ethnicity, % Cases/Controls	Diabetes, % Cases/Controls	Hypertension, % Cases/Controls	Dyslipidemia, % Cases/Controls	Hormonal Drug Use, % Cases/Controls	Smoking, % Cases/Controls
Zimba 2017(68)	52/46*	44/44	13	NR/NR	15/6	50/17*	8/8	Excluded/Excluded	4/4

* denotes significant difference; NR, not reported

Table S3. Results of Included Studies: Factor V Leiden.

References	Test Method	Odds Ratio (95%CI)	Number of All FVL (%)		Number of Homozygous FVL (%)		Number of Heterozygous FVL (%)	
			Cases	Controls	Cases	Controls	Cases	Controls
Anadure 2017(1)	PCR/RFLP	3.10 (0.61-15.7; <i>P</i> = .15)	6/120 (5)	2/120 (1.7)	0/120 (0)	0/120 (0)	6/120 (5)	2/120 (1.7)
Aznar 2004(2)	PCR/RFLP	2.62 (0.43-13.95)	2/49 (4.1)	5/294 (1.7)				
Belvis 2006(3)	PCR	NR	2/89 (2.2)	6/150 (4)	0/89 (0)	0/150 (0)	2/89 (2.2)	6/150 (4)
Biswas 2009(5)	PCR/RFLP	10.8 (1.3-229.5; <i>P</i> = .005)	10/120 (8.3)	1/120 (0.8)	4/120 (3.3)	0/120 (0)	6/120 (5)	1/120 (0.8)
Bolaman 2009(6)	PCR	0.431 (0.074-2.504; <i>P</i> > .05)	2/24 (8.3)	2/53 (3.8)	0/24 (0)	0/53 (0)	2/24 (8.3)	2/53 (3.8)
Buyru 2005(7)	PCR/RFLP	NR	1/29 (3.4)	0/20 (0)	1/29 (3.4)	0/20 (0)	0/29 (0)	0/20 (0)
Catto 1995(8)	PCR/RFLP	NR	16/386 (4.1)	14/247 (5.7)	0/386 (0)	0/247 (0)	16/386 (4.1)	14/247 (5.7)
Celiker 2009(9)	PCR/RFLP	NR	21/162 (13)	28/285 (9.8)	1/162 (0.6)	0/285 (0)	20/162 (12.3)	28/285 (9.8)
Cushman 1998(12)	PCR	0.76 (0.32-1.81)	8/149 (5.4)	34/482 (7.1)	0/149 (0)	0/482 (0)	8/149 (5.4)	34/482 (7.1)
D'Amico 1998(13)	PCR	NR	5/31 (16.1)	4/124 (3.2)	0/31 (0)	0/124 (0)	5/31 (16.1)	4/124 (3.2)
De Lucia 1999(14)	PCR	NR	11/50 (22)	2/100 (2)	2/50 (4)	0/100 (0)	9/50 (18)	2/100 (2)
Djordjevic 2012(15)	PCR/RFLP	1.45 (0.47-4.48)	6/73 (8.2)	7/120 (5.8)				
Erten 2015(17)	PCR/in situ hybridization	NR	32/212 (15.1)	21/238 (8.8)	3/212 (1.4)	0/238 (0)	29/212 (13.7)	21/238 (8.8)
Eterovic 2007(18)	PCR/RFLP	NR (<i>P</i> = .023)	10/120 (8.3)	3/120 (2.5)	0/120 (0)	0/120 (0)	10/120 (8.3)	3/120 (2.5)
Fan 2010(19)	PCR	0.82 (0.29-2.34)	6/156 (3.8)	262/5817 (4.5)				
Favaretto 2012(20)	PCR	NR	16/340 (4.7) PFO: 3/136 (2.2) Non-PFO: 13/204 (6.4)	14/272 (5.1)	0/340 (0) PFO: 0/136 (2.2) Non-PFO: 0/204 (6.4)	0/272 (0)	16/340 (4.7) PFO: 3/136 (2.2) Non-PFO: 13/204 (6.4)	14/272 (5.1)
Go 2003(21)	PCR/RFLP	Crude OR 1.59 (0.58-4.36) Adjusted OR for age and prior ischemic stroke: 1.81 (0.65-5.05)	8/137 (5.8)	8/214 (3.7)	0/137 (0)	0/214 (0)	8/137 (5.8)	8/214 (3.7)
Haeusler 2012(22)	PCR	NR (<i>P</i> = .073)	7/41 (17.1)	22/282 (7.8)	0/41 (0)	0/282 (0)	7/41 (17.1)	22/282 (7.8)
Halbmayer 1998(23)	PCR	NR (<i>P</i> = 1.00)	1/20 (5)	2/20 (10)	0/20 (0)	0/20 (0)	1/20 (5)	2/20 (10)
Hamedani 2013(24)	SNP array	1.00 (0.60-1.66; <i>P</i> = 1.00)	30/830 (3.6)	34/907 (3.7)				
Hankey 2001(25)	PCR/RFLP	2.1 (0.6-6.8)	10/219 (4.6)	4/205 (2)				
Juul 2002(28)	PCR/RFLP	Age-adjusted OR 0.92 (0.56-1.53)	17/231 (7.4)	629/7907 (8)	1/231 (0.4)	17/7907 (0.2)	16/231 (6.9)	612/7907 (7.7)
Kamberi 2016(29)	PCR/hybridization	1.15 (0.33-4.02; <i>P</i> = .83)	3/39 (7.7)	9/102 (8.8)	0/39 (0)	0/102 (0)	3/39 (7.7)	9/102 (8.8)
Karakus 2005(30)	PCR	NR	3/21 (14.3)	4/81 (4.9)	0/21 (0)	1/81 (1.2)	3/21 (14.3)	3/81 (3.7)
Karttunen 2003(31)	PCR	7.8 (0.8-71.3)	4/57 (7)	1/104 (1)	0/57 (0)	0/104 (0)	4/57 (7)	1/104 (1)
Kholodkova 2015(32)	PCR	NR	3/114 (2.6)	0/40 (0)	0/114 (0)	0/40 (0)	3/114 (2.6)	0/40 (0)
Krajcoviechova 2015(33)	PCR	Multivariate adjusted OR 1.22 (0.74-2.00; <i>P</i> =.43)	44/423 (10.4)	53/614 (8.6)	2/423 (0.5)	2/614 (0.3)	42/423 (9.9)	51/614 (8.3)
Kumar 2017(34)	PCR	1.80 (0.60-5.37; <i>P</i> =.29)	9/250 (3.6)	5/250 (2)	0/250 (0)	0/250 (0)	9/250 (3.6)	5/250 (2)
Linnemann 2008(35)	PCR	0.87 (0.42-1.79; <i>P</i> =.86)	10/41 (24.4)	278/1020 (27.3)	0/41 (0)	22/1020 (2.2)	10/41 (24.4)	256/1020 (25.1)
Longstreth 1998(36)	PCR/RFLP	0 (0-2.5)	0/40 (0)	16/388 (4.1)	0/40 (0)	0/388 (0)	0/40 (0)	16/388 (4.1)
Lopaciuk 2001(37)	PCR/RFLP	0.7 (0.2-2.6)	3/100 (3)	10/238 (4.2)	0/100 (0)	0/238 (0)	3/100 (3)	10/238 (4.2)
Martinelli 2006(38)	NR ("DNA analysis")	Crude OR 2.5 (0.8-7.5) Adjusted OR for age 2.6 (0.8-8.0)	6/105 (5.7)	7/293 (2.4)	0/105 (0)	0/293 (0)	6/105 (5.7)	7/293 (2.4)
Moskau 2010(41)	PCR	NR (<i>P</i> = .35)	11/167 (6.6)	30/500 (6)	1/167 (0.6)	1/500 (0.2)	10/167 (6)	29/500 (5.8)
Nagayama 1996(42)	PCR/RFLP	NR	0/106 (0)	0/37 (0)	0/106 (0)	0/37 (0)	0/106 (0)	0/37 (0)

References	Test Method	Odds Ratio (95%CI)	Number of All FVL (%)		Number of Homozygous FVL (%)		Number of Heterozygous FVL (%)	
			Cases	Controls	Cases	Controls	Cases	Controls
Pahus 2016(43)	PCR	Homozygous 4.06 (0.86-36.51) Heterozygous 1.02 (0.54-1.79)	15/207 (7.2)	282/4188 (6.7)	1/207 (0.5)	5/4188 (0.1)	14/207 (6.8)	277/4188 (6.6)
Pestana 2009(44)	PCR	2.60 (0.52-12.98)	4/54 (7.4)	4/134 (3)	0/54 (0)	0/134 (0)	4/54 (7.4)	4/134 (3)
Petrovic 2003(45)	PCR	1 (0.26-3.76; <i>P</i> = .97)	4/96 (4.2)	5/115 (4.3)	1/96 (1)	0/115 (0)	3/96 (3.1)	5/115 (4.3)
Pezzini 2005(46)	PCR	1.17 (0.35-3.92)	6/163 (3.7)	5/158 (3.2)	0/163 (0)	0/158 (0)	6/163 (3.7)	5/158 (3.2)
Pezzini 2007(47)	PCR	1.10 (0.51-5.70)	5/108 (4.6)	6/216 (2.8)	0/108 (0)	0/216 (0)	5/108 (4.6)	6/216 (2.8)
Press 1996(48)	PCR	NR	4/161 (2.5)	19/367 (5.2)	0/161 (0)	0/367 (0)	4/161 (2.5)	19/367 (5.2)
Pullmann 2004(49)	PCR	NR	2/23 (8.7)	3/71 (4.2)	0/23 (0)	0/71 (0)	2/23 (8.7)	3/71 (4.2)
Ranellou 2015(50)	PCR/ hybridization	NR (<i>P</i> = .20)	7/51 (13.7)	4/70 (5.7)	0/51 (0)	0/70 (0)	7/51 (13.7)	4/70 (5.7)
Ridker 1995(51)	PCR	Crude RR 0.7 (0.3-1.4; <i>P</i> =.3) Multivariate adjusted RR 1.0 (0.4-2.2; <i>P</i> =.9)	9/209 (4.3)	42/704 (6)	0/209 (0)	0/704 (0)	9/209 (4.3)	42/704 (6)
Ripoll 1997(53)	PCR	1.7 (0.8-3.4)	17/321 (5.3)	14/428 (3.3)	0/321 (0)	0/428 (0)	17/321 (5.3)	14/428 (3.3)
Rubattu 2005(55)	PCR/RFLP	NR (<i>P</i> = .27)	5/294 (1.7)	2/286 (0.7)	0/294 (0)	0/286 (0)	5/294 (1.7)	2/286 (0.7)
Rubattu 2005(56)	PCR/RFLP	NR	4/115 (3.5)	10/180 (5.6)	1/115 (0.9)	0/180 (0)	3/115 (2.6)	10/180 (5.6)
Sastry 2006(57)	NR	0.50 (0.15-1.66; <i>P</i> =.26)	4/101 (4)	8/101 (7.9)				
Shi 2008(58)	PCR/RFLP	NR	0/97 (0)	0/99 (0)	0/97 (0)	0/99 (0)	0/97 (0)	0/99 (0)
Slooter 2005(59)	PCR	Adjusted OR for age, index year, and residence: 1.8 (0.9-3.6)	14/179 (7.8)	42/763 (5.5)				
Supanc 2014(61)	PCR/RFLP	2.88 (1.01-8.20; <i>P</i> =.40)	14/155 (9)	5/150 (3.3)	2/155 (1.3)	0/150 (0)	12/155 (7.7)	5/150 (3.3)
Szolnoki 2003(62)	PCR	NR	72/867 (8.3)	49/743 (6.6)	3/867 (0.3)	2/743 (0.3)	69/867 (8)	47/743 (6.3)
Tatarsky 2010(63)	PCR/RFLP	NR	6/183 (3.3)	5/188 (2.7)	0/183 (0)	1/188 (0.5)	6/183 (3.3)	4/188 (2.1)
They-They 2012(64)	PCR/RFLP	NR	0/91 (0)	0/182 (0)	0/91 (0)	0/182 (0)	0/91 (0)	0/182 (0)
Tupitsyna 2013(65)	PCR	Russian: 0.9 (0.50-1.76) Ukrainian: 1.9 (0.40-8.97)	42/1450 (2.9)	16/577 (2.8)	0/1450 (0)	0/577 (0)	42/1450 (2.9)	16/577 (2.8)
Voetsch 2000(66)	PCR/RFLP	NR	5/153 (3.3)	8/225 (3.6)	0/153 (0)	0/225 (0)	5/153 (3.3)	8/225 (3.6)
Wypasek 2009(67)	SNP analysis	NR (<i>P</i> = .22)	9/100 (9)	5/107 (4.7)	0/100 (0)	0/107 (0)	9/100 (9)	5/107 (4.7)

NR, not reported; FVL, Factor V Leiden; PCR, polymerase chain reaction; RFLP, restriction fragment length polymorphism; OR, Odds ratio; RR, risk ratio; CI, confidence interval; SNP, Single nucleotide polymorphisms.

Table S4. Results of Included Studies: Prothrombin G20210A Mutation.

References	Test Method	Odds Ratio (95%CI)	Number of All PTM (%)		Number of Homozygous PTM (%)		Number of Heterozygous PTM (%)	
			Cases	Controls	Cases	Controls	Cases	Controls
Aznar 2004(2)	PCR/RFLP	3.75 (1.05-13.34)	4/49 (8.2)	7/294 (2.4)				
Belvis 2006(3)	PCR	NR	3/89 (3.4)	13/201 (6.5)	0/89 (0)	0/201 (0)	3/89(3.4)	13/201 (6.5)
Bentolila 1997(4)	NR	NR	8/125 (6.4)	5/134 (3.7)	0/125(0)	0/134 (0)	8/125 (6.4)	5/134 (3.7)
Biswas 2009(5)	PCR/RFLP	NR	0/120 (0)	0/120 (0)	0/120 (0)	0/120 (0)	0/120 (0)	0/120 (0)
Bolaman 2009(6)	PCR	NR	0/24 (0)	0/53 (0)	0/24 (0)	0/53 (0)	0/24 (0)	0/53 (0)
Celiker 2009(9)	PCR/RFLP	NR	3/162 (1.9)	5/182 (2.7)	0/162 (0)	0/182 (0)	3/162 (1.9)	5/182 (2.7)
Chatterjee 2013(10)	PCR/RFLP	1.00 (0.02-51.35; $P = 1$)	0/52 (0)	0/52 (0)	0/52(0)	0/52 (0)	0/52 (0)	0/52 (0)
Djordjevic 2012(15)	PCR/RFLP	1.33 (0.35-5.13)	4/73 (5.5)	5/120 (4.2)				
Egan 2000(16)	NR	NR	0/42 (0)	13/635 (2)	0/42(0)	0/635 (0)	0/42 (0)	13/635 (2)
Erten 2015(17)	PCR/in situ hybridization	NR	13/212 (6.1)	10/238 (4.2)	0/212 (0)	0/238 (0)	13/212 (6.1)	10/238 (4.2)
Eterovic 2007(18)	PCR/RFLP	NR ($P = .047$)	9/120 (7.5)	3/120 (2.5)	0/120 (0)	0/120 (0)	9/120 (7.5)	3/120 (2.5)
Fan 2010(19)	PCR	1.69 (0.26-10.82)	6/156 (3.8)	122/5817 (2.1)				
Favaretto 2012(20)	PCR	2.97 (1.32-6.69)	29/340 (8.5) PFO: 12/136 (8.8) Non-PFO: 17/204 (8.3)	9/272 (3.3)	0/340 (0) PFO: 0/136 (0) Non-PFO: 0/204 (0)	0/272 (0)	29/340 (8.5) PFO: 12/136 (8.8) Non-PFO: 17/204 (8.3)	9/272 (3.3)
Haeusler 2012(22)	PCR	NR ($P = >.995$)	1/38 (2.6)	10/282 (3.5)	0/38 (0)	0/282 (0)	1/38 (2.6)	10/282 (3.5)
Halbmayer 1998(23)	PCR	NR ($P = .46$)	2/20 (10)	0/20 (0)	0/20 (0)	0/20 (0)	2/20 (10)	0/20 (0)
Hankey 2001(25)	PCR	1.9 (0.5-6.2)	8/219 (3.7)	4/205 (2)				
Jiang 2014(27)	PCR	2.5 (0.9-6.5, $P = .07$)	14/397 (3.5)	6/426 (1.4)	1/397(0.3)	0/426 (0)	13/397 (3.3)	6/426 (1.4)
Kamberi 2016(29)	PCR	2.80 (0.33-23.53 $P = .32$)	1/39 (2.6)	7/102 (6.9)	0/39 (0)	0/102 (0)	1/39 (2.6)	7/102 (6.9)
Karakus 2005(30)	PCR	NR	1/21 (4.8)	1/81 (1.2)	1/21 (4.8)	0/81 (0)	0/21 (0)	1/81 (1.2)
Karttunen 2003(31)	PCR	1.0 (1.0-1.1)	2/57 (3.5)	0/104 (0)	0/57 (0)	0/104 (0)	2/57 (3.5)	0/104 (0)
Kholodkova 2015(32)	PCR	NR	6/114 (5.3)	0/40 (0)	0/114 (0)	0/40 (0)	6/114 (5.3)	0/40 (0)
Krajcoviechova 2015(33)	PCR	Multivariate adjusted OR 2.29 (1.04-5.02, $P = .04$)	22/423 (5.2)	15/614 (2.4)	1/423 (0.2)	0/614 (0)	21/423 (5)	15/614 (2.4)
Linnemann 2008(35)	PCR	0.63 (0.15-2.7, $P = .76$)	2/41 (4.9)	76/930 (8.2)	1/41 (2.4)	2/930 (0.2)	1/41 (2.4)	74/930 (8)
Longstreth 1998(36)	PCR/RFLP	1.6 (0.03-13.4)	1/41 (2.4)	6/382 (1.6)	0/41 (0)	0/382 (0)	1/41 (2.4)	6/382 (1.6)
Lopaciuk 2001(37)	PCR	0.9 (0.2-5.0)	2/100 (2)	5/238 (2.1)	0/100 (0)	0/238 (0)	2/100 (2)	5/238 (2.1)
Martinelli 2006(38)	NR ("DNA analysis")	Crude OR 0.9 (0.3-2.6) Adjusted OR for age 0.9 (0.1-11.2)	5/105 (4.8)	15/293 (5.1)	0/105 (0)	0/293 (0)	5/105 (4.8)	15/293 (5.1)
Moskau 2010(41)	PCR	NR ($P = .88$)	7/167 (4.2)	10/500 (2)	0/167 (0)	0/500 (0)	7/167 (4.2)	10/500 (2)
Pahus 2016(43)	PCR	Heterozygous 1.59 (0.32-4.81)	3/92 (3.3)	1377/64631 (2.1)	0/92 (0)	8/64631 (0)	3/92 (3.3)	1369/64631 (2.1)
Pezzini 2005(46)	PCR	2.68 (0.70-10.3)	9/163 (5.5)	3/158 (1.9)	1/163 (0.6)	0/158 (0)	8/163 (4.9)	3/158 (1.9)
Pezzini 2007(47)	PCR	6.52 (1.73-24.6)	10/108 (9.3)	3/216 (1.4)	1/108 (0.9)	0/216 (0)	9/108 (8.3)	3/216 (1.4)
Pullmann 2004(49)	PCR	NR	0/23 (0)	3/71 (4.2)	0/23 (0)	0/71 (0)	0/23 (0)	3/71 (4.2)
Ranellou 2015(50)	PCR	NR ($P = .70$)	7/51 (13.7)	4/70 (5.7)	0/51 (0)	0/70 (0)	7/51 (13.7)	4/70 (5.7)
Ridker 1999(52)	PCR	Crude RR 1.1 (0.6-2.1; $P = .8$)	11/259 (4.2)	69/1774 (3.9)	0/259(0)	1/1774 (0.1)	11/259 (4.2)	68/1774 (3.8)

		Multivariate adjusted RR 1.1 (0.5-2.4; $P=.7$)						
Rubattu 2005(55)	PCR/RFLP	NR ($P = .95$)	12/294 (4.1)	12/286 (4.2)	0/294 (0)	0/286 (0)	12/294 (4.1)	12/286 (4.2)
Rubattu 2005(56)	PCR	NR	8/115 (7)	10/180 (5.6)	0/115 (0)	0/180 (0)	8/115 (7)	10/180 (5.6)
Sastry 2006(57)	NR	NR	2/101 (2)	0/101 (0)				
Slooter 2005(59)	PCR	Adjusted OR for age, index year, and residence: 1.0 (0.3-3.0)	5/188 (2.7)	18/763 (2.4)				
Smiles 2002(60)	PCR	Crude OR 1.25 (0.46-3.39) Adjusted OR for sex, race, and clinic site: 1.40 (0.51-3.86)	6/182 (3.3)	12/453 (2.6)	0/182(0)	0/453 (0)	6/182 (3.3)	12/453 (2.6)
Supanc 2014(61)	PCR	3.50 (0.72-17.24; $P = .17$)	7/155 (4.5)	2/150 (1.3)	0/155 (0)	0/150 (0)	7/155 (4.5)	2/150 (1.3)
Szolnoki 2003(62)	PCR	NR	5/867 (0.6)	4/743 (0.5)	0/867 (0)	0/743 (0)	5/867 (0.6)	4/743 (0.5)
Tatarsky 2010(63)	PCR/RFLP	NR	8/183 (4.4)	3/188 (1.6)	0/183 (0)	0/188 (0)	8/183 (4.4)	3/188 (1.6)
They-They 2012(64)	PCR/RFLP	2.3 (0.97-5.8; $P = .60$)	11/91 (12.1)	10/182 (5.5)	0/91 (0)	0/182 (0)	11/91 (12.1)	10/182 (5.5)
Tupitsyna 2013(65)	PCR	Russian: 0.7 (0.35-1.30)	29/1450 (2)	20/817 (2.4)	0/1450 (0)	0/817 (0)	29/1450 (2)	20/817 (2.4)
Voetsch 2000(66)	PCR/RFLP	NR	7/153 (4.6)	5/229 (2.2)	0/153 (0)	0/229 (0)	7/153 (4.6)	5/229 (2.2)
Wypasek 2009(67)	SNP analysis	NR ($P = .15$)	1/100 (1)	1/107 (0.9)	0/100 (0)	0/107 (0)	1/100 (1)	1/107 (0.9)

NR, not reported; PTM, Prothrombin G20210A Mutation; PCR, polymerase chain reaction; RFLP, restriction fragment length polymorphism; OR, Odds ratio; RR, risk ratio; CI, confidence interval; SNP, Single nucleotide polymorphisms.

Table S5. Results of Included Studies: Protein C Deficiency.

References	Test Method	Definition of deficiency	Timing of test after stroke	Exclusion of anticoagulants	Odds Ratio (95%CI)	Number of PCD (%)	
						Cases	Controls
Biswas 2009(5)	ELISA	<65%	3-6 months after stroke	Yes	NR	4/120 (3.3)	0/120 (0)
Chatterjee 2013(10)	Clot-based assay	<70%	≥4 months after stroke	NR	0.15 (0.01-1.30; <i>P</i> = .12)	6/52 (11.5)	1/52 (1.9)
Chen 2003(11)	Chromogenic assay	<70%	NR (If a first test was abnormal, a second test was done 6 weeks later; a deficiency was defined by <70% in two tests)	Yes	5.29 (NR)	14/104 (13.5)	1/35 (2.9)
D'Amico 1998(13)	NR	NR	2-15 days after stroke	NR	NR	0/31 (0)	0/124 (0)
De Lucia 1999(14)	NR	<76.9% (<2.5th percentile)	At 3 months after stroke	Yes	NR	2/50 (4)	1/100 (1)
Favaretto 2012(20)	Chromogenic assay	<68%	1 month after discharge	NR	NR	0/340 (0)	0/272 (0)
Hankey 2001(25)	Chromogenic assay	<70%	Within 7 days and at 3-6 months (any low level is considered deficiency)	NR	0.7 (0.2-3.1; <i>P</i> = .6)	3/219 (1.4)	4/205 (2)
Jerrard-Dunne 2003(26)	Chromogenic assay	<-2SD of ethnic-specific controls	First test at presentation If abnormal, repeated test at ≥3 months after stroke Results of the repeated test were used	Yes	3.05 (0.60-15.39; <i>P</i> = .16)	6/130 (4.6)	2/130 (1.5)
Karakus 2005(30)	Chromogenic assay	NR	NR	NR	NR	1/21 (4.8)	1/81 (1.2)
Karttunen 2003(31)	Chromogenic assay	NR	>2 months after stroke	NR	NR	0/57 (0)	0/104 (0)
Linnemann 2008(35)	Chromogenic assay	<74%	NR (a deficiency was defined by repeatedly low activity)	Yes	0.98 (0.97-0.99; <i>P</i> = 1.00)	0/31 (0)	20/817 (2.4)
Pahus 2016(43)	Chromogenic and clot-based assay	<0.65 U/L (both assays)	Average 2 months after stroke (Repeated in cases of deficiency)	NR	1.90 (0.04-12.55)	1/363 (0.3)	14/9648 (0.1)
Romdhane 2011(54)	Chromogenic assay	NR	NR	NR	2 (NR; <i>P</i> = .6)	2/20 (10)	3/54 (5.6)
Sastry 2006(57)	Functional assay	NR	NR	Yes	NR	1/86 (1.2)	0/101 (0)
Zimba 2017(68)	Chromogenic assay	<70%	More than 48 hour, up to 1 month	Yes	NR (<i>P</i> = .06)	5/52 (9.6)	0/52 (0)

NR, not reported; PCD, protein C Deficiency; ELISA, enzyme-linked immunosorbent assay; CI, confidence interval.

Table S6. Results of Included Studies: Protein S Deficiency.

References	Test Method	Definition of deficiency	Timing of test after stroke	Exclusion of anticoagulants	Odds Ratio (95%CI)	Number of PSD (%)	
						Cases	Controls
Biswas 2009(5)	ELISA	<50%	3-6 months after stroke	Yes	NR	6/120 (5)	0/120 (0)
Chatterjee 2013(10)	Clot-based assay	<65%	≥4 months after stroke	NR	0.05 (0.01-0.38; <i>P</i> <.001)	15/52 (28.8)	1/52 (1.9)
Chen 2003(11)	Chromogenic assay	<60%	NR (If a first test was abnormal, a second test was done 6 weeks later; a deficiency was defined by <60% in two tests)	Yes	2.86 (NR)	22/104 (21.2)	3/35 (8.6)
D'Amico 1998(13)	NR	NR	2-15 days after stroke (Repeated at >6 months after stroke; a deficiency was defined by abnormalities in both tests)	NR	NR	2/31 (6.5)	0/124 (0)
De Lucia 1999(14)	NR	<74.6% (<2.5th percentile)	At 3 months after stroke	Yes	NR	3/50 (6)	1/100 (1)
Favaretto 2012(20)	Chromogenic assay	<62%	1 month after discharge	NR	NR	0/340 (0)	0/272 (0)
Hankey 2001(25)	Immunoelectrophoresis	<55%	Within 7 days and at 3-6 months (any low level is considered deficiency)	NR	0.9 (0.1-6.7; <i>P</i> = .5)	2/219 (0.9)	2/205 (1)
Jerrard-Dunne 2003(26)	Immunoassay	<-2SD of ethnic-specific controls	First test at presentation If abnormal, repeated test at ≥3 months after stroke Results of the repeated test were used	Yes	2.00 (0.36-11.1; <i>P</i> = .42)	4/130 (3.1)	2/130 (1.5)
Karakus 2005(30)	Immunoassay	NR	NR	NR	NR	1/21 (4.8)	0/81 (0)
Karttunen 2003(31)	Clot-based assay	NR	>2 months after stroke	NR	1.0 (1.0-1.1)	1/57 (1.8)	0/104 (0)
Linnemann 2008(35)	Clot-based assay	Male <70%, Female <60%	NR (a deficiency was defined by repeatedly low activity)	Yes	0.97 (0.13-7.39; <i>P</i> = 1.00)	1/30 (3.3)	28/788 (3.6)
Mayer 1993(39)	Free protein S: Immunoelectrophoresis	Free PS <20% of normal total PS	Cases: Average 2.4 days after onset of stroke Controls: Average 5.2 days after admission	Yes	1.1 (0.5-2.2)	20/94 (21.3)	19/94 (20.2)
Mochan 2005(40)	Clot-based assay	NR	Cases: Repeated at 3 months after stroke Controls: NR	NR	NR	11/33 (33.3)	12/33 (36.4)
Pahus 2016(43)	Clot-based assay, ELISA	Clot-based assay: <0.65 U/L AND Free PS ELISA: <0.13 U/L (until 2005) and <0.55 U/L (from 2006)	Average 2 months after stroke (Repeated in cases of deficiency)	NR	2.32 (0.24-11.27)	2/364 (0.5)	9/3788 (0.2)
Romdhane 2011(54)	Clot-based assay	NR	NR	NR	11.3 (NR; <i>P</i> = .003)	6/20 (30)	2/54 (3.7)
Sastry 2006(57)	Functional assay	NR	NR	Yes	NR	1/86 (1.2)	1/101 (1)
Zimba 2017(68)	Immunoassay	<60%	More than 48 hour, up to 1 month	Yes	NR (<i>P</i> = .42)	22/52 (42.3)	18/52 (34.6)

NR, not reported; PSD, protein S Deficiency; ELISA, enzyme-linked immunosorbent assay; CI, confidence interval.

Table S7. Results of Included Studies: Antithrombin Deficiency.

References	Test Method	Definition of deficiency	Timing of test after stroke	Exclusion of anticoagulants	Odds Ratio (95%CI)	Number of ATD (%)	
						Cases	Controls
Chatterjee 2013(10)	Chromogenic assay	<75%	≥4 months	NR	1.00 (0.02-51.35; <i>P</i> = 1)	0/52 (0)	0/52 (0)
Chen 2003(11)	Chromogenic assay	<70%	NR (If a first test was abnormal, a second test was done 6 weeks later; a deficiency was defined by <70% in two tests)	Yes	0.33 (NR)	1/104 (1)	1/35 (2.9)
De Lucia 1999(14)	NR	NR	3 months	Yes	NR	0/50 (0)	0/100 (0)
Favaretto 2012(20)	Chromogenic assay	<80%	1 month after discharge	NR	NR	0/340 (0)	0/272 (0)
Hankey 2001(25)	Chromogenic assay	82	Within 7 days and at 3-6 months (any low level is considered deficiency)	NR	1.3 (0.5-3.3; <i>P</i> = .6)	11/21 (5.2)	8/205 (3.7)
Jerrard-Dunne 2003(26)	Chromogenic assay	<-2SD of ethnic-specific controls	First test at presentation If abnormal, repeated test at ≥3 months after stroke Results of the repeated test were used	Yes	NR	0/130 (0)	0/130 (0)
Karakus 2005(30)	Chromogenic assay	NR	NR	NR	NR	1/21 (4.8)	0/81 (0)
Karittunen 2003(31)	Chromogenic assay	NR	>2 months after stroke	NR	0.9 (0.1-10.5)	1/57 (1.8)	2/104 (1.9)
Linnemann 2008(35)	Chromogenic assay	<86%	NR (a deficiency was defined by repeatedly low activity)	Yes	0.88 (0.12-6.60, <i>P</i> = 1.00)	1/40 (2.5)	29/993 (2.9)
Pahus 2016(43)	Functional assay	<0.60 x 10 ³ U/L	Average 2 months (Repeated in cases of deficiency)	NR	NR	0/288 (0)	16/9669 (0.2)
Romdhane 2011(54)	Chromogenic assay	NR	NR	NR	5.6 (NR; <i>P</i> = .01)	7/20 (35)	5/54 (9.3)
Sastry 2006(57)	Functional assay	NR	NR	Yes	NR	3/86 (3.5)	0/101 (0)

NR, not reported; ATD, Antithrombin Deficiency; CI, confidence interval.

Table S8. Components of Quality Assessment.

References	Research question	Study population	Target population and case representation	Sample size justification	Groups recruited from the same population	Inclusion and exclusion criteria prespecified and applied uniformly	Case and control definitions	Random selection of study participants	Concurrent controls	Exposure assessed prior to outcome measurement	Exposure measures and assessment	Blinding of exposure assessors	Statistical analysis	Study Quality
Anadure 2017(1)	Y	Y	NR	Y	CD	CD	Y	N	N	N	Y	NR	Y	Good
Aznar 2004(2)	Y	N	NR	N	Y	Y	Y	N	N	N	N	NR	N	Fair
Belvis 2006(3)	Y	Y	NR	N	N	N	Y	N	N	N	Y	Y	N	Fair
Bentolila 1997(4)	Y	Y	NR	N	CD	Y	Y	N	N	N	N	NR	N	Fair
Biswas 2009(5)	Y	N	NR	N	Y	Y	Y	N	N	N	N	NR	N	Fair
Bolaman 2009(6)	Y	N	NR	N	CD	Y	Y	N	N	N	Y	NR	N	Fair
Buryu 2005(7)	Y	N	NR	N	CD	CD	Y	N	N	N	Y	NR	N	Fair
Catto 1995(8)	Y	N	NR	N	Y	CD	Y	N	N	N	Y	NR	N	Fair
Celiker 2009(9)	Y	Y	NR	N	Y	N	Y	NA	N	N	Y	NR	N	Fair
Chatterjee 2013(10)	Y	N	NR	N	CD	Y	Y	N	N	N	N	NR	N	Fair
Chen 2003(11)	Y	N	NR	N	Y	CD	Y	N	N	N	Y	N	Y	Fair
Cushman 1998(12)	Y	Y	NR	N	Y	Y	Y	Y	N	Y	Y	NR	Y	Good
D'Amico 1998(13)	Y	Y	NR	N	Y	Y	Y	N	N	N	N	NR	N	Fair
De Lucia 1999(14)	Y	Y	NR	N	CD	Y	Y	N	N	N	N	Y	N	Fair
Djordjevic 2012(15)	Y	N	NR	N	Y	Y	Y	N	N	N	Y	NR	N	Fair
Egan 2000(16)	Y	Y	NR	N	Y	CD	Y	N	N	N	N	NR	N	Fair
Erten 2015(17)	Y	Y	NR	N	CD	Y	N	N	N	N	Y	NR	Y	Fair
Eterovic 2007(18)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	Y	Y	Good
Fan 2010(19)	Y	N	NR	N	Y	Y	N	NA	N	CD	Y	NR	Y	Fair
Favaretto 2012(20)	Y	Y	NR	N	Y	N	Y	N	N	N	Y	NR	Y	Fair
Go 2003(21)	Y	Y	NR	N	Y	Y	Y	NA	N	N	Y	NR	Y	Good
Haeusler 2012(22)	Y	N	NR	N	N	N	Y	N	N	N	Y	N	N	Fair
Halbmayer 1998(23)	Y	N	NR	N	CD	CD	N	N	N	N	Y	NR	N	Poor
Hamedani 2013(24)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	NR	Y	Fair
Hankey 2001(25)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	NR	Y	Good
Jerrard-Dunne 2003(26)	Y	N	NR	N	Y	CD	Y	Y	N	N	Y	NR	Y	Good
Jiang 2014(27)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	NR	Y	Fair
Juul 2002(28)	Y	Y	NR	N	Y	Y	Y	NA	N	Y	Y	NR	Y	Fair
Kamberi 2016(29)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	NR	N	Fair
Karakus 2005(30)	Y	N	NR	N	Y	Y	N	Y	N	N	N	NR	N	Fair
Karttunen 2003(31)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	Y	Y	Good
Kholodkova 2015(32)	Y	N	NR	N	Y	Y	Y	N	N	N	Y	NR	N	Fair
Krajcoviechova 2015(33)	Y	Y	NR	Y	Y	N	Y	N	N	N	Y	NR	Y	Fair
Kumar 2017(34)	Y	N	NR	Y	Y	Y	Y	NA	N	N	Y	Y	Y	Good
Linneman 2008(35)	Y	Y	NR	N	Y	Y	Y	NA	N	N	Y	NR	Y	Good

References	Research question	Study population	Target population and case representation	Sample size justification	Groups recruited from the same population	Inclusion and exclusion criteria prespecified and applied uniformly	Case and control definitions	Random selection of study participants	Concurrent controls	Exposure assessed prior to outcome measurement	Exposure measures and assessment	Blinding of exposure assessors	Statistical analysis	Study Quality
Longstreth 1998(36)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	Y	N	Good
Lopaciuk 2001(37)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	NR	N	Good
Martinelli 2006(38)	Y	Y	NR	N	Y	Y	Y	N	N	N	N	NR	Y	Fair
Mayer 1993(39)	Y	Y	NR	N	Y	Y	CD	N	N	N	Y	NR	N	Fair
Mochan 2005(40)	Y	Y	NR	N	N	N	Y	NA	N	N	N	NR	N	Fair
Moskau 2010(41)	Y	N	NR	N	N	CD	Y	N	N	N	Y	NR	N	Fair
Nagayama 1996(42)	Y	N	NR	N	Y	CD	Y	N	N	N	Y	NR	N	Fair
Pahus 2016(43)	Y	Y	NR	N	N	N	Y	NA	N	N	N	Y	N	Fair
Pestana 2009(44)	Y	N	NR	N	Y	Y	Y	Y	N	N	Y	NR	Y	Fair
Petrovic 2003(45)	Y	N	NR	N	Y	Y	Y	N	N	N	Y	Y	Y	Fair
Pezzini 2005(46)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	NR	Y	Good
Pezzini 2007(47)	Y	Y	NR	Y	Y	Y	Y	N	N	N	Y	NR	Y	Good
Press 1996(48)	Y	N	NR	N	Y	N	N	N	N	N	Y	NR	N	Fair
Pullmann 2004(49)	Y	N	NR	N	Y	Y	Y	NA	N	N	Y	NR	Y	Fair
Ranellou 2015(50)	Y	N	NR	N	Y	Y	Y	N	N	N	Y	NR	N	Fair
Ridker 1995(51)	Y	Y	NR	N	Y	Y	Y	Y	N	Y	Y	NR	Y	Good
Ridker 1999(52)	Y	Y	NR	N	Y	Y	Y	Y	N	Y	Y	NR	Y	Good
Ripoll 1997(53)	Y	N	NR	N	CD	CD	N	N	N	N	Y	NR	N	Fair
Romdhane 2011(54)	Y	N	NR	N	CD	CD	CD	N	N	N	N	NR	N	Poor
Rubattu 2005(55)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	Y	Y	Fair
Rubattu 2005(56)	Y	N	NR	N	Y	N	Y	N	N	N	Y	NR	Y	Fair
Sastry 2006(57)	Y	Y	NR	Y	Y	Y	Y	N	N	N	N	NR	Y	Fair
Shi 2008(58)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	NR	Y	Good
Slooter 2005(59)	Y	Y	NR	N	Y	Y	Y	Y	N	N	Y	Y	Y	Good
Smiles 2002(60)	Y	Y	NR	N	Y	Y	Y	NA	N	Y	Y	Y	Y	Good
Supanc 2014(61)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	NR	Y	Good
Szolnoki 2003(62)	Y	N	NR	N	Y	Y	Y	Y	N	N	Y	Y	Y	Good
Tatarsky 2010(63)	Y	N	NR	N	Y	CD	N	N	N	N	Y	NR	N	Fair
They-They 2012(64)	Y	Y	NR	N	Y	Y	Y	N	N	N	Y	N	Y	Good
Tupitsyna 2013(65)	Y	N	NR	N	Y	CD	N	N	N	N	Y	NR	N	Poor
Voetsch 2000(66)	Y	Y	NR	N	Y	N	Y	Y	N	N	Y	NR	N	Fair
Wypasek 2009(67)	Y	N	NR	N	NR	Y	Y	N	N	N	Y	NR	N	Fair
Zimba 2017(68)	Y	Y	NR	Y	Y	Y	Y	N	N	N	N	NR	Y	Good

Y, Yes; N, No; NR, not reported; CD, cannot determined; NA, not applicable

Table S9. Additional sensitivity analyses.

Types of studies that were excluded from the analysis	Excluded studies	Thrombophilias	Pooled OR (95%CI)	I²,%
Studies with enriched case population (those who were referred for thrombophilia testing because of a clinical indication or recruited from a thrombophilia center)	Aznar 2004, Martinelli 2006, Pahus 2016	FVL	1.24 (1.07, 1.44)	0
		PTM	1.47 (1.21, 1.77)	0
		PCD	2.17 (1.15, 4.11)	0
		PSD	2.30 (1.36, 4.07)	13.2
		ATD	1.37 (0.59, 2.94)	6.8
Studies that used self-reported history of stroke rather than imaging to define cases	Fan 2010	FVL	1.26 (1.09, 1.47)	0
		PTM	1.47 (1.20, 1.80)	0
Studies that were rated as poor quality	Halbmayer 1998, Romdhane 2011, Tupitsyna 2013	FVL	1.26 (1.09, 1.48)	0
		PTM	1.53 (1.26, 1.86)	0
		PCD	2.16 (1.14, 4.09)	0
		PSD	2.01 (1.22, 3.48)	0
		ATD	0.91 (0.38, 1.97)	0
Studies that included cases of recurrent ischemic stroke	Chatterjee 2013, Kumar 2017, They-They 2012	FVL	1.24 (1.08, 1.45)	0
		PTM	1.46 (1.20, 1.78)	0
		PCD	1.93 (1.01, 3.60)	0
		PSD	1.88 (1.19, 3.20)	0
		ATD	1.33 (0.59, 2.76)	8.0

FVL, Factor V Leiden; PTM, Prothrombin G20210A Mutation; PCD, Protein C Deficiency; PSD, Protein S Deficiency; ATD, Antithrombin Deficiency; CI, confidence interval

Figure S1. Forest plot showing pooled odds ratio for Factor V Leiden.




Figure S2. Forest plot showing pooled odds ratio for Factor V Leiden (Homozygous).




Figure S3. Forest plot showing pooled odds ratio for Factor V Leiden (Heterozygous).




Figure S4. Funnel plot of included studies.




Figure S5. Forest plot showing pooled odds ratio for Prothrombin G20210A Mutation.




Figure S6. Forest plot showing pooled odds ratio for Prothrombin G20210A Mutation (Homozygous).




Figure S7. Forest plot showing pooled odds ratio for Factor V Leiden (Heterozygous).




Figure S8. Forest plot showing pooled odds ratio for protein C deficiency.




Figure S9. Forest plot showing pooled odds ratio for protein S deficiency.




Figure S10. Forest plot showing pooled odds ratio for antithrombin deficiency.




Figure S11. Subgroup Analyses: Factor V Leiden.




Figure S12. Subgroup Analyses: Prothrombin G20210A Mutation.




Figure S13. Subgroup Analyses: Protein C Deficiency.




Figure S14. Subgroup Analyses: Protein S Deficiency.





Figure S15. Subgroup Analyses: Antithrombin Deficiency.



Supplemental References:

1. Anadure RC, Nagaraja D, Narayanan C. A genetic study of Factor V Leiden (G1691A) mutation in young ischemic strokes with large vessel disease in a South Indian population. *Journal of Clinical Neuroscience*. 2017;44:346-52.
2. Aznar JM, Vaya A, Corella D, Ferrando F, Villa P, Estelles A. Factor V Leiden and prothrombin G20210A mutations in young adults with cryptogenic ischemic stroke. *Thromb Haemost*. 2004;91:1031-4.
3. Belvis RS, A.; Marti-Fabregas, J.; Cocho, D.; Borrell, M.; Fontcuberta, J.; Marti-Vilalta, J. L. Diagnostic yield of prothrombotic state studies in cryptogenic stroke. *Acta Neurol Scand*. 2006;114:250-3.
4. Bentolila SR, L.; Drouet, L.; Mazoyer, E.; Woimant, F. Thrombophilia due to 20210 G-->A prothrombin polymorphism and cerebral ischemia in the young. *Stroke*. 1997;28:1846-7.
5. Biswas AR, R.; Meena, A.; Akhter, S.; Sharma, V.; Yadav, B. K.; Behari, M.; Saxena, R. Prothrombotic factors and the risk of acute onset non-cardioembolic stroke in young Asian Indians. *Thromb Res*. 2009;124:397-402.
6. Bolaman ZO, A.; Kiylioglu, N.; Kadikoylu, G.; Erturk, A.; Batun, S.; Akyol, A. Hereditary thrombophilic factors in stroke due to cerebral infarct. *Am J Med Sci*. 2009;337:11-3.
7. Buyru NA, J.; Somay, G.; Ulutin, T. Factor V Leiden mutation in cerebrovascular disease. *Clin Appl Thromb Hemost*. 2005;11:339-42.
8. Catto AC, A.; Ireland, H.; Bayston, T. A.; Philippou, H.; Barrett, J.; Lane, D. A.; Grant, P. J. Factor V Leiden gene mutation and thrombin generation in relation to the development of acute stroke. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 1995;15:783-5.
9. Celiker GC, U.; Verdi, H.; Yazici, A. C.; Ozbek, N.; Atac, F. B. Prevalence of thrombophilic mutations and ACE I/D polymorphism in Turkish ischemic stroke patients. *Clin Appl Thromb Hemost*. 2009;15:415-20.
10. Chatterjee TG, N.; Choudhry, V. P.; Behari, M.; Saxena, R.; Ashraf, M. Z. Prediction of ischemic stroke in young Indians: is thrombophilia profiling a way out? *Blood Coagul Fibrinolysis*. 2013;24:449-53.
11. Chen WHL, M. Y.; Chang, Y. Y.; Chen, S. S.; Liu, J. S. The prevalence of protein C, protein S, and antithrombin III deficiency in non-APS/SLE Chinese adults with noncardiac cerebral ischemia. *Clin Appl Thromb Hemost*. 2003;9:155-62.
12. Cushman MR, F. R.; Psaty, B. M.; Cook, E. F.; Valliere, J.; Kuller, L. H.; Tracy, R. P. Factor V Leiden is not a risk factor for arterial vascular disease in the elderly: results from the Cardiovascular Health Study. *Thromb Haemost*. 1998;79:912-5.
13. D'Amico DM, F.; Leone, M.; Ariano, C.; Ciusani, E.; Erba, N.; Grazi, L.; Ferraris, A.; Schieroni, F.; Bussone, G. Genetic abnormalities of the protein C system: shared risk factors in young adults with migraine with aura and with ischemic stroke? *Cephalalgia*. 1998;18:618-21; discussion 591.
14. De Lucia DP, M. L.; Ammendola, F.; Pezzella, S.; Del Giudice, V.; Marotta, R.; Renis, V.; Di Mauro, C.; Maisto, G.; Masi, S.; Nina, P.; Franco, A.; Schisano, G. Association of elevated levels of prothrombin fragment 1+2 and Arg506 to Gln mutation in patients with a history of ischemic stroke. *J Neurosurg Sci*. 1999;43:45-50; discussion -1.
15. Djordjevic VS, M.; Brankovic-Sreckovic, V.; Rakicevic, L.; Damnjanovic, T.; Antonijevic, N.; Radojkovic, D. Prothrombotic genetic risk factors in stroke: a

- possible different role in pediatric and adult patients. *Clin Appl Thromb Hemost.* 2012;18:658-61.
16. Egan RAK, J. M.; Press, R.; Lutsep, H. L. Lack of prothrombin gene mutation in young stroke patients. *Journal of Stroke and Cerebrovascular Diseases.* 2000;9:229-31.
 17. Erten ND, S.; Sütçü, R. Prevalence of thrombophilic mutations in ischemic stroke patients in isparta, Turkey. *Turk Noroloji Dergisi.* 2015;21:42-8.
 18. Eterovic DT, M.; Culic, V.; Zadro, R.; Primorac, D. Lower contribution of factor V Leiden or G202104 mutations to ischemic stroke in patients with clinical risk factors: pair-matched case-control study. *Clin Appl Thromb Hemost.* 2007;13:188-93.
 19. Fan AZF, J.; Yesupriya, A.; Chang, M.; Kilmer, G.; House, M.; Hayes, D.; Ned, R. M.; Dowling, N. F.; Mokdad, A. H. Gene polymorphisms in association with self-reported stroke in US adults. *Application of Clinical Genetics.* 2010;3:23-8.
 20. Favaretto ES, M.; Conti, E.; Legnani, C.; Palareti, G. G1691A factor V and G20210A FII mutations, acute ischemic stroke of unknown cause, and patent foramen ovale. *Thromb Res.* 2012;130:720-4.
 21. Go ASR, G. L.; Hylek, E. M.; Phillips, K. A.; Liu, L.; Henault, L. E.; Selby, J. V.; Singer, D. E. Factor V Leiden and risk of ischemic stroke in nonvalvular atrial fibrillation: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. *J Thromb Thrombolysis.* 2003;15:41-6.
 22. Haeusler KGH, J.; Hoppe, B.; Kasabov, R.; Malzahn, U.; Endres, M.; Koscielny, J.; Jungehulsing, G. J. Thrombophilia screening in young patients with cryptogenic stroke. Prevalence of gene polymorphisms compared to healthy blood donors and impact on secondary stroke prevention. *Hamostaseologie.* 2012;32:147-52.
 23. Halbmayr WMH, A.; Hermann, K. M.; Fischer, M. The 20210A allele of the prothrombin gene: a risk factor for juvenile stroke? Result of a pilot study. *Blood Coagul Fibrinolysis.* 1998;9:209-10.
 24. Hamedani AGC, J. W.; Cheng, Y.; Sparks, M. J.; O'Connell, J. R.; Stine, O. C.; Wozniak, M. A.; Stern, B. J.; Mitchell, B. D.; Kittner, S. J. Factor V Leiden and ischemic stroke risk: the Genetics of Early Onset Stroke (GEOS) study. *J Stroke Cerebrovasc Dis.* 2013;22:419-23.
 25. Hankey GJE, J. W.; van Bockxmeer, F. M.; Lofthouse, E.; Staples, N.; Baker, R. I. Inherited thrombophilia in ischemic stroke and its pathogenic subtypes. *Stroke.* 2001;32:1793-9.
 26. Jerrard-Dunne PE, A.; McGovern, R.; Hajat, C.; Kalra, L.; Rudd, A. G.; Wolfe, C. D.; Markus, H. S. Ethnic differences in markers of thrombophilia: implications for the investigation of ischemic stroke in multiethnic populations: the South London Ethnicity and Stroke Study. *Stroke.* 2003;34:1821-6.
 27. Jiang BR, K. A.; Hamedani, A.; Cheng, Y.; Sparks, M. J.; Koontz, D.; Bean, C. J.; Gallagher, M.; Hooper, W. C.; McArdle, P. F.; O'Connell, J. R.; Stine, O. C.; Wozniak, M. A.; Stern, B. J.; Mitchell, B. D.; Kittner, S. J.; Cole, J. W. Prothrombin G20210A mutation is associated with young-onset stroke: the genetics of early-onset stroke study and meta-analysis. *Stroke.* 2014;45:961-7.
 28. Juul KT-H, A.; Steffensen, R.; Kofoed, S.; Jensen, G.; Nordestgaard, B. G. Factor V Leiden: The Copenhagen City Heart Study and 2 meta-analyses. *Blood.* 2002;100:3-10.
 29. Kamberi BK, F.; Spiroski, M. Vascular genetic variants and ischemic stroke susceptibility in Albanians from the Republic of Macedonia. *Macedonian Journal of Medical Sciences.* 2016;4:556-64.

30. Karakuş ZG, E.; Başlamışlı, F.; Tanriverdi, K. Prothrombotic heritable risk factors for cerebral ischemic infarction, acute myocardial infarction and venous thrombosis in young adult Turkish patients. *Annals of Medical Sciences*. 2005;14:31-6.
31. Karttunen VH, L.; Rasi, V.; Vahtera, E.; Hillbom, M. Factor V Leiden and prothrombin gene mutation may predispose to paradoxical embolism in subjects with patent foramen ovale. *Blood Coagul Fibrinolysis*. 2003;14:261-8.
32. Kholodkova OLBV, M. K.; Torgalo, E. O.; Vovk, T. B.; Kravchenko, N. K.; Raksha, N. G.; Melnyk, V. S.; Savchuk, O. M.; Ostapchenko, L. I. FII G20210A and FV Leiden G1691A polymorphisms in patients with atherothrombotic and cardioembolic ischemic stroke from Ukraine. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2015;6:1568-71.
33. Krajcoviechova AW, P.; Mayer, O., Jr.; Vanek, J.; Hajkova, J.; Hlinovsky, D.; Kvasnicka, T.; Tremblay, J.; Hamet, P.; Filipovsky, J.; Kvasnicka, J.; Cifkova, R. Tobacco smoking strongly modifies the association of prothrombin G20210A with undetermined stroke: consecutive survivors and population-based controls. *Atherosclerosis*. 2015;240:446-52.
34. Kumar AM, S.; Sagar, R.; Kumar, P.; Yadav, A.; Talwar, P.; Raj, R.; Prasad, K. Relationship between factor v leiden gene variant and risk of ischemic stroke: A case-control study. *Annals of Indian Academy of Neurology*. 2017;20:284-8.
35. Linnemann BS, M.; Zgouras, D.; Erbe, M.; Jarosch-Preusche, M.; Lindhoff-Last, E. Are patients with thrombophilia and previous venous thromboembolism at higher risk to arterial thrombosis? *Thromb Res*. 2008;121:743-50.
36. Longstreth WT, Jr.; Rosendaal, F. R.; Siscovick, D. S.; Vos, H. L.; Schwartz, S. M.; Psaty, B. M.; Raghunathan, T. E.; Koepsell, T. D.; Reitsma, P. H. Risk of stroke in young women and two prothrombotic mutations: factor V Leiden and prothrombin gene variant (G20210A). *Stroke*. 1998;29:577-80.
37. Lopaciuk SB, K.; Kwiecinski, H.; Mickielewicz, A.; Czlonkowska, A.; Mendel, T.; Kuczynska-Zardzewialy, A.; Szlagowska, D.; Windyga, J.; Schroder, W.; Herrmann, F. H.; Jedrzejowska, H. Factor V Leiden, prothrombin gene G20210A variant, and methylenetetrahydrofolate reductase C677T genotype in young adults with ischemic stroke. *Clin Appl Thromb Hemost*. 2001;7:346-50.
38. Martinelli IB, T.; Burgo, I.; Di Domenico, S.; Mannucci, P. M. Oral contraceptive use, thrombophilia and their interaction in young women with ischemic stroke. *Haematologica*. 2006;91:844-7.
39. Mayer SAS, R. L.; Hurlet-Jensen, A.; Shi, T.; Mohr, J. P. Free protein S deficiency in acute ischemic stroke. A case-control study. *Stroke*. 1993;24(2):224-7.
40. Mochan AM, M.; Modi, G. Protein S deficiency in HIV associated ischaemic stroke: an epiphenomenon of HIV infection. *J Neurol Neurosurg Psychiatry*. 2005;76:1455-6.
41. Moskau SS, K.; Semmler, A.; Schweichel, D.; Harbrecht, U.; Muller, J.; Pohl, C.; Klockgether, T.; Linnebank, M. Common genetic coagulation variants are not associated with ischemic stroke in a case-control study. *Neurol Res*. 2010;32:519-22.
42. Nagayama TN, M.; Tsuda, M.; Yoshii, F.; Shinohara, Y. Low prevalence of activated protein C resistance and factor V Leiden in ischemic stroke in Japan. *Cerebrovascular Diseases*. 1996;6:356-9.
43. Pahus SHH, A. T.; Hvas, A. M. Thrombophilia testing in young patients with ischemic stroke. *Thromb Res*. 2016;137:108-12.
44. Pestana CIT, A.; Blanco, S.; Rojas, M. J.; Mendez, C.; Lopez, J. L.; de Bosch, N. B.; Porco, A. Factor V Leiden and the risk of venous thrombosis, myocardial infarction,

- and stroke: a case-control study in Venezuela. *Genet Test Mol Biomarkers*. 2009;13:537-42.
45. Petrovic DM, T.; Kobal, J.; Bregar, D.; Potisk, K. P.; Peterlin, B. Prothrombotic gene polymorphisms and atherothrombotic cerebral infarction. *Acta Neurol Scand*. 2003;108:109-13.
 46. Pezzini AG, M.; Del Zotto, E.; Archetti, S.; Spezi, R.; Vergani, V.; Assanelli, D.; Caimi, L.; Padovani, A. Cumulative effect of predisposing genotypes and their interaction with modifiable factors on the risk of ischemic stroke in young adults. *Stroke*. 2005;36:533-9.
 47. Pezzini AG, M.; Iacoviello, L.; Del Zotto, E.; Archetti, S.; Giossi, A.; Padovani, A. Inherited thrombophilia and stratification of ischaemic stroke risk among users of oral contraceptives. *J Neurol Neurosurg Psychiatry*. 2007;78:271-6.
 48. Press RDL, X. Y.; Beamer, N.; Coull, B. M. Ischemic stroke in the elderly. Role of the common factor V mutation causing resistance to activated protein C. *Stroke*. 1996;27:44-8.
 49. Pullmann R, Jr.; Skerenova, M.; Lukac, J.; Hybenova, J.; Melus, V.; Kubisz, P.; Rovensky, J.; Pullmann, R. Factor V Leiden and prothrombin G20210A mutations and the risk of atherothrombotic events in systemic lupus erythematosus. *Clin Appl Thromb Hemost*. 2004;10:233-8.
 50. Ranellou KP, A.; Kyriazopoulos, P.; Batistatou, A.; Evangelou, A.; El-Aly, M.; Zis, P.; Tavernarakis, A.; Charalabopoulos, K. Polymorphisms in prothrombotic genes in young stroke patients in Greece: a case-controlled study. *Blood Coagul Fibrinolysis*. 2015;26:430-5.
 51. Ridker PMH, C. H.; Lindpaintner, K.; Stampfer, M. J.; Eisenberg, P. R.; Miletich, J. P. Mutation in the gene coding for coagulation factor V and the risk of myocardial infarction, stroke, and venous thrombosis in apparently healthy men. *N Engl J Med*. 1995;332:912-7.
 52. Ridker PMH, C. H.; Miletich, J. P. G20210A mutation in prothrombin gene and risk of myocardial infarction, stroke, and venous thrombosis in a large cohort of US men. *Circulation*. 1999;99:999-1004.
 53. Ripoll LM, E.; Woimant, F.; Drouet, L. O. Different contribution of factor V Leiden-associated hypercoagulability to ischaemic cerebrovascular disease in the elderly. *Blood Coagul Fibrinolysis*. 1997;8:253-4.
 54. Romdhane NBB, H.; Lahmar, M.; Mahjoub, S.; Manai, Z. Deficiency of coagulation inhibitors in young adults with ischemic stroke. *Journal of Thrombosis and Haemostasis*. 2011;9:899.
 55. Rubattu SDA, E.; Nitsch, D.; Gigante, B.; Zanda, B.; Stanzione, R.; Evangelista, A.; Pirisi, A.; Rosati, G.; Volpe, M. Polymorphisms in prothrombotic genes and their impact on ischemic stroke in a Sardinian population. *Thrombosis and Haemostasis*. 2005;93:1095-100.
 56. Rubattu SS, R.; Ferrari, M.; Evangelista, A.; Beccia, M.; Stanzione, R.; Assenza, G. E.; Volpe, M.; Rasura, M. A role of TNF- α gene variant on juvenile ischemic stroke: A case-control study. *European Journal of Neurology*. 2005;12:989-93.
 57. Sastry SR, G.; Morris, J.; Taberner, D.; Cherry, N.; Heagerty, A.; McCollum, C. Young Adult Myocardial Infarction and Ischemic Stroke: the role of paradoxical embolism and thrombophilia (The YAMIS Study). *J Am Coll Cardiol*. 2006;48:686-91.
 58. Shi CK, X.; Wang, Y.; Zhou, Y. The coagulation factor V Leiden, MTHFRC677T variant and eNOS 4ab polymorphism in young Chinese population with ischemic stroke. *Clin Chim Acta*. 2008;396:7-9.

59. Slooter AJCR, F. R.; Tanis, B. C.; Kemmeren, J. M.; Van Der Graaf, Y.; Algra, A. Prothrombotic conditions, oral contraceptives, and the risk of ischemic stroke. *Journal of Thrombosis and Haemostasis*. 2005;3:1213-7.
60. Smiles AMJ, N. S.; Tang, Z.; Arnold, A.; Cushman, M.; Tracy, R. P. No association of plasma prothrombin concentration or the G20210A mutation with incident cardiovascular disease: results from the Cardiovascular Health Study. *Thromb Haemost*. 2002;87:614-21.
61. Supanc VS, Z.; Vukasovic, I.; Solter, V. V.; Zavoreo, I.; Kes, V. B. The role of classic risk factors and prothrombotic factor gene mutations in ischemic stroke risk development in young and middle-aged individuals. *J Stroke Cerebrovasc Dis*. 2014;23:e171-6.
62. Szolnoki ZS, F.; Kondacs, A.; Szabo, M.; Fodor, L.; Bene, J.; Melegh, B. Evaluation of the modifying effects of unfavourable genotypes on classical clinical risk factors for ischaemic stroke. *J Neurol Neurosurg Psychiatry*. 2003;74:1615-20.
63. Tatarsky PFK, A. M.; Kravchenko, S. A.; Shulzenko, D. V.; Kuznetsova, S. M.; Livshits, L. A. Ischemic stroke in Ukrainian population: Possible involvement of the F2 G20210A, F5 G1691A and MTHFR C677T gene variants. *Biopolymers and Cell*. 2010;26:299-305.
64. They-They TPB, O.; Slassi, I.; Rafai, M. A.; Katumbay, D. T.; Nadifi, S. Prothrombin G20210A and factor V Leiden polymorphisms in stroke. *J Mol Neurosci*. 2012;46:210-6.
65. Tupitsyna TVB, E. A.; Kravchenko, S. A.; Tatarsky P. F.; Shetova, I. M.; Shamalov, N. A.; Kuznetsova, S. M.; Shul'Zhenko, D. V.; Skvortsova, V. I.; Slominsky, P. A.; Livshits, L. A.; Limborska, S. A. Comparative analysis of associations between polymorphic variants of the F2, F5, GP1BA, and ACE genes and the risk of developing stroke in Russian and Ukrainian populations. *Molecular Genetics, Microbiology and Virology*. 2013;28:8-14.
66. Voetsch BD, B. P.; Camargo, E. C.; Massaro, A.; Bacheschi, L. A.; Scaff, M.; Annichino-Bizzacchi, J. M.; Arruda, V. R. Inherited thrombophilia as a risk factor for the development of ischemic stroke in young adults. *Thromb Haemost*. 2000;83:229-33.
67. Wypasek ES, E.; Pieculewicz, M.; Podolec, P.; Undas, A. Factor XIII Val34Leu polymorphism and ischaemic stroke in patients with patent foramen ovale. *Thromb Haemost*. 2009;102:1280-2.
68. Zimba SN, P. M.; Lakhi, S.; Atadzhanov, M. HIV infection, hypercoagulability and ischaemic stroke in adults at the University Teaching Hospital in Zambia: a case control study. *BMC Infect Dis*. 2017;17:354.