

Supplementary Tables

Table 1. Patient characteristics of the TOAST

	LAA (n=139)	CE (n=35)	SVD (n=31)	OE (n=16)	UE (n=78)	Total (%) (n=299)
Age ($\bar{x}\pm SD$)	59.9±11.1	64.7±17.8	55.6±8.6	40.2±15.0	57.2±13.3	58.3±13.5
Male (%)	85.6	60.0	71.0	68.8	76.9	77.9
Hypertention (%)	40.3	31.4	45.2	6.3	43.6	38.8
Diabetes (%)	20.1	17.1	12.9	6.3	16.7	17.4
Dyslipidemia (%)	70.5	69.0	63.3	64.3	65.8	68.1
Smoking (%)	56.1	45.7	51.6	37.6	52.6	52.6
CAHD (%)	10.1	22.9	16.1	6.3	9.0	11.7
MI (%)	4.3	5.7	3.2	0	3.9	3.3
AF (%)	0.7	17.1	0	0	1.3	2.7

Table 2. Patient characteristics of the SPARKLE

	LAA (n=165)	CE (n=32)	SVD (n=29)	OE (n=17)	UE (n=56)	Total (%) (n=299)
Age ($\bar{x}\pm SD$)	60.2±11.0	64.0±18.4	57.0±8.2	41.2±15.0	54.8±14.2	58.3±13.5
Male (%)	86.1	53.1	69.0	64.7	76.8	77.9
Hypertention (%)	42.4	28.1	41.4	5.9	42.9	38.8
Diabetes (%)	22.4	15.6	13.8	5.9	8.9	17.4
Dyslipidemia (%)	70.6	59.3	67.9	60.0	67.3	68.1
Smoking (%)	57.6	34.4	48.3	35.2	55.4	52.6
CAHD (%)	10.3	18.8	10.3	5.9	14.3	11.7
MI (%)	3.6	3.1	3.4	0	5.4	3.3
AF (%)	0.6	18.8	0	0	1.8	2.7

Table 3. Patient characteristics of the CISS

	LAA (n=139)	CE (n=35)	SVD (n=31)	OE (n=16)	UE (n=78)	Total (%) (n=299)
Age ($\bar{x}\pm SD$)	60.0±10.8	64.8±17.9	55.7±8.5	41.2±15.0	49.4±14.8	58.3±13.5
Male (%)	83.4	61.1	68.4	64.7	75.0	77.9
Hypertention (%)	44.7	30.6	36.8	5.9	28.6	38.8
Diabetes (%)	20.1	16.7	10.5	5.9	10.7	17.4
Dyslipidemia (%)	68.2	66.7	76.5	60.0	68.0	68.1
Smoking (%)	57.8	44.5	42.1	35.2	42.8	52.6
CAHD (%)	10.6	19.4	10.5	5.9	14.3	11.7
MI (%)	3.5	5.6	0	0	3.6	3.3
AF (%)	1.0	16.7	0	0	0	2.7