

**IL-21 regulates SOCS1 expression in autoreactive CD8⁺ T cells but is not required
for acquisition of CTL activity in the islets of non-obese diabetic mice**

Andrew P. R. Sutherland^{1,2*}, Kate L. Graham^{1,2*}, Michelle Papadimitriou¹, Gaurang
Jhala^{1,2}, Prerak Trivedi^{1,2}, Tara Catterall¹, Stacey Fynch¹, Thomas W. H. Kay^{1,2}, Helen E.
Thomas^{1,2}

¹St Vincent's Institute of Medical Research, Melbourne, Australia

²The University of Melbourne, Department of Medicine, St. Vincent's Hospital, Fitzroy,
Victoria, Australia

*Authors contributed equally

Address correspondence to:

A/Prof Helen Thomas

St. Vincent's Institute of Medical Research, 9 Princes Street, Fitzroy, VIC 3065,
Australia.

E-mail address: hthomas@svi.edu.au


Or

Dr. Andrew P. R. Sutherland


St. Vincent's Institute of Medical Research, 9 Princes Street, Fitzroy, VIC 3065,
Australia.

E-mail address: asutherland@svi.edu.au

PLN



ISLETS



Supplementary Figure 1: Quantification of CFSE dilution and hCD4 reporter expression in transferred 8.3 T cells in NOD pancreatic lymph nodes and islets.

Pancreatic lymph nodes and islets were isolated 5 days after 8.3 splenocyte transfer and

anti CD8 antibody staining and CFSE labelling used to identify the transferred 8.3 T cells.