


Supplemental Information


**A Bacterial Pathogen Senses Host Mannose
to Coordinate Virulence**

Lifan Wei, Haoxian Qiao, Brandon Sit, Kaiyu Yin, Guanhua Yang, Ruiqing Ma, Jiabao Ma, Chun Yang, Jun Yao, Yue Ma, Jingfan Xiao, Xiaohong Liu, Yuanxing Zhang, Matthew K. Waldor, and Qiya Wang

1 **Supplemental Materials**



4 **Supplemental Figure S1. Validation of the properties of the MKGR transposon insertion**
5 **mutants.** Related to Figure 1. **(A)** Antibiotic resistance of 6 insertion mutants (ETAE_1464,
6 ETAE_1797 (*hemR*), ETAE_2071 (*evrA*), ETAE_2200, ETAE_2366 (*hybA*), and EATE_3119)
7 that are predicted to generate active transcriptional fusions and 6 insertion mutants
8 (ETAE_1456 (*yoaA*), ETAE_1508 (*adhE*), ETAE_1591 (*gloA*), ETAE_2071 (*evrA*),
9 ETAE_2078, and ETAE_2022) that do not create transcriptional fusions. The 6 insertions
10 creating transcriptional fusions are resistant to Km as well as Col and Gm, whereas the latter 6
11 are not resistant to Km. **(B)** Detection of GFP and RFP fluorescence in Km^R and Km^S Tn
12 insertion mutants. Scale bar represents 5 μM.




15


16 **Supplemental Figure S2. Work flow for construction of the defined mutant library.**

17 Related to Figure 1. **(A)** Vector map of pMKGR. **(B)** *E. coli* SM10 λ pir and *E. piscicida* EIB202
 18 (Δ P) were used as the conjugation donor and recipient strains, respectively. To determine the
 19 transposon insertion sites, TAIL-PCRs were performed to amplify the junctions of the genome
 20 and transposon and the PCR products were sequenced and analyzed. The mutants were
 21 re-arrayed into different subset libraries for various screens. The master libraries and the
 22 subset libraries were duplicated for -80°C storage or as working libraries. **(C)** Schematic of
 23 TAIL-PCR amplification used for determination of the transposon insertion sites. The primer
 24 pair Sp1/AB2 was used for the first round of PCR and the primer pair Sp2/ABS was used for
 25 the second round of PCR. The Seq2 primer was used for sequencing of the amplification products.
 26 **(D)** Distribution of mapped transposon insertions on the *E. piscicida* chromosome. The most

27 outward circle represents the genes encoded in the two strands of DNA. The second circle
 28 displays the distribution of the transposon insertions, with the red and green bars representing
 29 the forward and reverse orientation of insertions within genes, respectively. The inner purple
 30 and olive circles corresponded to TA and GC contents on genome, respectively. **(E)**
 31 Sequencing saturation plot of the defined mutant library. The number of new ORFs disrupted
 32 (brown diamonds) and the total number of ORFs disrupted in the library (blue circles) are
 33 shown.




34
 35 **Supplemental Figure S3. Correlation of experimental replicates from TIS experiments.**
 36 Related to Figure 1. The sequencing data from experimental replicates of input libraries ($n = 3$)
 37 and output libraries ($n = 3$) recovered after growth in DMEM ($n = 3$), J774A.1 ($n = 3$), and
 38 turbot ($n = 3$) are shown.
 39



40


41 **Supplemental Figure S4. Fitness of *E. piscicida* transposon mutants in DMEM,**
 42 **macrophages and turbot.** Related to Figure 1. (A) COG categories of depleted genes in the
 43 three conditions. The number of depleted genes in each category is shown; the percentages

44 shown are the fraction of the total number of depleted genes represented by the number of
 45 depleted genes/category. **(B)** Heatmap of relative abundances of mutants in indicated
 46 conditions. Colored lines represent FC values of the genes across the genome (N). Four gene
 47 clusters, associated with LPS synthesis, T3SS, NADH, and T6SS, exhibiting reduced
 48 abundance in the turbot outputs, are highlighted. **(C)** Schematic model of genes and pathways
 49 essential for *in vivo* and *in vitro* growth in *E. piscicida*. Yellow, green, and pink colors
 50 respectively represented the growth conditions of DMEM, J774A.1 cells, and in turbot. The
 51 genes required for growth in specific conditions were highlighted in the colored ovals. The
 52 upstream regulatory network of T3/T6SS as well as the putative mannose metabolic pathways
 53 were highlighted in a dashed box.




54
 55 **Supplemental Figure S5. Intracellular growth of various *E. piscicida* mutants in J774A.1**
 56 **and turbot primary macrophages.** Related to Figure 1. **(A)** Intracellular bacterial counts of
 57 WT or individual T6SS deletion mutants grown in primary turbot macrophages; the means and
 58 SD from 3 replicates are shown. **(B)** Competition assays between the indicated deletion
 59 mutant vs. WT(ΔP) in primary turbot macrophages. Data presented are the mean \pm SD from 6

60 replicates. * $P < 0.05$, *** $P < 0.001$ based on ANOVA followed by Bonferroni's
61 multiple-comparison post-test comparing the data to the values from the WT/ WT Δp
62 competitions. **(C-D)** Intracellular *E. piscicida* CFU recovered (C), and cytotoxicity (D) of WT,
63 $\Delta evrA$ and the complemented strain towards J774A.1 cells. After incubation in DMEM for 12 h,
64 WT, $\Delta evrA$ and the complemented strain were used to infect J774A.1 cells at an MOI of 10.
65 Cytotoxicity was detected at 3 and 6 hours post infection using an LDH assay. Data were
66 shown as the mean \pm SEM of results for triplicate cultures or assays. *, $P < 0.05$ based on
67 student's *t*-test. **(E)** Western blot analysis of EseB and EvpP expression and secretion in the
68 indicated strains. The cell lysates (WCP) and extracellular proteins (ECPs) were analyzed by
69 western blotting with the indicated antibodies. DnaK levels were used as the loading control.




70 **Supplemental Figure S6. EvrA is a DeoR family protein.** Related to Figure 3. **(A)** Domain
 71 structure of EvrA. **(B)** The protein network of EvrA as predicted by STRING includes
 72 PTS-related proteins, e.g. *crr*, *nagE*, *fruK*, *ptxA*, *ptsN*, *sgcA*, 2301 (Fructose-specific PTS
 73 system IIBC component), 2090 (glucose-specific IIBC subunit), and 1718
 74 (Beta-glucoside-specific PTS system components IIABC). **(C)** Phylogenetic tree of EvrA
 75 multiple sequence alignments. The phylogenetic tree was inferred with a Neighbor-Joining
 76 algorithm built in MEGA (v6.0.6). **(D)** Conservation of EvrA in other bacterial species. The
 77 conserved residues are highlighted with various colors. The residues R7, R178 and R221 as
 78

79 well as the N-terminal HTH were as indicated. (E) The sequence alignment of the N-terminal
 80 HTH domain of EvrA. The conserved R7 residue was indicated by an asterisk. (F) Sequence
 81 alignment and predicted secondary structures between the indicated regions of EvrA (query)
 82 with the model RpiA (1LK7) (template).



83
 84 **Supplemental Figure S7. Mannose stimulates evrA-dependent induction of esrB**
 85 **expression.** Related to Figure 4. (A) Optical density at 600 nm (OD_{600}) of 24 h cultures of WT,
 86 $\Delta evrA$ and $\Delta manX$ in DMEM medium supplemented with 5 mg/mL of the indicated sugar.
 87 Results shown are mean \pm SD from 3 replicates, * $P < 0.05$; ***, $P < 0.001$ based on the
 88 student's *t*-test. (B) P_{esrB} -luxAB activity in the indicated strains grown in DMEM supplemented
 89 with glucose (glu) or mannose (man). (C) Auto-aggregation and protein expression in *E.*
 90 *piscicida* strains bearing an empty FLAG vector or a vector with EvrA-FLAG. (D) EMSA of


91 EvrA binding to *esrB* promoter in the presence of increasing concentrations of mannose. **(E)**
92 qRT-PCR analyses of *manX* and *evrA* transcript levels in cells grown in glucose or mannose
93 compared to those in WT cells grown in glucose. *gyrB* was used as a control. $n = 3$, * $P < 0.05$
94 based on Student's *t*-test. NA, not applicable. **(F)** EMSA of EvrA binding to the predicted *manX*
95 promoter (P_{manX}). 5-fold excess of un-labeled P_{manX} probe was added to lane to confirm the
96 specificity of the mobility shift. The putative binding motif in P_{manX} was mutated (T to G and A to
97 C) in P_{manX}^{Mut} .



98

99 **Supplemental Figure S8. Binding behaviors of EvrA or EvrA^{R141A} binding to esrB in the**
100 **presence of various sugars.** Related to Figure 5. (A-G) Isothermal titration calorimetry (ITC)
101 analyses of interaction of EvrA protein with mannose-6P (man-6P) (A), mannose-1P (man-1P)
102 (B), mannose (man) (C), and GDP-mannose (GDP-man) (D). EvrA^{R221A} (E), EvrA^{R7A} (F), and
103 EvrA^{R178A} (G) were also tested for interactions with man-6P. The top panel shows the raw

104 calorimetric data for the interaction and bottom panel the integrated heat changes, corrected
105 for heat of dilution, and fitted to a single-site binding model. K_d , dissociation constant. (H)
106 EMSA of EvrA or EvrA^{R221A} binding to *esrB* promoter probe in the absence or presence of
107 excess man-6P. In EMSA experiments, 20 ng of each Cy5-labeled probe was added to the
108 EMSA reactions. B: bound DNA; U: unbound DNA.




109

110 **Supplemental Figure S9.** Related to Figure 5. Mass spectrometry of EvrA in complex


111 with Man-6P after denatured with the addition of formic acids (A and B) or in the native
 112 condition (non-formic acid-treated) (C and D). Mass spectrometry B and D were zoomed in
 113 from the respective peak clusters with $z = 7$ (A and B).

114



115

116 **Supplemental Figure S10. Characterization of critical Arg residues in EvrA regulation of**
 117 **virulence gene expression.** Related to Figure 5. **(A)** Western blot of expression levels of
 118 EvrA or EvrA-HA in various *E. piscicida* strains used in Figures 5 and 6. DnaK was used as a
 119 loading control. **(B)** Auto-aggregation phenotypes of WT, ΔevrA, ΔevrA⁺, and various ΔevrA
 120 strains expressing R7A, R178A and R221A of EvrA variants. **(C)** Detection of eseB expression
 121 in HeLa cells infected by the indicated *E. piscicida* strains carrying a P_{eseB}-EGFP reporter
 122 plasmid. Bacteria were pre-incubated in glucose or mannose before infection at an MOI of 10
 123 for 8 h. mCherry fluorescence (red) was used to identify intracellular bacterial cells. DAPI was
 124 used to label cell nuclei (blue). Scale bar represents 10 μM. Quantification results are shown
 125 for at least 9 cells. ***P < 0.0001 based on Student's t-test.



126

127 **Supplemental Figure S11. Activation of *esrB* expression by Mannose of physical**
128 **concentrations.** Related to Figure 6. Chromosomal P_{esrB} -*luxAB* reporter activity in WT grown
129 for 12 h in DMEM medium supplemented with mannose at indicated concentrations. Data
130 shown are the mean \pm SEM of results for triplicate assays. * $P < 0.05$; **, $P < 0.01$ based on
131 the student's *t*-test.

132

Table S1 Characteristics of the *E. piscicida* EIB202 derived defined transposon mutant library

Data set	N
Master library	
Transposon mutants isolated	34,560
Mutants with well-defined insertion locations	24,470
Mutants in unique genomic locations	20,346
Insertions inside ORFs	18,128
Insertions between ORFs	6,342
Insertions inside 5-85% ORFs	14,028
Insertions outside 5-85% ORFs	4,100
Transcriptional fusion	12,295
Transposon with - frame insertions ^a	12,383
Transposon with + frame insertions ^b	12,087
Annotated ORFs in <i>E. piscicida</i>	3,599
ORFs disrupted	2,806
ORFs without hits	793
Average hits per annotated ORF	5.04
Predicted essential genes	496
Conditional essential genes/ intergenic regions in DMEM medium	52/20
Conditional essential genes/ intergenic regions in J774A.1	67/62
Conditional essential genes/ intergenic regions in turbot	258/108
Subset libraries ^c	
1 st NR subset (single mutant for each disrupted ORF) ^d	2,759
2 nd NR subset (single mutant for each disrupted ORF) ^e	2,235
3 rd NR subset (Tn transcriptionally fused to the disrupted gene) ^f	3,705
4 th NR subset (mutants with intergenic insertions) ^g	2,305
5 th NR subset (composite library of equally mixed 1 st , 2 nd , and 4 th subsets)	7,299

^{a,b} "+" and "-" indicates that the direction of transposon insertion is consistent with or reverse with the direction of gene, respectively.

^c the number indicated the mutants included in the specific subset:

^{d,e} No. 1 and 2 non-redundant (NR) libraries are parallel containing distinct insertion (20-80% region in the gene) mutants in the same allelic genes:

^{f,g} No more than two transcriptional fusion mutants for each disrupted ORF were selected for the 3rd and 4th subset.

Table S7 List of 5th subset of defined transposon insertion mutant library

Name of Composite Collection	Annotation
1st subset composite collection	composite collection of 1st subset, Col' and Gm', consist of 10 μ l bacterium culture which contains 2×10^6 c.f.u of 2,759 defferent mutants.
2nd subset composite collection	composite collection of 2nd subset, Col' and Gm', consist of 10 μ l bacterium culture which contains 2×10^6 c.f.u of 2,235 defferent mutants.
4th subset composite collection	composite collection of 4th subset,Col' and Gm',consist of 10 μ l bacterium culture which contains 2×10^6 c.f.u of 2,305 defferent mutants.

Table S8 Conditional essential genes grown in DMEM medium

Gene_ID	Gene	Annotation-nr	No.TA	LB_01	LB_02	LB_03	DMEM_01	DMEM_02	DMEM_03	DMEM/LB	P value
ETAE_0075	waaQ	heptosyl III transferase	45	260	311	282	37.63220892	35.3103448	43.93933988	-2.856865972	8.37407E-05
ETAE_0076	walW	lipopolysaccharide biosynthesis	46	211	178	154	26.22850925	20.50434783	18.69759144	-3.033742991	0.000673513
ETAE_0078	walR	putative glycosyltransferase	52	254	259	309	45.61479869	64.93043478	33.65566459	-2.503221036	0.000337238
ETAE_0079	wabK	putative glycosyltransferase	78	161	172	160	24.2898803	16.5173913	15.61248885	-3.104561798	6.61277E-06
ETAE_0083	rfaD	ADP-L-glycero-D-manno-heptose-6-epimerase	45	3	2	4	0.540369967	0	0.934879572	-2.008433752	0.015635015
ETAE_0185	purD	phosphoribosylamine-glycine ligase	37	227	225	199	42.19368879	52.4	47.67885816	-2.186744767	5.48257E-05
ETAE_0186	purH	ribosylaminomimidazole carboxamide formyltransferase/IV	51	780	741	680	167.6343852	119.6086957	153.3202498	-2.318332308	5.38948E-05
ETAE_0204	pgi	glucose-6-phosphate isomerase	60	50	45	55	2.04369967	2.04369967	2.04369967	-2.118770112	0.000377832
ETAE_0246	0	hypothetical protein	9	222	272	288	3.42109902	4.556521739	3.739518287	-5.943288669	0.000204106
ETAE_0358	purA	adenylosuccinate synthase	48	52	46	64	13.68443961	12.53043478	12.15343443	-2.052408584	0.001674465
ETAE_0368	rplI	50S ribosomal protein L9	15	32	25	36	1.140369967	4.556521739	2.804638715	-3.296101516	0.001275789
ETAE_0459	folB	bifunctional dihydronopterin aldolase/dihydronopterin tri	10	40	31	37	2.280739935	1.139130433	1.869759144	-4.114148411	0.000221066
ETAE_0593	carA	carbamoylphosphate synthase small subunit	26	183	168	180	46.75516866	43.28695652	32.72078501	-2.105702307	2.44036E-05
ETAE_0594	carB	carbamoylphosphate synthase large subunit	89	1,304	1,340	1,268	366.0587595	389.5826087	203.8037467	-2.026405802	9.22998E-05
ETAE_0771	pspE	rhodanese-related sulfurtransferase	9	292	304	258	55.44069641	40.49565217	39.74843886	-2.653415605	0.000519648
ETAE_0774	purL	phosphoribosylformylglycinamide synthase	74	1,631	1,646	1,640	458.4287269	439.7043478	330.0124888	-2.000454698	6.86623E-06
ETAE_0796	prob	gamma-glutamyl kinase	27	260	265	263	60.43960827	54.67826087	56.09277431	-2.19596383	1.01296E-07
ETAE_0797	proA	gamma-glutamyl phosphate reductase	24	436	470	423	84.38737758	67.20869565	75.72524532	-2.54196643	1.66033E-05
ETAE_0802	0	hypothetical protein	8	225	226	197	28.50924918	35.31034348	35.52542373	-2.693134931	4.85458E-05
ETAE_0808	0	hypothetical protein	3	38	61	43	10.52665941	12.6695652	11.89295272	-2.113968356	0.013236819
ETAE_0837	0	putative phospholipid biosynthesis acyltransferase	26	473	452	476	119.3014146	110.6173913	118.3621766	-2.211522028	5.18511E-05
ETAE_0891	0	hypothetical protein	6	12	14	6	1.140369967	14.177391304	2.804638715	-2.562229382	0.030152916
ETAE_1010	acrB	RND family, acridine/multidrug efflux pump/acriflavin resist	104	809	875	824	84.38737758	103.6608696	114.0553078	-3.049164806	4.70073E-06
ETAE_1011	acrA	efflux transporter, RND family, MFP subunit	42	458	423	424	109.4755169	63.79130435	99.09723461	-2.256245111	4.39294E-05
ETAE_1020	dnaX	DNA polymerase III, subunits gamma and tau	34	20	28	23	1.140369967	1.139130433	4.674397858	-3.166150504	0.001156244
ETAE_1053	purB	adenylosuccinate lyase	29	344	300	333	66.14145811	54.67826087	67.31132917	-2.369860097	4.67743E-05
ETAE_1086	purM	phosphoribosylformylglycinamide cyclo-ligase	20	307	300	318	77.54515778	67.20869565	68.24620874	-2.111973611	2.70844E-06
ETAE_1098	purC	phosphoribosylaminomimidazole-succinocarboxamide synth	17	132	138	158	21.66702938	46.70434783	36.4603033	-2.016136161	0.000540506
ETAE_1248	pyrD	dihydroorotate dehydrogenase 2	47	187	231	209	39.91294886	58.09565217	43.0044603	-2.145275549	0.000316366
ETAE_1520	0	ferritin Dps family protein	22	877	892	871	106.0544007	186.8173913	208.4781445	-2.394696475	2.33859E-05
ETAE_1563	0	hypothetical protein	7	166	190	173	40.43960827	39.23478261	34.50669045	-2.014143443	9.14137E-05
ETAE_1689	pdxH	pyridoxamine 5'-phosphate oxidase	20	27	21	13	1.140369967	1.139130433	5.609277431	-2.812698256	0.014153307
ETAE_1867	pspA	rock protein A (IM30), suppresses sigma54-dependent tran	10	715	705	688	21.66702938	38.73043478	44.87421945	-4.310206022	3.81692E-07
ETAE_1894	0	hypothetical protein	11	1,203	1,245	1,143	86.66811752	72.90434783	101.9018733	-3.774829304	3.48145E-06
ETAE_1951	wza	polysaccharide export-related protein	14	1,372	1,373	1,295	117.4581066	76.32173913	125.2738626	-3.658414808	2.03634E-06
ETAE_1961	0	hypothetical protein	11	75	84	75	9.122959739	9.113043478	5.609277431	-3.244205294	2.74928E-05
ETAE_2082	pyrF	OMP decarboxylase; OMPDCase; OMPdecase	16	87	76	74	23.94776931	19.36521739	14.95807315	-2.00373805	0.000250875
ETAE_2138	0	hypothetical protein	8	244	245	309	25.08813928	18.22608696	31.78590544	-3.392868407	0.00038499
ETAE_2159	mukE	condesin subunit E	22	2	3	1	0	0	0	-2.885963624	0.021610065
ETAE_2160	mukF	chromosome segregation and condensation protein	46	6	9	7	1.140369967	0	0.934879572	-2.883327281	0.001283873
ETAE_2256	0	hypothetical protein	22	3	4	4	0	0	0	-3.566250389	0.000585947
ETAE_2262	0	hypothetical protein	12	9	9	7	0	0	0	-4.683175648	0.000117254
ETAE_2274	0	hypothetical protein	2	8	6	5	1.280739935	1.27826087	1.869759144	-2.653415605	0.010712579
ETAE_2391	0	aminotransferase class I and II	31	539	597	548	139.125136	119.6086957	82.26940232	-2.300582757	5.34221E-05
ETAE_2409	purF	amidophosphoribosyltransferase	56	996	1,017	881	249.7410229	213.0173913	164.5388046	-2.20403196	0.000103139
ETAE_2519	0	hypothetical protein	14	582	477	690	21.66702938	29.6173913	35.52542373	-4.316392181	0.00853337
ETAE_2700	purE	phosphoribosylaminomimidazole carboxylase catalytic subu	8	57	54	54	13.68443961	17.08695652	8.413916146	-2.047184308	0.000102292
ETAE_2701	purK	phosphoribosylaminomimidazole carboxylase	22	285	245	254	65.00108814	95.68695652	34.59054416	-2.000616641	0.000776871
ETAE_2736	0	inorganic polyphosphate/ATP-NAD kinase	23	10	4	7	0	1.139130435	0.934879572	-2.866993196	0.01891615
ETAE_3124	pyrI	aspartate carbamoyltransferase regulatory subunit	9	241	257	273	60.43960827	58.09565217	63.57181088	-2.076427757	3.19392E-05
ETAE_3125	pyrB	aspartate carbamoyltransferase	24	369	364	387	78.68552775	58.09565217	44.87421945	-2.616665381	1.3228E-05
ETAE_3261	aroK	shikimate kinase I	12	15	10	17	0	2.27826087	3.739518287	-2.569333197	0.008092284

Table S9 Conditional essential genes grown in J774A.1 cells

Gene_ID	Gene	Annotation-nr	No.TA	Input_01	Input_02	Input_03	J774A.1_01	J774A.1_02	J774A.1_03	J774A.1/Input	P value
ETAE_0147	ubiE	ubiquinone/menaquinone biosynthl	30	337	245.3632144	364.569027	52	49.01921317	17.51487414	-2.98699558	0.001843319
ETAE_0217	ubiA	4-hydroxybenzoate polyprenyltrar	26	10	11.10240789	4.873917473	1	0	1.251062439	-3.07361928	0.016011176
ETAE_0228	citA	sensor histidine kinase	51	15,575	15291.34639	14869.34743	2179	2095.571363	2468.346192	-2.761697089	6.29927E-07
ETAE_0314	groEL	chaperonin GroEL (HSP60 family	49	112	145.4415434	113.0748854	8	12.25480329	66.30630925	-2.086046428	0.012039748
ETAE_0340	poxA	lysyl-tRNA synthetase, class II	27	6	4.440963156	2.924350484	0	1.750686185	0	-2.40826283	0.024368157
ETAE_0345		hypothetical protein	8	8	6.661444735	6.823484463	1	0	0	-3.510960361	0.000370855
ETAE_0348		putative ATPase	19	15	9.992167102	7.798267957	0	0	1.251062439	-3.888652536	0.006983543
ETAE_0381	fbp	fructose-1,6-bisphosphatase	38	1,127	1101.1358863	1073.236628	311	248.5974382	226.4423014	-2.068955919	9.30721E-06
ETAE_0405	infB	translation initiation factor 2	73	2	1.110240789	3.899133979	0	0	0	-3.015807311	0.044556486
ETAE_0409	pnp	polyribonucleotide nucleotidyltran	66	223	135.4493763	180.3349465	31	26.26029277	12.51062439	-2.93240182	0.003842202
ETAE_0494		conserved hypothetical protein	34	2,133	2167.19002	2076.288844	422	434.1701738	321.5230467	-2.435754806	2.58749E-06
ETAE_0578	nhaA	Na+/H+ antiporter	31	9	3.330722367	11.69740194	0	0	0	-4.665765179	0.031036263
ETAE_0600	ksgA	dimethyladenosine transferase (rl	19	45	68.83492892	49.71395823	3	7.002744739	8.757437071	-3.055157261	0.003097679
ETAE_0735	glnD	PII uridylyl-transferase	67	59	97.70118944	111.1253184	17	5.252058554	27.54037267	-2.428617931	0.029865787
ETAE_0743	cdsA	CDP-diglyceride synthetase	25	15	11.10240789	3.899133979	0	0	0	-4.933236128	0.034874421
ETAE_0859	esaH	putative type III secretion apparat	6	326	445.2065564	463.9969435	35	43.76715462	61.3020595	-3.131867732	0.001128871
ETAE_0860	esaG	putative type III secretion system	7	1,716	1646.48709	1628.86322	298	320.3755718	130.1104936	-2.735750711	2.74968E-05
ETAE_0861	esrC	putative transcriptional regulator	19	9,378	9548.070786	9813.145441	833	85.0773102	870.7394573	-3.500291293	2.68681E-07
ETAE_0881	esaS	type III secretion apparatus	6	1,971	2061.717145	2830.771268	475	540.9620311	582.9950964	-2.101048748	0.003078484
ETAE_0882	esaT	type III secretion system EscT ho	24	8,523	8790.886568	10104.60571	1309	1253.491308	942.0500163	-2.967488623	9.18513E-05
ETAE_0883	esaU	type III secretion system EscU ho	24	25,085	23348.36379	25658.25115	1773	1685.910796	2047.989212	-3.749764336	5.33059E-06
ETAE_0884		putative transglycosylase signal p	29	22,079	22106.00435	23736.95288	1780	1755.938243	2249.410265	-3.553164088	3.45632E-06
ETAE_0885	esrA	two-component sensor/regulator	69	28,010	26593.59762	27164.29165	2687	2718.815645	3121.400785	-3.261240893	6.01382E-07
ETAE_0886	esrB	two-component sensor/regulator	15	3,963	3894.724688	3782.159959	423	395.6550778	499.173913	-3.141890979	5.98632E-07
ETAE_0956	ahpC	alkyl hydroperoxide reductase, sn	23	82	81.0475776	115.9992359	1	3.50137237	48.79143511	-2.367407153	0.017561024
ETAE_0975	ribH	riboflavin synthase beta-chain	14	3	5.551203945	2.924350484	0	0	0	-3.654629928	0.012441266
ETAE_0981	xseB	exonuclease VII small subunit	6	193	203.1740644	212.5020818	40	26.26029277	48.79143511	-2.394045236	4.34837E-05
ETAE_0982	thil	thiamine biosynthesis ATP pyropl	41	622	659.4830287	643.3571065	116	140.0548948	125.1062439	-2.332897594	2.46509E-06
ETAE_1011	acrA	efflux transporter, RND family, Mf	42	1,836	1708.660574	1838.441671	397	409.6605672	455.4717228	-2.092453133	3.56829E-05
ETAE_1179		hypothetical protein	3	1	3.330722367	3.899133979	0	0	0	-3.212450606	0.034395192
ETAE_1207	neuB	sialic acid synthase	81	246	253.1348999	194.9566989	15	24.50960659	96.33180778	-2.345389683	0.004132014
ETAE_1258		hypothetical protein	6	27	38.85842762	23.39480387	6	3.50137237	7.012749264	-2.434590438	0.020812777
ETAE_1439	ruvA	holliday junction resolvosome, DN	10	25	26.64577894	12.67218543	1	0	3.753187316	-3.503834232	0.011998685
ETAE_1489		hypothetical protein	9	85	89.92950392	75.05832909	0	0	60.05099706	-2.040847657	0.036458424
ETAE_1518	cls	cardiolipin synthetase	50	214	235.3710473	269.0420445	19	17.50686185	50.04249755	-3.038873742	0.000391098
ETAE_1553	cysB	transcriptional regulator	35	22	8.881926313	10.72261844	1	0	0	-4.407065168	0.029325456
ETAE_1554	cspC	cold shock protein	6	67	44.40963156	97.47834947	4	8.753430924	33.77868585	-2.139344416	0.039699662
ETAE_1617		hypothetical protein	5	68	66.61444735	41.91569027	11	8.753430924	16.27231121	-2.289212007	0.012457067
ETAE_1654		hypothetical protein	5	34	23.31505657	32.16785532	4	0	2.502124877	-3.597042675	0.001458571
ETAE_1675		hypothetical protein	5	59	31.08674209	34.11742231	6	5.252058554	12.01699902	-2.419068426	0.037475334
ETAE_1725		chain length determinant protein	60	131	95.48070786	88.70529801	7	1.750686185	27.52337365	-3.084356147	0.003691597
ETAE_1787		hypothetical protein	17	98	61.0632434	52.63830871	12	5.252058554	31.27656097	-2.100472168	0.026801951
ETAE_1792	aroH	3-deoxy-7-phosphoheptulonate s:	25	208	112.1343197	185.208864	29	22.7589204	36.28081072	-2.507241479	0.008836873
ETAE_1831		hypothetical protein	5	11	33.30722367	20.47045339	0	0	0	-6.048996085	0.026652392
ETAE_1906		hypothetical protein	33	1,832	2100.575573	1334.478604	300	309.8714547	447.8803531	-2.315023038	0.003620538
ETAE_1967		HpcH/HpaI aldolase family proteir	9	116	113.2445605	68.234844643	9	10.50411711	32.55312194	-2.516609241	0.03758009
ETAE_2030		hypothetical protein	10	21	18.87409341	7.798267957	0	1.750686185	0	-4.138930251	0.020389422
ETAE_2176	cmk	cytidylate kinase	19	9	2.220481578	7.798267957	0	0	0	-4.34859903	0.002066975
ETAE_2305		hypothetical protein	7	57	59.95300261	75.05832909	12	14.00548948	13.77018634	-2.272479392	0.002027673
ETAE_2374	nuoM	NADH:ubiquinone oxidoreductase	45	106	153.2132289	184.2340805	17	15.75617566	45.03824779	-2.495901617	0.007809592
ETAE_2376	nuoK	NADH:ubiquinone oxidoreductase	7	28	34.41746446	40.94090678	5	8.753430924	8.757437071	-2.199844664	0.002320357
ETAE_2377	nuoJ	NADH:ubiquinone oxidoreductase	12	43	72.16565129	76.03311258	0	15.01274926	3.582410392	0.007401223	
ETAE_2381	nuoF	NADH:ubiquinone oxidoreductase	29	81	74.38613287	94.55399898	5	5.252058554	8.757437071	-3.648998663	0.000220093
ETAE_2383	nuoD	NADH:ubiquinone oxidoreductase	49	220	229.8198433	233.9480387	2	3.50137237	135.1147434	-2.274364101	0.014988342
ETAE_2385	nuoA	NADH:ubiquinone oxidoreductase	15	22	27.755601973	39.96612328	5	8.753430924	1.251062439	-2.500223919	0.012525015
ETAE_2412	folC	bifunctional folylpolyglutamate sy	31	8	11.10240789	17.5461029	1	3.50137237	1.251062439	-2.491135459	0.021687059
ETAE_2549		hypothetical protein	6	19	28.86626052	36.0669893	2	0	0	-4.818341079	0.005853485
ETAE_2650	lipA	lipoate synthase	16	24	22.20481578	18.5208864	0	0	1.251062439	-4.865709546	0.000210904
ETAE_2745		phage/plasmid primase	3	45	59.95300261	51.66352522	13	10.50411711	12.51487414	-2.117579482	0.001483348
ETAE_2858	sdhC	succinate dehydrogenase cytochr	7	109	65.50420656	45.81482425	7	7.002744739	8.757437071	-3.220389352	0.024439934
ETAE_2859	alaS	alanyl-tRNA synthetase	72	12	15.54337105	10.72261844	1	0	1.251062439	-3.61155638	0.001134151
ETAE_2873	rpoS	RNA polymerase sigma factor	24	830	749.4125326	865.6077433	42	31.51235133	41.28506048	-4.401035826	2.32167E-05
ETAE_3055		truncated integrase	5	11	13.32288947	17.5461029	1	1.750686185	5.004249755	-2.306271159	0.00621337
ETAE_3147		hypothetical protein	7	113	87.70902234	86.75573102	26	15.75617566	6.255312194	-2.558665358	0.001539308
ETAE_3211	secY	preprotein translocase subunit Se	67	11	33.30722367	23.39480387	0	0	0	-6.111348871	0.022930809
ETAE_3474		transcriptional regulator, TelR fan	19	14,913	14799.50972	15576.06546	3132	3119.722781	2394.533508	-2.38865778	3.73911E-06
ETAE_3539	pstC	ABC-type phosphate transport sy	37	195	178.748767	160.8392766	58	43.02470265	15.01274926	-2.204925796	0.001870957

Table S10 Conditional essential genes grown in turbot

Gene_ID	Gene	Annotation-n	No.TA	Input_01	Input_02	Input_03	Turbot_01	Turbot_02	Turbot_03	Turbot/Input_P value
ETAE_0023		conserved hypothetical protein	20	1058.3721	1049.0489	1100.714	28.6	70.41	15.9	-4.767325 1.452E-06
ETAE_0026	gltS	Na ⁺ /glutamate symporter	35	684.67741	628.76223	643.16231	205.43	132.02	101.76	-2.147757 0.000134
ETAE_0074	waaG	lipopolysaccharide core biosynthesis protein	40	181.2634	175.95336	85.25138	18.01	14.62	13.14	-3.191256 0.013268
ETAE_0075	waaQ	heptosyl III transferase	45	353.07706	332.72696	372.30033	64.09	102.28	89.35	-2.036103 7.522E-05
ETAE_0076	walW	lipopolysaccharide biosynthesis	46	133.15558	129.25484	112.22967	16.22	20.61	18.16	-2.703136 8.352E-05
ETAE_0101	rfr	UDP-N-acetylmuramyl pentapeptide phosphotransferase	50	2675.9977	2650.975	2437.7578	52.01	44.01	65.86	-5.392832 5.048E-06
ETAE_0103	wecB	UDP-N-acetylglucosamine 2-epimerase	35	627.11984	668.78953	675.53625	91.01	28.61	111.3	-3.077361 3.747E-05
ETAE_0104	wecC	UDP-N-acetyl-D-mannosaminuronate dehydrogenase	31	525.74978	607.08077	547.11962	104.01	138.63	130.38	-2.162111 8.002E-05
ETAE_0105	rfbB	dTDP-D-glucose 4,6-dehydratase	36	768.00703	806.38339	858.98859	62.41	66.01	60.42	-3.666793 9.255E-06
ETAE_0106	rfaB	glucose-1-phosphate thymidylyltransferase	30	391.73513	376.08987	403.59514	13	4.4	9.54	-5.293731 1.371E-06
ETAE_0107	wecD	TDP-D-fucosamine acetyltransferase	15	214.76707	219.31627	226.61759	5.2	0	0	-6.338766 5.846E-07
ETAE_0108	wecE	TDP-4-oxo-6-deoxy-D-glucose transaminase	38	814.39672	751.34585	875.17556	13	15.4	0	-6.282288 2.396E-05
ETAE_0109	wzxE	membrane protein involved in the export of O-antigen and teichoic acid	45	214.676707	39.193402	21.582628	2.6	0	0	-3.928247 0.0111494
ETAE_0110	wecF	4-alpha-L-fucosyltransferase	31	492.24612	497.00569	459.70979	15.6	0	3.18	-6.058856 2.96E-06
ETAE_0112	wecG	UDP-N-acetyl-D-mannosaminuronic acid transferase	25	1117.6478	1102.4187	1019.7792	241.83	195.84	171.72	-2.404738 1.821E-05
ETAE_0121		putative lipoprotein	6	66.148257	60.874856	69.064409	10.4	0	0	-3.893098 0.0001251
ETAE_0126	uvrD	DNA-dependent helicase II	62	1102.1846	1097.4152	1130.9297	161.22	44.01	254.4	-2.84912 0.0001013
ETAE_0131	recQ	ATP-dependent DNA helicase	54	1637.3841	1729.5131	1638.1215	411.99	452.54	273.44	-2.133979 3.214E-05
ETAE_0137	zntA	zinc/cadmium/mercury/lead-transporting ATPase	43	1553.1954	1618.6041	1587.4023	465.46	107.82	407.05	-2.275881 0.0003606
ETAE_0147	ubiE	ubiquinone/menaquinone biosynthesis methyltransferase	30	238.82098	226.82139	240.6463	5.2	8.8	3.18	-5.135375 9.936E-07
ETAE_0151	tatB	Sec-independent protein secretion pathway component	8	183.84061	205.97383	208.27236	52.01	33.01	54.06	-2.08087 0.0001185
ETAE_0152	tatC	twin arginine-targeting protein translocase	27	2030.8374	2067.2435	2218.6941	725.5	189.23	343.45	-2.325099 0.0005729
ETAE_0158	frd	FMN reductase	23	77.316144	72.549488	74.460066	5.2	0	12.72	-3.441806 6.336E-05
ETAE_0185	purD	phosphoribosylamine-glycine ligase	37	853.91386	869.75995	887.04601	137.82	211.24	149.46	-2.381725 9.029E-06
ETAE_0186	purH	bifunctional phosphoribosylaminoimidazolecarboxamide formyltransferase	51	1780.8485	1819.5745	1810.7825	161.22	402.67	181.26	-2.85535 3.779E-05
ETAE_0204	pgi	glucose-6-phosphate isomerase	60	47.248755	42.52901	57.193964	11.4	13	9.5	-2.022998 0.0010548
ETAE_0220	dgkA	diacylglycerol kinase	12	688.11368	659.61661	773.73721	101.41	178.23	76.32	-2.565206 0.0002151
ETAE_0340	poxA	lysyl-tRNA synthetase, class II	27	19.75857	17.511945	9.7121826	5.2	0	3.18	-2.134928 0.0193541
ETAE_0341		putative membrane transport protein	67	1832.3926	1883.785	1783.8042	440.87	363.07	499.27	-2.077357 2.351E-05
ETAE_0351	niaA	tRNA delta(2)-isopentenylpyrophosphate transferase	21	35.221799	35.857793	52.877438	7.8	0	12.72	-2.43238 0.0073778
ETAE_0360	rrrR	exoribonuclease R	72	3864.9481	3793.4209	3534.1553	731.15	992.38	878.04	-2.103805 2.203E-05
ETAE_0379		conserved hypothetical protein	16	156.35042	141.76337	140.28708	52.01	28.61	15.9	-2.149031 0.000633
ETAE_0384	argR	arginine repressor	13	42.953413	50.868032	62.589621	2.6	13.2	12.72	-2.338412 0.003075
ETAE_0396	greA	transcription elongation factor	9	322.1506	316.04892	296.76113	83.21	88.82	54.06	-2.048008 0.0003635
ETAE_0397	yhbY	RNA-binding protein	6	67.007325	58.373151	33.453073	0	0	3.18	-4.71076 0.0068413
ETAE_0406	rbfA	ribosome-binding factor A	11	18.899502	22.515358	34.532205	0	6.6	4.72	-2.74612 0.0343798
ETAE_0409	pnp	polyribonucleotide nucleotidyltransferase	66	61.852915	60.040956	69.064409	2.6	0	3.18	-4.465328 2.978E-05
ETAE_0410	nplI	lipoprotein NplI, contains TPR repeats	47	572.13947	538.7008	544.96135	132.62	96.82	101.76	-2.311358 8.343E-06
ETAE_0411	deAD	ATP-dependent RNA helicase	53	538.6358	509.51422	502.87523	117.02	72.61	118.54	-2.320235 2.427E-05
ETAE_0412		hypothetical protein	7	32.645494	35.857793	22.661759	10.4	4.4	3.18	-2.16613 0.0058712
ETAE_0479	prfC	peptide chain release factor RF-3	50	113.39701	103.40387	123.02098	0	0	0	-6.836353 3.687E-05
ETAE_0488	deoB	phosphopentomutase	40	1005.1099	1015.6928	1095.3184	249.63	211.24	265.71	-2.100558 0.0001505
ETAE_0494		conserved hypothetical protein	34	301.53296	316.88282	334.53073	96.21	24.2	111.3	-2.026051 0.0010837
ETAE_0497	rpoN	RNA polymerase factor sigma-54	43	734.50337	761.35267	721.9389	52.01	55.01	143.1	-3.133183 3.37E-05
ETAE_0501	kdsC	3-deoxy-D-manno-octulonate 8-phosphate phosphatase	14	133.15558	125.91923	111.15053	62.41	0	15.9	-2.198542 0.0079791
ETAE_0502	yrbH	D-arabinose 5-phosphate isomerase	24	161.50483	179.28896	213.66802	43.21	35.21	47.7	-2.110161 0.0008186
ETAE_0517	sspA	stringent starvation protein A	18	126.28304	136.75995	108.83401	2.6	2.2	25.44	-3.487708 0.0006291
ETAE_0530		hypothetical protein	10	188.99502	175.11945	160.79058	26	6.6	28.62	-3.039184 0.0001336
ETAE_0543	trpR	Trp operon repressor	8	123.70583	150.10239	123.73316	26	36.01	25.44	-2.216872 0.0034108
ETAE_0570	talD	transaldolase B	34	542.07208	527.86007	637.76665	111.82	33.01	50.88	-3.105849 0.0002753
ETAE_0600	ksgA	dimethyladenosine transferase (rRNA methylation)	19	23.194843	26.684869	16.186971	0	0	0	-4.524956 0.0020409
ETAE_0601	pdxA	4-hydroxythreonine-4-phosphate dehydrogenase	32	106.52447	121.74972	111.15053	0	0	0	-6.834681 1.494E-05
ETAE_0618	leuD	3-isopropylmalate dehydratase small subunit	17	376.2719	376.08987	367.98381	123.45	77.25	64.63	-2.065722 9.474E-05
ETAE_0627	fruR	DNA-binding transcriptional regulator	35	370.25842	356.91013	426.2569	49.41	129.82	47.7	-2.330406 0.0008546
ETAE_0629	mraW	S-adenosyl-methyltransferase	22	142.60533	128.42093	96.042694	0	0	12.72	-4.557121 0.0012091
ETAE_0644		conserved hypothetical protein	17	97.933783	108.40728	131.165403	39.01	13.2	6.36	-2.469452 0.0027042
ETAE_0661		hypothetical protein	6	15.463229	12.505822	15.10784	3.8	2.2	0	-2.84937 0.0116221
ETAE_0670	hpt	hypoxanthine-guanine phosphoribosyltransferase	24	88.484032	64.210466	97.121826	5.2	2.2	6.36	-3.914996 0.0013724
ETAE_0671	cynT	carbonic anhydrase	21	18.899502	24.183163	31.29481	7.8	6.6	0	-2.152826 0.0099271
ETAE_0697		hypothetical protein	11	18.899502	13.342435	20.503497	2.6	4.4	0	-2.478854 0.0037511
ETAE_0712	recD	exonuclease V subunit alpha	35	224.21682	230.157	192.08539	48.01	28.61	59.41	-2.24768 0.0016972
ETAE_0746	ompH	periplasmic chaperone	19	72.161735	80.88851	99.280088	7.8	2.2	6.36	-3.721214 0.0006499
ETAE_0751	rnhB	ribonuclease HII	14	139.16906	150.93629	139.20795	39.81	11	50.88	-2.077853 0.0027116
ETAE_0769		hypothetical protein	15	384.86258	401.10694	332.37247	52.01	114.42	76.32	-2.189963 0.0004505
ETAE_0774	purL	phosphoribosylformylglycinamide synthase	74	3371.843	3461.5279	3226.6029	702.1	889.2	833.17	-2.052885 4.836E-05
ETAE_0783	apbE	membrane-associated lipoprotein involved in thiamine biosynthesis	29	240.53912	245.16724	247.30891	2.6	4.4	34.98	-4.012843 2.956E-05
ETAE_0854	esal	putative type III secretion apparatus	30	660.6235	662.95222	673.37799	135.22	94.22	92.22	-2.786588 1.391E-05
ETAE_0856		conserved hypothetical protein	15	390.01699	428.62571	423.01951	0	19.8	0	-5.770577 7.7E-06
ETAE_0857	esaj	putative type III secretion system apparatus	22	693.26809	692.95222	731.65109	13	24.2	31.8	-4.859964 5.249E-06
ETAE_0859	esah	putative type III secretion apparatus	6	94.49751	80.88851	76.618329	2.6	0	3.18	-4.860157 0.0001157
ETAE_0860	esat	putative type III secretion system needle protein	7	180.40434	191.7975	209.35149	0	11	12.72	-4.451343 3.715E-05
ETAE_0861	esrC	putative transcriptional regulator	19	842.74597	830.56655	805.03202	5.2	35.21	15.9	-5.386703 5.671E-07
ETAE_0863	esad	type-III secretion protein	28	680.38207	611.25028	625.89621	10.4	0	9.54	-6.387494 7.656E-06
ETAE_0864	esAC	Type II secretion pathway, component PufD like protein	47	197.32798	2003.8669	1860.4225	156.4621	6.21	6.36	-6.358696 1.836E-06
ETAE_0865	esAB	two-component sensor/regulator	14	317.85526	361.07964	371.22122	26	26.4	22.26	-3.761404 3.832E-05
ETAE_0868	eseE	type III secretion system effector protein E	12	396.88954	441.13424	373.37946	7.8	4.4	15.9	-5.287189 4.019E-05
ETAE_0869	eseD	type III secretion system effector protein D	18	534.34046	527.86007	495.32131	148.22	74.81	15.9	-2.689367 0.0003948
ETAE_0870	eseC	type III secretion system effector protein C	30	445.85643	486.99886	498.5587	15.6	4.4	168.54	-2.904744 0.0016989
ETAE_0871	escA	type III secretion low calcium response chaperone	15	683.81834	688.80319	731.65109	13	46.21	168.54	-3.190974 0.0002289
ETAE_0872	esEB	EsP family secreted protein	22	752.5438	733	749.9				

ETAE_1021	hypothetical protein	7	47.248755	62.542662	57.193964	15.6	19.8	0	-2.146232	0.004279
ETAE_1025	adk adenylate kinase	22	18.899502	10.006826	16.186971	2.6	0	3.18	-2.453543	0.0095307
ETAE_1086	purM phosphoribosylformylglycinamide cyclo-ligase	20	851.33665	842.24118	887.04601	106.61	107.82	130.38	-2.893025	1.18E-06
ETAE_1095	gcvR glycine cleavage system transcriptional repressor	19	151.19602	136.75995	162.94884	44.21	8.8	34.98	-2.31861	0.0007491
ETAE_1098	purC phosphoribosylaminoimidazole-succinocarboxamide synthase	17	485.37357	444.46985	488.84652	49.41	39.61	114.48	-2.783393	0.0001232
ETAE_1128	hypothetical protein	15	290.36507	292.69966	268.70372	46.81	30.81	117.66	-2.107994	0.0013998
ETAE_1131	ptsl phosphoenopyruvate-protein phosphotransferase	43	47.248755	58.373151	59.352227	15.6	19.8	0	-2.129061	0.0038179
ETAE_1167	hypothetical protein	16	4.2953413	9.1729238	5.395657	0	0	2.18	-3.11323	0.0452115
ETAE_1169	hypothetical protein	122	56.698506	45.864619	39.927862	2.6	2.2	6.36	-3.361037	0.001007
ETAE_1201	galE UDP-glucose 4-epimerase	56	1262.8304	1299.2196	1101.7932	75.41	72.61	66.78	-4.073463	4.581E-05
ETAE_1207	neuA sialic acid synthase	81	24.91298	25.017065	25.899153	2.6	0	3.18	-3.166437	2.213E-05
ETAE_1208	neuA acylneuraminate cytidylyltransferase	118	42.094435	52.535836	39.927862	5.2	2.2	9.54	-2.786303	0.0009056
ETAE_1212	gnd 6-phosphogluconate dehydrogenase	48	768.8661	738.83732	789.92418	75.41	8.8	181.26	-3.099195	0.0002058
ETAE_1407	hemK N5-glutamine S-adenosyl-L-methionine-dependent methyltransferase	17	233.66657	223.48578	207.18144	96.21	15.4	0	-2.527635	0.0043904
ETAE_1445	msbB lipid A biosynthesis (KDO)2-(lauroyl)-lipid IVA acyltransferase	28	298.09669	341.89989	338.84726	57.21	17.6	15.9	-3.389219	0.0001105
ETAE_1449	zwf glucose-6-phosphate 1-dehydrogenase	38	758.55728	791.37315	786.86879	199.27	110.26	237.77	-2.087982	0.000108
ETAE_1450	devB 6-phosphoglucuronolactonase	14	340.19103	326.88965	331.29334	67.61	45.23	66.78	-2.455064	5.113E-06
ETAE_1461	minE cell division topological specificity factor	11	21.476707	40.861206	19.424365	0	2.2	0	-4.026839	0.018129
ETAE_1462	minD septum site-determining protein	24	420.94345	381.09329	343.16378	83.21	26.4	63.6	-2.704007	0.0003169
ETAE_1463	minC septum site-determining protein	25	749.9666	747.17634	679.85278	202.83	85.82	149.81	-2.311821	0.0030053
ETAE_1468	dsbB disulphide bond formation protein	12	188.13595	202.63823	207.19323	20.8	15.4	34.98	-3.018185	2.801E-05
ETAE_1472	fadR fatty acid metabolism regulator	36	45.530618	44.196815	41.006993	10.4	0	12.72	-2.356144	0.0009716
ETAE_1486	pncA nicotinamidase/pyrazinamidase	11	487.95078	478.656984	466.18476	41.61	72.61	143.1	-2.463492	0.0002167
ETAE_1518	cs cardiolipin synthetase	50	130.57838	147.60068	125.17924	23.4	24.2	25.44	-2.417922	8.423E-05
ETAE_1545	rlnB ribosomal large subunit pseudouridine synthase B	22	23.194843	30.020478	30.215679	7.8	0	3.18	-2.628184	0.001718
ETAE_1553	cysB transcriptional regulator	35	10.308819	15.010239	15.10784	3.2	0	3.18	-2.663687	0.0081815
ETAE_1554	cspQ cold shock protein	6	67.866393	73.38339	87.409643	10.4	0	9.54	-3.336068	0.0004868
ETAE_1557	manZ PTS system, mannose-specific IID component	31	658.90536	669.62344	659.34928	130.63	183.85	176.66	-2.010413	8.092E-06
ETAE_1558	manY PTS system, mannose-specific IIC component	25	724.19455	795.54266	663.66581	188.87	160.87	163.41	-2.082697	0.0001419
ETAE_1559	manX PTS system, mannose-specific IIB component	34	877.96777	831.40046	834.16857	179.88	244.48	199.27	-2.022851	1.249E-05
ETAE_1614	regulatory prophage protein cl	20	115.97422	107.57338	111.15053	15.6	46.21	12.72	-2.122908	0.001394
ETAE_1615	cra putative phage transcriptional regulatory protein	11	85.906827	94.239044	124.10011	18.2	11	25.44	-2.414214	0.0025085
ETAE_1632	hypothetical protein	3	60.993847	59.207053	46.40265	2.6	0	31.8	-2.181056	0.0169463
ETAE_1655	prc carboxy-terminal protease	61	174.39086	168.44824	176.97755	7.8	11	9.54	-4.060228	4.347E-07
ETAE_1675	hypothetical protein	5	32.645494	28.352673	36.690467	5.2	0	12.72	-2.266933	0.0038064
ETAE_1695	rseX SoxR-reducing system protein	17	91.920305	84.224118	90.647037	33.8	4.4	6.36	-2.504024	0.0016318
ETAE_1696	rnfG electron transport complex protein	16	34.362731	33.356086	14.028708	5.2	0	0	-3.369475	0.020289
ETAE_1697	rnfD electron transport complex protein	23	46.389668	40.861206	46.40265	7.8	2.2	0	-3.393939	0.0001554
ETAE_1698	rnfC electron transport complex protein	46	157.20949	151.77019	142.44534	44.21	33.01	15.9	-2.241134	0.0002112
ETAE_1700	Na(+) -translocating NADH-quinone reductase subunit E	20	29.208321	19.17975	21.582628	0	0	0	-4.604283	0.0015192
ETAE_1705	add adenosine deaminase	23	353.07706	321.88623	311.86897	28.6	11	54.06	-3.356193	7.179E-05
ETAE_1712	hypothetical protein	9	55.839437	41.695108	61.51049	5.2	18.61	6.36	-2.398249	0.0146758
ETAE_1786	predicted permease	34	693.26809	696.3083	656.11189	202.83	72.61	128.46	-2.340516	0.00217
ETAE_1792	aroH 3-deoxy-7-phosphoheptulonate synthase	25	81.611485	80.88851	90.647037	0	0	0	-6.415866	1.139E-05
ETAE_1822	pkfB 6-phosphofructokinase	26	125.24397	123.99716	123.88428	310.12	201.66	298.53	-2.199332	9.649E-06
ETAE_1839	hypothetical protein	4	125.42397	112.57679	108.99227	10.4	24.2	12.72	-2.798122	0.0001087
ETAE_1840	pabB anthranilate/para-aminobenzoate synthases component I	31	584.16642	615.4198	575.17703	18.2	13.2	12.72	-5.23758	1.245E-06
ETAE_1851	hypothetical protein	14	21.476707	18.345848	7.5539198	0	0	0	-4.069716	0.0200376
ETAE_1883	potC spermidine/putrescine ABC transporter membrane protein	22	159.7867	203.47213	187.76886	44.21	19.8	56.32	-2.195128	0.0027405
ETAE_1910	hypothetical protein	35	972.46528	1040.7099	966.90173	231.43	160.63	318	-2.064706	0.0001232
ETAE_1967	HpcH/HpaI aldolase family protein	9	121.98769	117.5802	90.647037	13	6.6	38.16	-2.455259	0.002714
ETAE_1981	hypothetical protein	3	16.322297	19.17975	14.028708	5.2	2.6	0	-2.666779	0.0073317
ETAE_1982	IeuD 3-isopropylmalate dehydratase small subunit	19	469.05127	483.66325	444.60213	86.27	62.65	52.45	-2.776498	1.246E-05
ETAE_2044	uvrC excinuclease ABC subunit C	51	923.49839	969.82821	1095.3184	289.32	185.54	252.25	-2.034753	0.0002255
ETAE_2071	transcriptional regulator, DeoR family	13	450.15177	456.14448	473.73686	60.05	62.63	54.25	-2.94237	7.32E-07
ETAE_2079	DedA-family membrane protein	12	232.8075	240.16382	254.67501	75.41	8.8	79.5	-2.131833	0.0013896
ETAE_2099	fabG 3-ketoacyl-(acyl-carrier-protein) reductase	28	12.886024	10.006826	21.582628	2.2	2.2	3.18	-2.55274	0.034601
ETAE_2114	tumA NADH-lyase, F-e-S type, tartrate/fumarate subfamily, beta subunit	59	927.39737	885.6041	91.86603	101.41	26.4	238.5	-2.885079	0.0002415
ETAE_2176	cmt cytidylate kinase	19	24.059312	27.518771	25.899153	10.4	0	6.36	-2.025901	0.0031556
ETAE_2177	aroA 3-phosphoshikimate 1-carboxyvinyltransferase	35	572.99854	569.55518	467.26389	2.6	2.2	3.18	-7.198561	0.0001039
ETAE_2178	serC phosphoserine aminotransferase	36	11.167887	9.172938	5.395657	2.2	0	0	-3.548239	0.047556
ETAE_2202	trxB thioiredoxin reductase (NADPH)	29	329.02315	317.17672	311.86897	26	8.8	57.24	-3.38843	4.337E-05
ETAE_2241	grxA glutaredoxin 1	11	10.119181	91.72938	76.618329	26	19.8	12.72	-2.197424	0.0022761
ETAE_2260	undecaprenyl-diphosphatase	17	250.84793	281.85893	283.81156	18.2	0	69.96	-3.168304	0.0004969
ETAE_2281	hisF imidazole glycerol phosphate synthase subunit	16	413.21184	437.79863	405.7534	75.22	96.82	120.84	-2.101332	3.442E-05
ETAE_2298	NLP/P60 protein	29	852.19572	833.90216	836.32683	91.01	123.22	111.3	-2.942429	3.066E-07
ETAE_2306	hypothetical protein	12	19.75857	23.349261	22.661759	5.8	6.6	0	-2.407081	0.0030077
ETAE_2324	putative transcriptional regulator	20	871.09522	778.03072	883.80861	137.82	11	152.64	-3.058193	0.0001852
ETAE_2326	transcriptional regulator, AraC family	13	271.46557	277.66842	317.63158	93.61	26.4	50.88	-2.307481	0.0005486
ETAE_2327	rheB ATP-dependent RNA helicase	36	149.0741	143.43778	141.15039	291.88	491.87	224.32	-2.118352	0.0001989
ETAE_2334	rcsB two-component system, NarL family, captular synthesis response regulator	14	140.02813	134.25825	105.75488	20.8	35.21	0	-2.698468	0.0018366
ETAE_2346	moaC molybdenum cofactor biosynthesis protein C	10	615.95195	585.39932	512.58741	86.46	102.06	97.85	-2.568851	0.0001052
ETAE_2347	moaA molybdenum cofactor biosynthesis protein A	22	382.28558	434.43603	382.01251	79.65	43.48	47.65	-2.798916	8.022E-05
ETAE_2359	menF isochorismate synthase	29	652.89188	676.29465	693.88149	190.28	155.25	120.11	-2.112131	2.476E-05
ETAE_2373	nuoN NADH:ubiquinone oxidoreductase subunit 2 (chain N)	38	109.10167	102.56997	85.25138	2.6	4.4	3.18	-4.50817	0.0001796
ETAE_2374	nuoM NADH:ubiquinone oxidoreductase subunit 4 (chain M)	45	170.95459	180.12287	183.45234	7.8	0	3.18	-5.264908	2.348E-06
ETAE_2375	nuoL NADH:ubiquinone oxidoreductase subunit 5 (chain L)/multisubunit Na+/H+	56	267.17023	260.17747	295.682	5.2	4.4	9.54	-5.221467	1.67E-05
ETAE_2376	nuoK NADH:ubiquinone oxidoreductase subunit 11 or 4L (chain K)	7	36.939936	50.03413	46.40265	0	0	3.18	-4.463847	0.0004274
ETAE_2377	nuoJ NADH:ubiquinone oxidoreductase subunit 6 (chain J)	12	90.202168	95.064846	83.093117	13	6.6	19.08	-2.702782	0.0001067
ETAE_2378	nuoI formate hydrogenlyase subunit 6/NADH:ubiquinone oxidoreductase 23 kD	18	42.094435	50.868032	36.690467	0	0	0	-5.466549	0.0004718
ETAE_2380	nuoG NADH dehydrogenase/NADH:ubiquinone oxidoreductase 75 kD subunit (c	68	349.64079	371.08646	397.12035</					

ETAE_2441	evpM type VI secretion system protein EvpM	38	1110.7753	1144.9477	1225.8933	10.4	39.61	25.44	-5.473083	5.483E-06
ETAE_2442	evpN type VI secretion system protein EvpN	20	970.74714	999.84869	1038.1244	20.8	50.61	34.98	-4.783035	1.412E-06
ETAE_2443	evpO type VI secretion system protein EvpO	86	2226.705	2208.1729	2314.7368	41.61	50.61	47.7	-5.562167	3.007E-07
ETAE_2445	aroC chorismate synthase	19	374.55377	376.08987	336.689	88.41	35.21	115.36	-2.185832	0.00263
ETAE_2448	sixA phosphohistidine phosphatase	10	53.262233	47.532423	53.95657	7.8	0	12.72	-2.745691	0.0004502
ETAE_2488	hypothetical protein	9	25.772048	19.17975	32.373942	5.2	2.2	9.54	-2.010197	0.0099396
ETAE_2524	hypothetical protein	12	21.476707	25.017065	26.978285	7.8	4.4	3.18	-2.056796	0.0007999
ETAE_2547	uvrB excinuclease ABC subunit B	43	1812.634	1826.2457	1890.6382	455.24	340.04	281.56	-2.357119	1.24E-05
ETAE_2575	tolR colicin uptake protein	13	3.7181365	1.6678043	4.3165256	0	0	0	-2.01	0.0425803
ETAE_2582	sucC succinyl-CoA synthetase subunit beta	27	422.66159	406.94425	374.45859	46.81	57.21	38.16	-3.055588	2.028E-05
ETAE_2583	sucB 2-oxoglutarate dehydrogenase, E2 subunit, dihydrolipoamide succinyltrans	28	173.53179	162.61092	185.6106	13	6.6	6.36	-4.179506	1.897E-05
ETAE_2584	sucA component of the 2-oxoglutarate dehydrogenase complex, thiamin-binding	59	824.70554	778.03072	796.39897	0	8.8	9.54	-6.814613	5.663E-07
ETAE_2585	sdhB succinate dehydrogenase iron-sulfur subunit	20	297.23762	288.53015	285.96982	78.81	63.62	64.94	-2.055921	3.082E-06
ETAE_2586	sdhA succinate dehydrogenase catalytic subunit	33	977.61969	1006.5199	1015.4626	218.43	288.25	157.58	-2.169892	3.887E-05
ETAE_2587	sdhI succinate dehydrogenase cytochrome b556 small membrane subunit	15	902.88075	927.2992	904.31211	179.42	74.81	168.54	-2.684709	2.281E-05
ETAE_2588	sdhC succinate dehydrogenase cytochrome b556 large membrane subunit	11	193.29036	180.12287	173.74015	57.61	33.01	34.98	-2.123109	0.0004052
ETAE_2589	gltA type II citrate synthase	41	1353.0325	1366.7656	1427.6908	176.82	244.24	337.08	-2.447048	2.617E-05
ETAE_2593	edwR AHL-dependent transcriptional regulator	32	275.76091	287.69625	247.12109	67.61	35.21	38.16	-2.498413	0.0001484
ETAE_2617	nagB glucosamine-6-phosphate deaminase	33	446.7155	467.81911	468.34303	140.42	68.21	114.48	-2.087369	9.26E-05
ETAE_2619	nagC putative N-acetylglucosamine regulatory protein	28	168.37738	185.96018	173.74015	41.61	8.8	12.72	-3.005546	0.0001802
ETAE_2649	lipE lipote-protein ligase B	23	329.88222	331.05916	240.6463	101.41	28.61	87.66	-2.050259	0.0057991
ETAE_2700	purE phosphoribosylaminoimidazole carboxylase catalytic subunit	8	164.94411	156.77381	149.99926	7.8	2.2	22.26	-3.750954	3.788E-05
ETAE_2701	purK phosphoribosylaminoimidazole carboxylase	22	752.54538	804.71569	735.96761	187.23	170.65	181.26	-2.088647	0.0001005
ETAE_2724	lepA GTP-binding protein	48	79.034281	70.881684	85.25138	26	6.6	6.36	-2.504889	0.0010659
ETAE_2731	srmA ATP-dependent RNA helicase	31	399.46674	342.73379	362.58815	46.81	17.6	47.7	-3.266598	6.846E-05
ETAE_2738	smpA SmpA/OmlA domain protein	14	79.893349	82.556314	66.906146	10.4	6.6	19.08	-2.571833	0.00045
ETAE_2739	hypothetical protein	7	18.899502	16.678043	39.927862	7.8	0	0	-2.861761	0.045251
ETAE_2786	guA GMP synthase, large subunit	31	108.2426	105.07167	135.97056	15.6	2.2	0	-4.082087	0.0005442
ETAE_2787	guAB inositol-5-monophosphate dehydrogenase	35	621.96543	642.10466	614.02576	0	0	0	-9.292395	1.903E-07
ETAE_2812	hscB co-chaperone Hsc20	11	20.617638	15.010239	20.503497	3.8	0	6.36	-2.465907	0.0098739
ETAE_2855	gshA glutamate-cysteine ligase	47	134.01465	123.41752	133.81229	2.6	8.8	25.44	-3.306801	0.0001034
ETAE_2876	surE stationary phase survival protein	14	315.27805	311.04551	305.39418	57.42	50.61	85.86	-2.264655	0.0005739
ETAE_2877	truD tRNA pseudouridine synthase D	18	500.8368	474.49033	433.81082	111.82	63.81	124.02	-2.222159	0.0001596
ETAE_2879	ispD 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase	13	18.040434	32.522184	49.640044	0	0	0	-5.104374	0.0216312
ETAE_2947	Z-ring-associated protein	13	348.78172	346.9033	339.92639	110.82	63.81	50.88	-2.199219	0.0012334
ETAE_2964	metK S-adenosylmethionine synthetase	30	14.604161	12.50852	8.6330512	4.8	0	0	-2.896665	0.0410327
ETAE_3036	hypothetical protein	12	28.349253	17.511945	14.028708	7.8	0	3.18	-2.169464	0.0286353
ETAE_3148	panF sodium/panthothenate symporter	37	285.21067	335.22867	328.05594	65.01	59.41	79.5	-2.201123	0.00012
ETAE_3248	trpD anthranilate synthase component II	17	702.71784	636.26735	623.73795	31.2	15.4	85.86	-3.859121	4.742E-05
ETAE_3278	envZ osmolarity sensor protein	44	786.90653	751.34585	841.72249	109.21	79.21	159	-2.765607	4.248E-05
ETAE_3312	glpD glycerol-3-phosphate dehydrogenase	41	42.094345	65.044369	31.29481	7.8	2.2	0	-3.443541	0.0138098
ETAE_3327	cell division protein	26	2007.6425	2027.2162	1773.0129	518.88	255.25	575.59	-1.999998	0.0004977
ETAE_3328	ftsE cell division protein	17	1525.7052	1475.1729	1310.0655	405.66	239.84	386.55	-2.062491	0.0004073
ETAE_3367	gor glutathione-disulfide reductase	42	809.24231	883.93629	793.16158	260.04	116.62	146.28	-2.242793	0.0002275
ETAE_3400	procC pyrrole-5-carboxylate reductase	16	210.47173	203.47213	199.63931	5.2	2.2	9.54	-4.950558	8.151E-07
ETAE_3438	menA 1,4-dihydroxy-2-naphthylidenepropenyltransferase	33	1108.1981	1129.1035	1148.1958	313.91	231.28	226.47	-2.129011	9.101E-06
ETAE_3446	fpr ferredoxin-NADP reductase	23	394.31234	361.07964	353.9551	41.61	22	66.78	-3.059885	5.398E-05
ETAE_3450	pflB 6-phosphofructokinase	35	240.53912	276.02162	279.49503	65.01	30.81	69.96	-2.243152	0.0002762
ETAE_3461	gpm phosphoglycerate mutase	45	173.53179	150.93629	152.15753	62.41	0	22.26	-2.451754	0.0026717
ETAE_3462	membrane-bound metallopeptidase	35	1879.6414	1886.2867	2006.1053	309.44	72.61	314.82	-3.04467	4.677E-05
ETAE_3476	transcriptional regulator, LysR family	29	1140.8427	1159.9579	1083.4479	72.81	2.2	216.24	-3.525	0.0001043
ETAE_3481	argB acetylglutamate kinase	19	1043.7679	1050.7167	1016.5418	117.47	359.38	244.47	-2.104082	0.0003523
ETAE_3492	yihK GTP-binding protein	64	111.67887	116.7463	98.200957	0	2.2	3.18	-5.297737	4.449E-05
ETAE_3493	glnA glutamine synthetase	44	944.9751	953.98407	1026.254	252.23	55.01	155.82	-2.651433	0.000194
ETAE_3531	altpA F-type H ⁺ -transporting ATPase alpha chain	50	5.1544096	8.3390216	5.395657	0	0	1.18	-4.000694	0.0237262
ETAE_3540	pstA ABC-type phosphate transport system, permease component	26	24.053912	19.17975	21.582628	5.2	0	0	-3.047936	0.0008805
ETAE_3554	lpnH UDP-2,3-diacylglycosamine pyrophosphatase	24	18.899502	17.511945	22.661759	0	2.2	0	-3.577387	0.0003716

Table S11 Regions for genes of interests after in vivo and in vitro screening

ID	loc-s	loc-e	gene	Annotation-nr	DMEM/Input	J774A.1/Input	Turbot/Input	category
ETAE_0073	65362	66456	wabH	putative glycosy	-2.131967136	0.753208244	1.042644337	region1
ETAE_0074	66460	67587	waaG	lipopolysacchari	-1.12998707	1.177650735	-3.191256379	region1
ETAE_0075	67584	68660	waaQ	heptosyl III transf	-2.856865972	0.586506554	-2.036103053	region1
ETAE_0076	68793	69764	walW	lipopolysaccahr	-3.033742991	-0.904675783	-2.703135853	region1
ETAE_0077	69850	71061		putative glycosy	-0.012558773	-0.235049314	0.780377638	region1
ETAE_0078	71268	72374	walR	putative glycosy	-2.503221036	0.126252918	1.592898884	region1
ETAE_0079	72378	73535	wabK	putative glycosy	-3.104561798	0.255657819	0	region1
ETAE_0080	73537	74670	waal	lipid A core - O-c	0.11381022	-0.867720963	0.1398864	region1
ETAE_0081	74660	75625	waaC	ADP-heptose:LPs	1.097024454	4.07980996	-1.315193724	region1
ETAE_0082	75613	76674	waaF	ADP-heptose:LPs	0.952243775	0.383626693	-1.191401195	region1
ETAE_0083	76698	77627	rfaD	ADP-L-glycero-D	-1.659230386	0.639753601	-1.355002573	region1
ETAE_00860	942694	942915	esaG	putative type III s	0.335681578	-2.735750711	-4.451342723	region2
ETAE_00861	942934	943626	esrC	putative transcript	0.684854525	-3.500291293	-5.386702891	region2
ETAE_00862	943711	943935		hypothetical pro	0.40428455	0.289725369	0.764671248	region2
ETAE_00863	943946	945136	esAD	type-III secretion	0.030868954	0.183084454	-6.387494218	region2
ETAE_00864	945144	946628	esAC	Type II secretory	0.081003617	-0.488940103	-6.358695937	region2
ETAE_00865	946625	947104	esAB	two-component	-0.074744379	-0.89731476	-3.761403766	region2
ETAE_00866	947143	948162	eseG	two-component	-0.132075936	0.233248623	-1.015808013	region2
ETAE_00867	948050	948541	escB	type III secretion	0.210345642	0.08486143	-1.413532895	region2
ETAE_00868	948697	949080	eseE	type III secretion	0.102403366	-0.139261225	-5.287189162	region2
ETAE_00869	949091	949672	esED	type III secretion	0.121914919	0.030338547	-2.689366966	region2
ETAE_00870	949688	951208	eseC	type III secretion	0.057012458	-0.015564898	-2.90474386	region2
ETAE_00871	951121	951588	esCA	type III secretion	0.000339695	0.099214783	-3.190974411	region2
ETAE_00872	951599	952195	eseB	EspA family secre	0.303516655	0.083900032	-2.003910926	region2
ETAE_00873	952220	952567		conserved hypo	-0.04188396	-0.448116749	-3.48914182	region2
ETAE_00874	952871	953785	esaQ	translocation prc	0.125934069	-0.383922713	-5.315107022	region2
ETAE_00875	953763	954236	esap	putative major fc	-0.073484791	0.072366716	-4.792360448	region2
ETAE_00876	954233	954601	esaO	type III secretion	0.007363707	-1.358719487	-6.542351654	region2
ETAE_00877	954595	955911	esAN	type III secretion	-0.061162966	0.109063679	-6.287144834	region2
ETAE_00878	955898	957955	esAV	type III secretion	-0.1003326418	-0.140884208	-5.765091606	region2
ETAE_00879	957939	958319	esAM	type III secretion	-0.016531647	0.653106873	-5.101363541	region2
ETAE_00880	958499	959146	esar	type three secre	-0.160763926	1.154581364	-7.27607721	region2
ETAE_00881	959165	959434	esas	type III secretion	-0.039179404	-2.101048748	-6.492993654	region2
ETAE_00882	959434	960216	esAT	type III secretion	0.087116512	-2.967488623	-7.111083855	region2
ETAE_00883	960213	961271	esAU	type III secretion	0.050794596	-3.749764336	-5.299346927	region2
ETAE_00884	961264	961914		putative transgly	0.172018165	-3.553164088	-3.487438937	region2
ETAE_00885	961860	964661	esrA	two-component	0.344473177	-3.261240893	-6.427381621	region2
ETAE_00886	964658	965302	esrB	two-component	0.0993314503	-3.141890979	-1.667898125	region2
ETAE_2373	2498661	2500118	nuoN	NADH:ubiquinon	0.029350574	-1.694218146	-4.508169964	region3
ETAE_2374	2500125	2501648	nuoM	NADH:ubiquinon	-0.061208448	-2.495901617	-5.264908444	region3
ETAE_2375	2501726	2503570	nuoL	NADH:ubiquinon	-0.050097379	-1.71155172	-5.221466542	region3
ETAE_2376	2503567	2503869	nuoK	NADH:ubiquinon	-0.350862071	-1.924347325	-4.463846693	region3
ETAE_2377	2503866	2504438	nuoJ	NADH:ubiquinon	-0.430819243	-3.582410392	-2.702781594	region3
ETAE_2378	2504449	2504991	nuoI	formate hydroge	0.093756537	-2.221482716	-5.4665493	region3
ETAE_2379	2505009	2505986	nuoH	NADH:ubiquinon	0.171767748	-0.594618037	-3.324725804	region3
ETAE_2380	2505983	2508718	nuoG	NADH dehydrog	0.044950791	-1.717393614	-5.063424172	region3
ETAE_2381	2508779	2510125	nuoF	NADH:ubiquinon	0.114994175	-3.648998663	-5.04518692	region3
ETAE_2382	2510122	2510622	nuoE	NADH:ubiquinon	0.399321671	-1.204341271	-5.066651638	region3
ETAE_2383	2510625	2512421	nuoD	NADH:ubiquinon	0.110122605	-2.274364101	-4.757854957	region3
ETAE_2384	2512521	2513195	nuoB	NADH:ubiquinon	-0.117939029	-1.810318535	-6.109447314	region3
ETAE_2385	2513259	2513618	nuoA	NADH:ubiquinon	0.445872139	-2.500223919	-4.422818233	region3
ETAE_2428	2554573	2555136	evpP	type VI secretion	-0.011017199	-0.192785759	-2.417594876	region4
ETAE_2429	2555386	2555901	evpA	type VI secretion	0.194115069	-0.699248533	-7.934881853	region4
ETAE_2430	2555898	2557385	evpB	type VI secretion	-0.052361621	-0.402694053	-3.065273183	region4
ETAE_2431	2557455	2557946	evpC	type VI secretion	-0.044305356	-0.429432845	-5.276353601	region4
ETAE_2432	2558043	2559248	evpD	type VI secretion	0.039331342	-0.079935946	-4.951615624	region4
ETAE_2433	2559257	2559733	evpE	type VI secretion	0.001068584	-0.160079486	-5.620110996	region4
ETAE_2434	2559737	2561578	evpF	type VI secretion	-0.092801513	-0.021723099	-6.092925209	region4
ETAE_2435	2561575	2562600	evpG	type VI secretion	-0.055207938	-0.417009205	-4.85852982	region4
ETAE_2436	2562719	2565331	evpH	type VI secretion	0.0317927	-0.165080774	-6.114718663	region4
ETAE_2437	2565331	2567316	evpI	type VI secretion	0.051936205	-0.262586565	-5.739158632	region4
ETAE_2438	2567411	2567713	evpJ	type VI secretion	0.123923695	0.138112081	-6.904822177	region4
ETAE_2439	2567727	2568794	evpK	type VI secretion	0.027725446	-0.232445393	-5.83709582	region4
ETAE_2440	2568715	2569422	evpL	type VI secretion	0.299428151	0.058126482	-5.887241041	region4
ETAE_2441	2569522	2570910	evpM	type VI secretion	0.211728325	-0.087730055	-5.473082509	region4
ETAE_2442	2570907	2571557	evpN	type VI secretion	-0.531149416	-0.106168872	-4.783034998	region4
ETAE_2443	2571545	2575336	evpO	type VI secretion	0.030628451	-0.184312318	-5.56216652	region4

Table 12 Intracellular growth deficient mutants

Strain with mutation in	Annotation	Gene name	Intracellular growth	Autoaggregation
WT	/		+	+
ETAE_0854	Putative type III secretion apparatus	esaL ETAE_0854	-	-
ETAE_0856	Uncharacterized protein	ETAE_0856	-	-
ETAE_0857	Lipoprotein	esaJ ETAE_0857	-	-
ETAE_0860	Putative type III secretion system needle protein	esaG ETAE_0860	-	-
ETAE_0861	Putative transcriptional regulator EsrC	esrC ETAE_0861	-	-
ETAE_0863	Type-III secretion protein	esaD ETAE_0863	-	-
ETAE_0864	Type II secretory pathway, component PulD like protein	esaC ETAE_0864	-	-
ETAE_0865	Uncharacterized protein	esaB ETAE_0865	-	-
ETAE_0872	EspA family secreted protein	esaB ETAE_0872	-	-
ETAE_0875	Putative major facilitator family transporter	esaP ETAE_0875	-	-
ETAE_0877	Type III secretion system ATPase	esaN ETAE_0877	-	-
ETAE_0878	Type III secretion protein, HrcV family	esaV ETAE_0878	-	-
ETAE_0879	Type III secretion apparatus protein	esaM ETAE_0879	-	-
ETAE_0880	Type three secretion apparatus protein R	esaR ETAE_0880	-	-
ETAE_0881	Type III secretion apparatus	esaS ETAE_0881	-	-
ETAE_0882	Type III secretion apparatus protein	esaT ETAE_0882	-	-
ETAE_0883	Type III secretion apparatus protein	esaU ETAE_0883	-	-
ETAE_0884	Putative transglycosylase signal peptide protein	ETAE_0884	-	-
ETAE_0885	Two-component sensor/regulator EsrA	esrA ETAE_0885	-	-
ETAE_0886	Two-component response regulator EsrB, LuxR family	esrB ETAE_0886	-	-
ETAE_0861-62	/	/	-	-
ETAE_0863-64	/	/	-	-
ETAE_0873-74	/	/	-	-
ETAE_0879-80	/	/	-	-
ETAE_2071	Transcriptional regulator, DeoR family	ETAE_2071	-	-
ETAE_0323	Putative invasin	ETAE_0323	-	-
ETAE_1437	Probable transcriptional regulatory protein ETAE_1437	ETAE_1437	-	-
ETAE_1039	Copper transporter	ybaR ETAE_1039	-	-
ETAE_1720	Putative porin	ETAE_1720	-	-
ETAE_2185	Uncharacterized protein	ETAE_2185	-	-
ETAE_3320	Glycerol-3-phosphate transporter permease	ugpA ETAE_3320	-	-
ETAE_1709	DNA replication terminus site-binding protein	tus ETAE_1709	-	-
ETAE_3474	HTH-type transcriptional repressor FabR	fabR ETAE_3474	-	-
ETAE_3493	Glutamine synthetase	glnA ETAE_3493	-	-

Table S14 Bacterial strains and plasmids used in this study

Strain or plasmid	Genotype
<i>Escherichia coli</i>	
SM10 λpir	<i>thi thr leu tonA lacy supE recA::RP4-2-Tc::Mu, pirR6K, Kan^r</i>
DH5a λpir	λpir lysogen D(ara-leu) araD D(lacX74) phoA20 <i>thi-1 rpoB argE (am) recA1</i>
BL21(DE3)	Host for protein expression
<i>Plasmid</i>	
pMKGR	Mariner Himar1 transposon encoding a gentamycin resistance gene. Suicide plasmid, <i>pir</i> dependent, R6K, Gm ^r
pDM4	Suicide plasmid, <i>pir</i> dependent, R6K, SacBR, Cm ^r
pDMK	pDM4 derivative with Kan resistance gene inserted in SalI site, Kan ^r , Cm ^r
pUTt	Complementation vector, Am ^r
pET28b-HisSumo	Protein expression vector, Km ^r
pET28b-HisSumo- <i>evrA</i>	Protein expression vector, Km ^r
pET28b-HisSumo- <i>evrA</i> ^{R221A}	Protein expression vector, Km ^r
pET28b-HisSumo- <i>evrA</i> ^{R7A}	Protein expression vector, Km ^r
pET28b-HisSumo- <i>evrA</i> ^{R178A}	Protein expression vector, Km ^r
pUTt- <i>evrA</i>	Complementation vector, ETAE_2071, Amp ^r
pUTt-P _{<i>eseB</i>} -EGFP	Complementation vector, <i>eseB</i> promoter driven EGFP expression, Amp ^r
pUTt-P _{<i>eseB</i>} - <i>luc</i>	Complementation vector, <i>eseB</i> promoter driven luciferase expression, Amp ^r
pUTt-P _{<i>esrB</i>} - <i>luxAB</i>	Complementation vector, <i>esrB</i> promoter driven luxAB expression, Amp ^r
pUTt-P _{<i>esrC</i>} - <i>luxAB</i>	Complementation vector, <i>esrC</i> promoter driven luxAB expression, Amp ^r
<i>Edwardsiella piscicida</i>	
EIB202	Wild type strain, CCTCC M 208068, Col ^r , Cm ^r , Kan ^s
wt ΔP	EIB202, pEIB202 cured, Col ^r , Cm ^s , Kan ^s
EIB202::Tn	EIB202, transposon insertion mutant, labelling mCherry, locus between ETAE_3351 and ETAE_3352
esrB::Tn	EIB202, transposon insertion mutant
<i>eseB</i> ::Tn	EIB202, transposon insertion mutant
2071::Tn	EIB202, transposon insertion mutant
pdhR::Tn	EIB202, transposon insertion mutant
2342::Tn	EIB202, transposon insertion mutant
esaB::Tn	EIB202, transposon insertion mutant
esaM::Tn	EIB202, transposon insertion mutant
ΔwaaQ	EIB202, in-frame deletion of ETAE_0075
ΔwalW	EIB202, in-frame deletion of ETAE_0076
ΔwabK	EIB202, in-frame deletion of ETAE_0079
ΔesrA	EIB202, in-frame deletion of <i>esrA</i>
ΔesrB	EIB202, in-frame deletion of <i>esrB</i>
ΔesrC	EIB202, in-frame deletion of ETAE_0861
ΔeseB	EIB202, in-frame deletion of ETAE_0872
ΔesaM	EIB202, in-frame deletion of ETAE_0879
ΔnuoM	EIB202, in-frame deletion of ETAE_2374
ΔnuoJ	EIB202, in-frame deletion of ETAE_2377
ΔnuoI	EIB202, in-frame deletion of ETAE_2378
ΔevpP	EIB202, in-frame deletion of ETAE_2428
ΔevpC	EIB202, in-frame deletion of ETAE_2431
ΔevpI	EIB202, in-frame deletion of ETAE_2437
ΔmanX	EIB202, in-frame deletion of ETAE_1559
ΔevrA	EIB202, in-frame deletion of ETAE_2071
ΔevrA+pUTt	EIB202, in-frame deletion of ETAE_2071 complemented with pUTt vector
ΔevrA+pUTt-2071	EIB202, in-frame deletion of ETAE_2071 complementation
ΔevrAΔesrB	EIB202, double in-frame deletion of ETAE_2071 and <i>esrB</i>
ΔevrAΔesrB+pUTt-P _{<i>esrB</i>} - <i>esrB</i>	EIB202, double in-frame deletion of ETAE_2071 and <i>esrB</i> complemented with <i>esrB</i> driven by its own promoter
ΔevrAΔesrB+pUTt-P _{<i>lac</i>} - <i>esrB</i>	EIB202, double in-frame deletion of ETAE_2071 and <i>esrB</i> complemented with <i>esrB</i> driven by <i>lac</i> promoter
ΔevrA+P _{<i>eseB</i>} - <i>luc</i>	EIB202, in-frame deletion of ETAE_2071 complemented with luciferase driven by <i>eseB</i> promoter
WT+P _{<i>eseB</i>} - <i>luc</i>	EIB202, wild type complemented with luciferase driven by <i>eseB</i> promoter

Table S15 Primers used in this study

primer name	primer sequence
transposon mutant library construction and validation	
evpP-F	TCATCGCACATAACAGAATAAACGCC
evpP-R	CCGTAACATTCTTACAACACTGCG
pG1	AGGTGATGCTACATACGGAAAG
pmh1	AGCGCATGAACTCCTTGATG
PpMar1	AAAAGTCCGCTGGCAAAG
PpMar2	CCCTTCAAGAGCGATACAAC
Sp1	GCTCCGTAGTAAGACATTATCGCG
Sp2	GCTTACGTTCTGCCAAGTTGAG
ABS	GGCCACGCGTCGACTAGTAC
AB2	GGCCACGCGTCGACTAGTACNNNNNNNNNCTGG
Seq2	CAATTGTTCAAGCCGAGATCG
mutants construction	
pDMK-2071-P1	cccccccgagctcaggtaaccggatctatGGCCAGATAGCTCAGCGCATCG
pDMK-2071-P2	CGCAGGATCACATAGTGCCTGTGCGGGCAG
pDMK-2071-P3	GGACACTATGTGATCCTGCGCAGAGAAAAATC
pDMK-2071-P4	gagtacgcgtactagtggggcccttagCGGGATGCGTCTGGATAACT
pDMK-2071-in-F	AGCTACCTCAAGCGGGTGCA
pDMK-2071-in-R	GAACTTGGAGGCCTCGGTCA
pDMK-2071-out-F	GCTCGGCAACCAGCAGCGTAT
pDMK-2071-out-R	GTGCTAATCCCACCGTCCC
pDM4-esrA-P1	cccccccgagctcaggtaaccggatctatGGTGCTCCGCTTAAATGG
pDM4-esrA-P2	AGTTTAATCCATAGGGGATTCTTTATG
pDM4-esrA-P3	ATCCCCTATGGATTAAAATCCAGAACCCC
pDM4-esrA-P4	gagtacgcgtactagtggggcccttagCGCGTGTGACGTGATCCGTC
pDM4-esrA-in-F	GCCGAAACGGTCTATGAGC
pDM4-esrA-in-R	CCTCGTAAAATTACTCTCC
pDM4-esrA-out-F	TGGAGAATATTCCGCTGGC
pDM4-esrA-out-R	GGGCTGGCCGTTATGAGG
pDM4-esrB-P1	gtggaattcccgaggagactCACCCGAGAAAAACCAACG
pDM4-esrB-P2	GGCCTCCTTACATAGTGCCTCCTGAG
pDM4-esrB-P3	GAGCACTATGTAAGGAGGCCACGATGACCG
pDM4-esrB-P4	aagcttatcgataccgtcgaGAAGTGACGCAGTATTTCCC
pDM4-esrB-in-F	CGATAGCATCATGTCCGAC
pDM4-esrB-in-R	CTGGGTGACAAAGTCGGAGC
pDM4-esrB-out-F	CGGCTGGACGATGGCTGG
pDM4-esrB-out-R	GGTGTGCTGCTGCGACTGC
pDM4-esaM-P1	gtggaattcccgaggagactAGCTTTGTCGCCACCTG
pDM4-esaM-P2	TAGCCAGCTACATGAATATCCTCCCGCGATC
pDM4-esaM-P3	GATATTCATGTAGCTGGCTACACAAACTC
pDM4-esaM-P4	aagcttatcgataccgtcgaCTGCACGACGTAATGATGG
pDM4-esaM-in-F	GCAAACCGAACCTTGGCTAC
pDM4-esaM-in-R	CATGGGGATTCTCCATCACG
pDM4-esaM-out-F	CGACACCACATCATCATCCCC
pDM4-esaM-out-R	AGAGCTGGCTCTTTTGC
pDM4-waaQ-P1	gtggaattcccgaggagactATAGCCGCTGTAGTCACTAC
pDM4-waaQ-P2	AATTGGGTACATTCTTTGCCTTGACTG
pDM4-waaQ-P3	CAAAAGAATGTGACCCAATTCAAGATTGGC
pDM4-waaQ-P4	aagcttatcgataccgtcgaTGGCATTGTAGATGACGTGG
pDM4-waaQ-in-F	GGATATGTTGTCGGCTAATC
pDM4-waaQ-in-R	CGGAGGTGAGCAAAATAGGG
pDM4-waaQ-out-F	CAGGTGATTAAACAGCGGGAG
pDM4-waaQ-out-R	GACATAGATCAGGCAGGAGG
pDM4-walW-P1	gtggaattcccgaggagactGCTATATCCCATGGTGACGC
pDM4-walW-P2	AACGAAATTACATACTGATACGTCTTCTTC
pDM4-walW-P3	TATCAGTATGTAATTCTGTTGGCGCATG
pDM4-walW-P4	aagcttatcgataccgtcgaCTAGAGGCTGGAACCTATGG

pDM4-walW-in-F CCGCTATTGCCTCGATTCC
pDM4-walW-in-R GCGTCCACAAAATAGGCG
pDM4-walW-out-F GAGAGCATTAAATGACCTCGC
pDM4-walW-out-R CCGCATCATTGCTATATCG
pDM4-wabK-P1 gtggaaattcccgaggagactGCATGTTTATTAGCTGGC
pDM4-wabK-P2 TCATTATCTACATGTTACTTGCTCCGAT
pDM4-wabK-P3 GTATAACATGTAGATAATGAATATGCCAC
pDM4-wabK-P4 aagcttatcgataccgtcgaGGGAGCAACCAGCGTTTATC
pDM4-wabK-in-F GTTGAGCGTCTTATTAGAC
pDM4-wabK-in-R GCAATCAGGGGAAGTAATCC
pDM4-wabK-out-F CGATAGATTCTATCCAATCC
pDM4-wabK-out-R TCCACATAATAGTTTGCCC
pDM4-nuoM-P1 gtggaaattcccgaggagactCCGAATGATCTTCATCGTGT
pDM4-nuoM-P2 TGGCGATTACATGGCGTTGGTTCCCTT
pDM4-nuoM-P3 AAACGCCATGTAAATGCCATGACAATAAC
pDM4-nuoM-P4 aagcttatcgataccgtcgaCTCTATCCCCAGGAACAGCG
pDM4-nuoM-in-F GCTGAAAGCGCCGCGCTGG
pDM4-nuoM-in-R CGATCAGGAAGTACATCGGC
pDM4-nuoM-out-F GCAGAACATCTCAAGATG
pDM4-nuoM-out-R CGCGAAGGAGAGATCGC
pDM4-nuoJ-P1 gtggaaattcccgaggagactGCTGGCGTATTGCCCTGCC
pDM4-nuoJ-P2 AACGGGATCACATGCTCGGCTCCTTAGGG
pDM4-nuoJ-P3 GCCGAGCATGTGATCCGTTACAACATGGG
pDM4-nuoJ-P4 aagcttatcgataccgtcgaCGGGAAAAGGCCAGGATCAG
pDM4-nuoJ-in-F CCGTGTGGCGACGATCCG
pDM4-nuoJ-in-R CAGCATCGAGACCAGCTCC
pDM4-nuoJ-out-F CACCTGTTCTCGGCGGC
pDM4-nuoJ-out-R CGGTAAAGTTACCGACCAGC
pDM4-nuol-P1 gtggaaattcccgaggagactCTGTTCTCCTGATGATGGC
pDM4-nuol-P2 CGGCTCCTTATGTCATGGTTACACTCACC
pDM4-nuol-P3 AACCATGACATAAGGAGCCGAGCATGGAAT
pDM4-nuol-P4 aagcttatcgataccgtcgaCAATGTGGAAGGCCACGAC
pDM4-nuol-in-F GTTGGTTCCGGCACCCAAAG
pDM4-nuol-in-R CCATACGGTAAAAGTTGTAG
pDM4-nuol-out-F GATCAAGATGTTCTCAAGG
pDM4-nuol-out-R CCTTATTGCTCAGTACCTC
pDM4-evpC-P1 gtggaaattcccgaggagactGCACACCTTACCACCGATG
pDM4-evpC-P2 CGTCTTACTTCATAGCGGACCTCTCTTG
pDM4-evpC-P3 GTCCGCTATGAAGTAAGACGGTCAAACAGG
pDM4-evpC-P4 aagcttatcgataccgtcgaCCAGCACATTGGACGCCGTC
pDM4-evpC-in-F CGATAAGCACA AAAAATGG
pDM4-evpC-in-R CATCGTAAATTGTAGATG
pDM4-evpC-out-F CAGGCCTATGCCAAATATGG
pDM4-evpC-out-R ATTGGTCAGCGCGATATAGG
pDM4-evpl-P1 gtggaaattcccgaggagactGTTCTCCAGGTGTTGAC
pDM4-evpl-P2 AAAATCAGTTCAAGGATGCCTTACAGGTG
pDM4-evpl-P3 GGCATCCTGAAGTGTGTTCGCTATCCGC
pDM4-evpl-P4 aagcttatcgataccgtcgaGTCATACAGCGGGTCGAAGC
pDM4-evpl-in-F GGCATTTCTACTGGTCG
pDM4-evpl-in-R GAGACAGATAAAAGTGGTC
pDM4-evpl-out-F CAACCTGCGGAGTTTCAG
pDM4-evpl-out-R CAGCTGCAGATCTTGCTGC
pDM4-manX-P1 gtggaaattcccgaggagactCATGGGTTATTCTGCCCTCG
pDM4-manX-P2 CGGCATATCAAATACTCACTCGCTACCTCC
pDM4-manX-P3 AGTGAGTATTGATATGCCGTAAGGCATTG
pDM4-manX-P4 aagcttatcgataccgtcgaGGTACGTACGATGATGGTC
pDM4-manX-in-F GTAACGTTGGCTTATCGAC
pDM4-manX-in-R GGTTCCCTGGTCCAGCG

pDM4-manX-out-F	ATCAGCGCTAAACTGAGC
pDM4-manX-out-R	GTGACGACCTCCGGAAATGG
pUTt-2071-F	ctcatccgccaaaacagccaGCAGCGGCTTCCACAGGTGG
pUTt-2071-R	ttaaaaattaaggaggaattTCAGCGGATGCCGTTTCAGAG
pUTt-P _{eseB} -luc-1	ctcatccgccaaaacagccaGTTAGTCCTCGGCTGGTCTGG
pUTt-P _{eseB} -luc-2	cgtctccatAGTGAACACCTCCTATTGACTAC
pUTt-P _{eseB} -luc-3	aggttcactATGGAAGACGCCAAAAACATAA
pUTt-P _{eseB} -luc-4	ttaaaaattaaggaggaattTTACACGGCGATCTTCCGCC
pUTt-P _{eseB} -EGFP-1	ctcatccgccaaaacagccaGTTAGTCCTCGGCTGGTCTGG
pUTt-P _{eseB} -EGFP-2	tgctagccatAGTGAACACCTCCTATTGACTAC
pUTt-P _{eseB} -EGFP-3	aggttcactATGGCTAGCAAAGGAGAAG
pUTt-P _{eseB} -EGFP-4	ttaaaaattaaggaggaattTTATTGTACAGTTCATCCATG
pET28b-Sumo-2071-F	tcacagagaacagattgggtgatccATGAACGCCAGACAACACG
pET28a-Sumo-2071-R	agtgggtgggtgggtcgatTCAGCGGATGCCGTTTCAG
pUTt-luxAB-F	TGGCTGTTGGCGGATGAG
pUTt-luxAB-R	CAAATAAGGAAATGTTATGAAAT
pUTt-P _{esrB} -luxAB-F	ctcatccgccaaaacagccaGTTATCGGCATATAAAATATT
pUTt-P _{esrB} -luxAB-R	atttcataacattccattttgATTAAAGGGTACTCCGAATC
pUTt-P _{esrC} -luxAB-F	ctcatccgccaaaacagccaGCCAACGCCCTGATCGACTGC
pUTt-P _{esrC} -luxAB-F	atttcataacattccattttgAGGTGCTCCTGACTGAGGTAC
qRT-PCR	
qPCR-esrA-F	TAGCGCCGTAGAGAAAACCC
qPCR-esrA-R	TCGCGGCAGATGGAGAATAC
qPCR-esrB-F	CGACCAGCTTGAGAATTGCG
qPCR-esrB-R	GTAGCCTCGTCCGATATGGC
qPCR-esrC-F	CCATGCCGAACTTGTCGTTG
qPCR-esrC-R	GAGTGTCAACGGACCTCCAC
qPCR-eseB-F	CCCGCTTCTTGAACTTGGC
qPCR-eseB-R	ACGCTATTCAACCGATCTGGC
qPCR-eseE-F	ATAACGGCCTGTCATCGTC
qPCR-eseE-R	TTTCTCATGGGACAGCGCAT
qPCR-evpP-F	GAATGGGGACGACTCACCTC
qPCR-evpP-R	AAATCCACCGAACCGAGGCAT
qPCR-evpC-F	CGATAAGCACA AAAATGG
qPCR-evpC-R	CTCCATCGTGCATTATTGC
qPCR-pdhR-F	ATCTGTTGGAAACCCGCCAT
qPCR-pdhR-R	TCGCGCAGACGAACAAAATC
qPCR-2342-F	CCATCCCAGACAGGACGAAC
qPCR-2342-R	AGGTGTATGGCTGGATGTGC
qPCR-2071-F	AAAACGCAGCTACCTCAAGC
qPCR-2071-R	AAGATGGTCTCCCCGTCGTT
qPCR-esaM-F	CTGAAATCCACAGCGCATCG
qPCR-esaM-R	GGGATCGCGACCGTATCTT
RT-gyrB-F	CCGATGATGGTACGGGTCTG
RT-gyrB-R	GCTTTTCAGACAGGGCGTTC
EMSA	
EMSA-P _{esrB} -F	tgcctgcaggtcgacgatCATGATGATCCATACTCCAAAG
EMSA-P _{esrB} -R	AAACTTCATATAGCTCGCTCG
EMSA-P _{manX} -F	tgcctgcaggtcgacgatCGGCTACATTGTTACGTC
EMSA-P _{manX} -R	ACGCCGTAAAATAAACACGG
Cy5-F	tgcctgcaggtcgacgat
FAM-F	tgcctgcaggtcgacgat

133 **Transparent Methods**

134 **Bacterial strains and culture conditions**

135 A list of the strains used in this study is in [Table S14](#). *E. piscicida* EIB202 cured of the
136 endogenous plasmid pEIB202, (*E. piscicida* EIB202 ΔP), was used as the parental strain for
137 generation of the transposon insertion mutant library. *E. coli* SM10λpir was used as the donor
138 for conjugations. *E. piscicida* strains were grown in Luria-Bertani broth (LB) or on LB agar
139 (LBA) (Oxoid, England) and in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen, USA)
140 at 28°C. DMEM was used to induce T3/T6SS production as well as to yield auto-aggregation
141 phenotypes mimicking in macrophages and *in vivo* conditions ([Zheng and Leung 2007](#)).
142 Where required, antibiotics were supplemented at the following concentrations: polymyxin B
143 (Col, 20 µg/mL), ampicillin (Amp, 100 µg/mL), kanamycin (Km, 25 µg/mL), streptomycin (Str,
144 100 µg/mL) and gentamicin (Gm, 15 µg/mL).

145 **Generation of the MKGR transposon**

146 The transposon derivatives were constructed within pMar2xT7, a vector containing the
147 *mariner* transposase outside the *Himar1* transposon ([Jacobs et al., 2003](#)). First, an *mcherry*
148 gene with a constitutive promoter, P_{tetA}, was inserted into the *Dra*III site in pMar2xT7 (yielding
149 PMmch). Promoterless *km* and *egfp* with their respective ribosome binding sequences (RBS)
150 were amplified from the appropriate templates ([Dennis and Zylstra 1998; Gu et al., 2016](#)) and
151 fused together with overlap PCR ([Table S15](#)) and then the triplet terminator TGACTAGCTAA
152 and a 48-bp T7 terminator were introduced at the 5' and at 3' ends, respectively. After
153 sequence verification, the complete amplicon was cloned into the *Nhe*I site of pMmch, yielding
154 pMKGR.

155 **Preparation of the transposon mutant library**

156 Transposon insertion mutants were generated by conjugative transfer of pMKGR into *E.*
157 *piscicida* EIB202 (Δp) from *E. coli* SM10λpir on LBA plates containing Gm and Col ([Yang et al.,](#)
158 [2017](#)). The mutants were then manually picked into 96-well plates, containing 200 µL LB per
159 well. After incubation at 28°C for 16-20 h, 20% glycerol was added to the wells and the plates
160 were stored at -80°C. Transposon insertion sites for each mutant were determined by thermal
161 asymmetric interlaced PCR (TAIL-PCR) ([Liu and Chen 2007](#)) as follows: the first round of PCR
162 was performed with a transposon-specific primer SP1 and a degenerate primer AB2 ([Table](#)
163 [S15](#)). The PCR products were then diluted 12-fold and used as the template for a second
164 round of PCR, using a nested transposon-specific primer SP2 and a primer ABS that
165 hybridizes to the defined portion of AB2. The third nested transposon-specific primer Seq2

166 was used for sequencing and the sequencing results were batch mapped by local BLAST
167 (<https://blast.ncbi.nlm.nih.gov>).

168 Five subset libraries were assembled from the original set of 20,346 unique insertion
169 mutants to facilitate targeted studies with smaller libraries ([Tables S1-S7](#)). The 1st (2,759
170 mutants) and 2nd (2,235 mutants) subset libraries each contained distinct insertion mutants
171 for each disrupted gene, with insertion sites located within the central 40-60% of each coding
172 sequence. Mutants in genes that only contained a single insertion were preferentially selected
173 for inclusion in the first subset library. The 3rd subset library (3,705 mutants) contained
174 transcriptional fusions and priority was given to mutants that had insertions close to beginnings
175 of ORFs in the creation of this library. The 4th library (2,305 mutants) consisted of insertions
176 within intergenic regions. The 5th library was a composite composed of equally mixture of
177 mutants from the 1st, 2nd and 4th subset libraries and contained 7,299 distinct insertion
178 mutants ([Table S1](#)).

179 **Construction of deletion mutants and complemented strains**

180 In-frame deletion mutants were generated using *sacB*-based allelic exchange ([Yin et al., 2018](#)). Overlap PCR was used to generate appropriate DNA fragments for creating in-frame
181 deletions in each target gene. The fragments were inserted into the suicide vector pDM4 with
182 Gibson assembly ([Gu et al., 2016](#)) and the resulting plasmids were introduced into *E. coli*
183 SM10 λpir for conjugation into *E. piscicida* EIB202. Transconjugants were selected on LBA
184 containing Km and Col. Double-crossover events were subsequently selected on LBA
185 containing 12% sucrose. To complement the *evrA* deletion in Δ*evrA*, an intact *evrA* containing
186 the putative promoter region was amplified and introduced into plasmid pUTt ([Yin et al., 2018](#));
187 the sequence of the insert was subsequently verified.

189 **Growth conditions for TIS studies**

190 To prepare input inocula for the TIS studies, the 5th composite library ([Table S7](#)) was
191 grown in LB medium with shaking at 28°C for 12 h, collected by centrifugation at 8,000 × g for
192 2 min and washed twice with filter sterilized PBS. Then, the input library was subjected to three
193 selective growth conditions as follows. For growth in DMEM, the input was diluted 1:100 into
194 DMEM and incubated at 28°C for 24 h after which genomic DNA extraction was performed.
195 For growth in macrophages, the input was inoculated into J774A.1 cell cultures at an MOI of
196 10 at 37°C for 2 h, followed by killing of extracellular bacteria with Gm. The J774A.1 cells were
197 incubated for an additional 4 h and lysed with 1% Triton X-100 as described ([Okuda et al., 2009](#)). Bacterial cells from the lysate were pelleted and resuspended into 30 mL LB with

199 shaking at 28°C until OD₆₀₀ reached 1.0 (~ 3 h), eliminating DNA contamination from dead *E.*
200 *piscicida* cells before genomic DNA extraction. For growth in turbot, the input was used to
201 intraperitoneally (i.p.) inoculate six-month old naïve turbot fish (~100 g), with 3 x 10⁶ CFU/fish.
202 At 14-days post infection (d.p.i.), each output library was generated from pooled livers of 5
203 cohoused fish. The pooled livers were homogenized and plated on the LBA plates
204 supplemented with Col and Gm. After overnight incubation at 28°C, bacterial colonies were
205 scraped, resuspended into fresh LB and ~1:10 back-diluted into 30 mL LB with shaking at
206 28°C until OD₆₀₀ 1.0 before genomic DNA was extracted ([Fu et al., 2013](#)).

207 ***Transposon insertion sequencing (TIS)***

208 TIS libraries were analyzed by the STAT-Tn-seq bioinformatic pipeline ([Fu et al., 2013](#)).
209 Briefly, genomic DNA was extracted and fragmented by sonication. Then, DNA fragments
210 were end-repaired, A-tailed and P5/P7 adaptor sequences were added by two rounds of PCR.
211 Experimental triplicate libraries of inputs and outputs were sequenced on an Illumina MiSeq
212 platform; ~ 2 million reads were generated for each library. The sequencing results were
213 processed with adapter trimming, mapping to the genome, and tallying as described ([Fu et al.,](#)
214 [2013](#)). The read counts for each locus were normalized among the three libraries according to
215 sequencing depth. The fold change of each locus was generated by dividing the output over
216 the input read counts.

217 ***HeLa, J774A.1, and turbot macrophage infection assays***

218 HeLa or macrophage cells (J774A.1 and turbot primary macrophages) were seeded at a
219 density of 1.0 × 10⁵ and 3.0 × 10⁵ cells/well, respectively, in 24-well plates and incubated
220 overnight at 37°C with 5% CO₂. *E. piscicida* cultures were inoculated into fresh DMEM and
221 statically grown for 12 h at 28°C. J774A.1 or HeLa cells were infected with *E. piscicida* at an
222 MOI of 10 or 100, respectively, followed by centrifugation at 600 × g for 10 min to facilitate
223 bacterial attachment to cells. After 2 h infection, cells were washed twice with PBS. DMEM
224 was added into cell cultures with 50 µg/mL Gm to kill extracellular bacteria and cells were
225 incubated at 37°C with 5% CO₂ for another 4-6 h. Then the cultures were treated for 10 min
226 with 1% Triton X-100 to disrupt the cells. Intracellular bacteria were enumerated by serial
227 dilution plating on LBA. For lactate dehydrogenase (LDH) detection, supernatants were
228 transferred to a new centrifuge tube and spun at 5,000 × g for 5 min at 4°C and LDH was
229 measured with a CytoTox 96 assay kit (Promega, USA). For fluorescence microscopy, cells
230 were washed twice with PBS and fixed with 4% paraformaldehyde for 10 min, washed with
231 PBS, permeabilized with 0.1% Triton X-100 for 5 min, and washed with PBS again. Nuclei

232 were stained with DAPI (Beyotime Biotechnology, China) for 30 s, washed 3x with PBS, and
233 coverslips were sealed with nail polish. Images were acquired by a Nikon A1R confocal
234 microscope and analyzed with NIS-Elements Viewer (Nikon, Japan).

235 **Total RNA extraction and qRT-PCR**

236 Overnight cultures of WT and Δ evrA were cultured statically in DMEM or DMEM
237 supplemented with various sugars at 28°C for 12 h, respectively. RNA samples were extracted
238 with a commercial RNA isolation kit (Tiangen, China) and mRNA was reverse-transcribed into
239 cDNA using the FastKing RT kit (Tiangen, China). qRT-PCR was performed with an Applied
240 Biosystems 7500 cycler (Applied Biosystems, USA) with triplicate reactions for each sample.
241 The comparative C_T ($2-\Delta\Delta C_T$) method ([Gu et al., 2016](#)) was used to quantify the relative levels
242 of each transcript with the housekeeping *gyrB* gene as an internal control and the specific
243 primer pair ([Yin et al., 2018](#)).

244 **RNA-seq**

245 For preparation of mRNA for RNA-seq, the Ribo-Zero-rRNA kit (Epicentre, USA) was
246 initially used to remove rRNA from the RNA samples. The final concentration of RNA samples
247 was determined with a Qubit 2.0 Fluorometer (Thermo Fisher, USA). The VAHTS Stranded
248 mRNA-seq Library Prep Kit for Illumina (Vazyme, China) was used to construct strand-specific
249 RNA-seq libraries, and sequencing was conducted on an Illumina HiSeq 2500 platform,
250 yielding 101-bp paired end-reads. Adapter sequences and low-quality bases (PHRED quality
251 scores ≤ 5) were trimmed with the Trimmomatic package ([Bolger et al., 2014](#)) using the default
252 parameters and reads smaller than 35 bp were discarded. The RNA-seq data processing
253 procedures and statistical analysis were performed ([Tjaden 2015](#)). The SRA accession
254 number for the RNA-seq data is SRP156435.

255 **SDS-PAGE and Western blotting analysis**

256 Whole cell proteins (WCPs) and extracellular proteins (ECPs) were extracted and
257 concentrated ([Yin et al., 2018; Zheng and Leung 2007](#)). Overnight cultures of *E. piscicida* were
258 subcultured into 50 mL fresh DMEM and statically incubated for 24 h at 28°C; bacteria were
259 then harvested by centrifugation at 5,000 $\times g$ for 10 min at 4°C for WCPs. For ECPs, culture
260 supernatants were filtered with 0.22 μm filters (Millipore, USA), and concentrated using 10 kDa
261 cutoff centrifugal filter devices (Millipore, USA). Proteins were separated by 12% SDS-PAGE,
262 followed by Coomassie Blue staining or Western blotting. For Western blots, separated
263 proteins were wet transferred onto PVDF membranes (Millipore, USA) and incubated with a
264 1:1000 dilution of mouse anti-EseB (GL Biochem, China). HRP-conjugated anti-mouse IgG

265 (Santa Cruz Biotechnology, CA) was used at a 1:2,000 dilution as a secondary antibody.
266 Proteins were visualized with TMB substrate (Amresco, USA). Mouse anti-DnaK (Santa Cruz
267 Biotechnology, USA) was used as a cytoplasmic protein control.

268 **Luminescence and fluorescence assays**

269 For assays involving luminescence and fluorescence, pUTt derivative plasmids bearing
270 P_{esrB} -*luxAB*, P_{esrC} -*luxAB* and P_{eseB} -*luc* (luciferase) reporters were introduced into WT and
271 Δ evrA *E. piscicida*. For luminescence assays, strains were inoculated into 50 mL DMEM and
272 statically incubated at 28°C. Pellets of 150 µL culture of each strain were mixed with 40 µL
273 capraldehyde dissolved in ethanol as substrate, and OD₆₀₀ was determined using a Microplate
274 Reader (Bio-Tek, USA) and luminescence values were monitored using a Microplate
275 Luminometer Orion II (Titertek-Berthold, Germany) every 2 hours.

276 For *in vivo* fluorescence detection, overnight cultures of WT+ P_{eseB} -*luc* and
277 Δ evrA+ P_{eseB} -*luc* were i.p. injected into turbot. At 5 d.p.i., the fish were anesthetized with
278 tricaine methanesulfonate (MS-222) (Sigma-Aldrich, USA) and i.p. injected 100 µL of a 1
279 mg/mL beetle luciferin solution (Promega, USA). At 10 min post-injection, fluorescence was
280 detected with a Kodak In-Vivo Multispectral System FX (Carestream Health, USA). After
281 fluorescence measurements were taken, the fish were killed with overdose of MS-222 and
282 livers from each fish were obtained for bacterial CFU plating ([Yin et al., 2018](#)).

283 **Protein purification**

284 Recombinant EvrA (WT and mutant variants) with N-terminal HisSumo tags were purified
285 from BL21 (DE3) *E. coli*. Expression of the EvrA-HisSumo fusion was induced by growth in LB
286 medium supplemented with 0.2 mM isopropyl β-D-1-thiogalacto-pyranoside at 22°C at 200
287 rpm. At 18 h post-induction, the cells were harvested by centrifugation, resuspended in lysis
288 buffer (20 mM Tris, 500 mM NaCl, pH 9.0) and lysed with a French press (Glen Mills, USA).
289 Following centrifugation at 12,000 rpm for 30 min, the supernatant was loaded onto a
290 pre-packed Ni-NTA column (GE Healthcare, Sweden) for purification. The loaded column was
291 washed with a lysis buffer gradient supplemented with 40-100 mM imidazole on an ÄKTA
292 protein purification system (GE, Healthcare, Sweden). Proteins were eluted with lysis buffer
293 supplemented with 500 mM imidazole. The SUMO protease ULP1 (Thermo Fisher, USA) was
294 added into the purified protein at the ratio 1:500 to digest HisSumo tag at 4°C in statics. After
295 digestion of the HisSumo tag with ULP1 for 10 h, reaction mixtures were reloaded on to a
296 Ni-NTA column to isolate EvrA protein in the flowthrough. Purified proteins were stored in 10%
297 glycerol at 4°C and their purity was confirmed by SDS-PAGE. Protein concentrations were

298 determined by the bicinchoninic acid protein assay (Thermo Fisher, USA).

299 ***Electrophoretic mobility shift assay (EMSA) and DNase I footprinting assay***

300 For EMSAs, purified EvrA protein was incubated with Cy5-labeled DNA probes (P_{esrB} ,
301 P_{esrC} , and P_{manX}) (Genewiz, China) in 20 μ L of binding buffer (10 mM Tris, 50 mM KCl, 5 mM
302 $MgCl_2$, 0.1 mM DTT, pH 7.4). In all EMSA analysis, excess (10-fold) of the nonspecific
303 competitor poly(dI-dC) was used to determine the specificity of the binding. After incubation at
304 25°C for 30 min, the samples were loaded on a 6% polyacrylamide gel and electrophoresed in
305 0.5 × TBE (Tris/Boric acid/EDTA) buffer at 4°C at 100 V for 120 min. Gels were imaged using a
306 Typhoon FLA 9500 (GE Healthcare, Sweden) with the Cy5 channel set at a 531 nm excitation
307 wavelength (Gu et al., 2016).

308 Dye primer-based DNase I footprinting assays were performed (Gu et al., 2016). Briefly,
309 the promoter region of *esrB* was PCR-amplified to include a 6-FAM moiety at the 5' end (S14
310 Table). For each assay, 200 ng of probe was incubated without or with the presence of 200 ng
311 EvrA in a total volume of 40 μ L. After the mixture was incubated for 30 min at 25°C, a 10 μ L
312 solution containing approximately 0.015 units of DNase I (Promega, USA) and 100 nmol of
313 freshly prepared $CaCl_2$ was added. The mixture was then incubated for 1 min at 25°C. The
314 reaction was stopped by adding 140 μ L of DNase I stop solution (200 mM unbuffered sodium
315 acetate, 30 mM EDTA and 0.15% SDS). The samples were first extracted using
316 phenol/chloroform and then precipitated using ethanol, and the pellets were dissolved in 10 μ L
317 of MilliQ water. Approximately 2 μ L of digested DNA was added to 7.9 μ L of HiDi formamide
318 (Applied Biosystems. USA) and 0.1 μ L of GeneScan-500 LIZ size standards (Applied
319 Biosystems, USA). The samples were analyzed using a 3730 DNA Analyzer with a G5 dye set
320 that was run on an altered default genotyping module that increased the injection time to 30 s
321 and the injection voltage to 3 kV. The results were analyzed using GeneMapper 4.0 (Applied
322 Biosystems, USA).

323 ***Chromatin immunoprecipitation quantitative PCR (ChIP-qPCR)***

324 ChIP-qPCR was performed (Liu et al., 2017). Briefly, strains expressing functional
325 FLAG-tagged EvrA or FLAG only vectors (WT+EvrA-Flag and WT+Flag) (Fig. S7C) were
326 cultured in DMEM at 28°C for 12 h without shaking. Bacteria were treated with 1%
327 formaldehyde at room temperature for 10 min and the cross-linking reaction was stopped with
328 125 mM glycine. The bacteria were then washed twice with cold sterile PBS and resuspended
329 in 5 mL of SDS lysis buffer (Liu et al., 2017). Next, the bacteria were sonicated, and the DNA
330 was fragmented to 100-500 bp at 200 W (Diaenode, USA). Insoluble cellular debris was

331 removed via centrifugation, and the supernatant was used as the input sample in IP
332 experiments. Both the input and the IP samples were washed with 50 μ L protein G beads for 1
333 h, and incubated overnight with 30 μ L anti-Flag M2 magnetic beads (Sigma-Aldrich, USA).
334 The beads were washed twice with 1 mL of each of the following buffers: low salt wash buffer,
335 high salt wash buffer, LiCl wash buffer, and standard TE buffer ([Liu e al., 2017](#)). The beads
336 were resuspended in 200 μ L of elution buffer, incubated at 65°C for 2 h, and then centrifuged
337 at 5,000 $\times g$ for 1 min. The supernatants containing the immunoprecipitated DNA were
338 collected, and 8 μ L of 5 M NaCl was added to all of the tubes (IPs and Inputs). The tubes were
339 then incubated at 65°C overnight to reverse the DNA-protein crosslinks. After treatment with
340 RNase A (10 μ g/mL) and Proteinase K (1 mg/mL), the enriched DNA was purified using
341 phenol-chloroform and amplified using qPCR.

342 For each DNA target, ΔC_T of the Input fraction and IP fraction was calculated in both the
343 WT+EvrA-Flag and WT+Flag samples. Each value was then divided by the corresponding ΔC_T
344 that was obtained for the non-specific *gyrB* intragenic region in the strains. Then, the
345 enrichment ratio was calculated from the $\Delta\Delta C_T$ value in WT+EvrA-Flag strain divided by that of
346 WT+Flag strain. The formula of EvrA binding is as following: $(\Delta\Delta C_T) = \text{IP}\{[C_T(\text{WT+EvrA-Flag}) - C_T(\text{WT+Flag})] - C_{T\text{gyrB}}\} - \text{Input}\{[C_T(\text{WT+EvrA-Flag}) - C_T(\text{WT+Flag})] - C_{T\text{gyrB}}\}$.

348 ***Electrospray ionization mass spectrometry (ESI-MS)***

349 ESI-MS based determination of EvrA interaction with man-6P ([Zhou et al., 2018](#)). Purified
350 EvrA (20 μ M) was buffer-exchanged into 100 mM ammonium acetate (pH 7.5) using a
351 centrifugal buffer exchange column (Micro Bio-Spin 6, Bio-Rad, USA), and one aliquot was
352 denatured by adding formic acid to a final concentration of 0.1%. Both native (non-formic
353 acid-treated) and denatured protein samples were analyzed by direct infusion. Specifically, 15
354 μ L of each protein sample was loaded into a nano-flow borosilicate emitter (NanoES spray
355 capillaries, Thermo Scientific, USA) and sprayed into an Orbitrap Fusion mass spectrometer
356 through a Nanospray FLEX Ion Source (Thermo Scientific, USA). The mass spectrometer
357 settings were: spray voltage 1.3 kV for native samples and 2.2 kV for denatured samples;
358 S-lens RF level of 150, SID 100 for complete desolvation of the native protein sample; capillary
359 temperature at 150°C for native protein samples or 300°C for denatured samples; scan range
360 1500–5500 m/z for native samples and 1000–5000 for denatured samples; intact protein mode
361 with trapping gas pressure set as 0.2. Mass spectra were analyzed using Thermo Scientific
362 Protein Deconvolution software (Thermo Scientific, USA). The parameters for spectra analysis
363 were specified according to the mass spectrometer settings. The minimum adjacent range of

364 charges was 4–8 for native proteins or 5–10 for the denatured proteins, and mass tolerance
365 was 30 p.p.m. The deconvoluted mass of the most abundant ion was selected as the mass of
366 the target protein. The mass of the bound ligand was calculated as the difference between the
367 native protein and the denatured protein.

368 ***Turbot virulence and competitive index assays***

369 Turbot experiments were performed according to protocols approved by the Animal Care
370 Committee of the East China University of Science and Technology (2006272) and the
371 Experimental Animal Care and Use Guidelines from the Ministry of Science and Technology of
372 China (MOST-2011-02). Healthy turbot weighing 30.0 ± 3.0 g (~2 months' old and ~1:1 female
373 to male) were obtained from a commercial farm (Yantai, China) and acclimatized to laboratory
374 conditions for at least 7 days. Competitive assays were performed between WT or the
375 indicated gene-deletion mutant strains and WT(Δp), the WT strain cured of its endogenous Cm
376 and Str resistance plasmid pEIB202 ([Wang et al., 2009](#)). WT(Δp) does not exhibit impaired
377 growth in LB, DMEM, J774A.1 or turbot. Inocula were prepared using fresh cultures of bacteria
378 that were diluted and mixed at a 1:1 ratio. The i.p. injection dose was $\sim 10^5$ CFU/fish in a 100
379 μL inoculum. At 8 d.p.i., the livers from fish in each group (5 animals/group) were sampled,
380 homogenized and plated on LBA plates with or without the presence of 34 $\mu\text{g/mL}$
381 chloramphenicol (Cm) to distinguish WT(Δp) (Cm^s) or other strains (Cm^r) ([Yang et al., 2017](#))
382 and enumerate the ratio of the competing strains. The ratios of the bacterial counts were used
383 to determine competitive indices.

384 For fish survival assays, overnight cultures were harvested by centrifugation at $8,000 \times g$
385 for 2 min at 4°C and washed three times with PBS. A total of 2.0×10^4 CFU bacteria
386 suspended in PBS containing 5 mg/mL mannose or glucose was i.p. injected into each fish;
387 PBS or PBS supplemented with 5 mg/mL mannose and glucose was used as negative control.
388 A total of 30 fish were injected with each strain and fish mortality was monitored daily. The
389 infection experiments were performed at least three independent times.

390 ***Turbot macrophage separation***

391 The separated head kidney of turbot fishes was homogenized with woven nylon mesh,
392 and macrophages obtained from the organ were collected using the continuous gradient
393 Percoll separation method ([Vray and Plasman 1994](#)). Nine volumes of Percoll with 1 volume of
394 sterile 1.5 M NaCl were mixed to make stock isotonic Percoll (SIP). The initial density of 1.065
395 g/mL was obtained by mixing 4 mL of the head kidney cell suspension in pH 7.6 heparin/L-15
396 with 4.2 ml of SIP. To make a self-generated continuous gradient, the prepared sample

397 solution was centrifuged for 20 min, 20,000 × g, at 5°C. Macrophages were obtained from the
398 third band with a density distribution of 1.069-1.075 g/mL. The collected cell suspended
399 solution was washed three times by centrifugation for 10 min at 300 × g and at 4°C with 10 mL
400 of incomplete L-15. The washed cell suspension was counted with a 0.4% solution of Trypan
401 Blue in HBSS to determine cell viability.

402 ***Isothermal titration calorimetry (ITC)***

403 All ITC titrations were performed at 25°C using a MicroCal iTC200 instrument (Malvern
404 Panalytical Ltd). A binding buffer consisting of 20 mM Tris and 500 NaCl (pH 9.0) was used for
405 all measurements. Assays were performed with 20 total injections with 100 µM EvrA in the
406 sample cell and 2 mM sugar in the injection syringe. Buffer-only runs were performed to
407 quantify the heat of dilution for background subtraction from binding runs. Thermal data were
408 fitted to the One Set of Sites binding model with the N value fixed at 1 to yield the equilibrium
409 dissociation constant (K_d) according to the modeling alignment to RpiA ([Ishikawa et al., 2002](#)).
410 Each binding run was carried out at least 2 independent times.

411 ***Quantification of mannose and man-6P/man-1P in bacterial cells and fish tissues***

412 Metabolites quantification was performed ([Cheng et al., 2019; Guo et al., 2014](#)). After 5 d.p.i.
413 with *E. piscicida* WT or PBS as a control, turbot livers and intestines were harvested and
414 weighted. To extract mannose and its derivatives, 1 mL methanol was added to the liver,
415 vortexed for 10 s, ultrasonicated for 20 min, and centrifuged for 20 min with 12,000 rpm at 4°C.
416 700 µL of the resulting supernatant was taken for UPLC-TQS MS/MS (Waters, USA).
417 UPLC-TQS MS/MS settings were set as follows: column temperature 35°C, sample chamber
418 temperature 10°C, carrier speed 0.3 ml/min, electronic press of spray 2.5 kV. The sample of
419 bacterial cells cultivated in DMEM condition was treated with the same procedures.

420 ***Bioinformatics analysis of EvrA***

421 Structural homology modeling of EvrA was performed with Phyre 2.0 ([Kelley et al., 2015](#)).
422 The structural model with the highest confidence was from an alignment to the structure of
423 RpiA (PDB: 1LK7) ([Ishikawa et al., 2002](#)). The Phyre-predicted structure of WT EvrA was used
424 for ligand (man-6P) docking with AutoDock (<http://autodock.scripps.edu>) with the default
425 docking parameters and the top ligand-binding site with lowest ΔG was selected. Multiple
426 sequence alignments were performed using ClusterX 1.8, and the phylogenetic tree was
427 drawn using MEGA 6.0 ([Tamura et al., 2013](#)). COG counts were based on the previous COG
428 annotation of EIB202 genome ([Wang et al., 2009](#)). EvrA was searched against Uniprot
429 database (<https://www.uniprot.org>) to predict its conserved domains. STRING analysis was

430 performed with the default parameters ([Szklarczyk et al., 2019](#)).

431 **Statistical analysis**

432 GraphPad Prism (version 6.0) was used to perform the statistical analyses. Data are
433 presented as the mean \pm SD of triplicate samples per experimental condition unless otherwise
434 noted. Statistical analyses were performed using unpaired two-tailed Student's *t* test for the
435 metabolite level analysis, One-way ANOVA analyses followed by Bonferroni's
436 multiple-comparison post-test comparing the data of CI values, or Kaplan-Meier survival
437 analysis with a log-rank test. Differences were considered significant at **P* < 0.05, ***P* < 0.01,
438 and ****P* < 0.001.

439

440 **Supplemental References**

- 441 Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina
442 sequence data. *Bioinformatics* 30, 2114-20.
- 443 Cheng, Z.X., Guo, C., Chen, Z.G., Yang, T.C., Zhang, J.Y., Wang, J., Zhu, J.X., Li, D., Zhang,
444 T.T., Li, H., Peng, B., and Peng, X.X. (2019). Glycine, serine and threonine metabolism
445 confounds efficacy of complement-mediated killing. *Nat. Commun.* 10, 3325.
- 446 Dennis, J.J., and Zylstra, G.J. (1998). Plasposons: modular self-cloning minitransposon
447 derivatives for rapid genetic analysis of Gram-negative bacterial genomes. *Appl. Environ.
448 Microbiol.* 64, 2710-5.
- 449 Fu, Y., Waldor, M.K., and Mekalanos, J.J. (2013). Tn-Seq analysis of *Vibrio cholerae* intestinal
450 colonization reveals a role for T6SS-mediated antibacterial activity in the host. *Cell Host
451 Microbe* 14, 652-63.
- 452 Gu, D., Guo, M., Yang, M.J., Zhang, Y.X., Zhou, X.H., and Wang, Q.Y. (2016). A σ^E -Mediated
453 temperature gauge controls a switch from LuxR-mediated virulence gene expression to
454 thermal stress adaptation in *Vibrio alginolyticus*. *PLoS Pathog.* 12, e1005645.
- 455 Guo, C., Huang, X.Y., Yang, M.J., Wang, S., Ren, S.T., Li, H., and Peng, X.X. (2014).
456 GC/MS-based metabolomics approach to identify biomarkers differentiating survivals from
457 death in crucian carps infected by *Edwardsiella tarda*. *Fish Shellfish Immunol.* 39,
458 215-222.
- 459 Ishikawa, K., Matsui, I., Payan, F., Cambillau, C., Ishida, H., Kawarabayasi, Y., Kikuchi, H.,
460 and Roussel, A. (2002). A hyperthermostable D-ribose-5-phosphate isomerase from
461 *Pyrococcus horikoshii* characterization and three-dimensional structure. *Structure* 10,
462 877-886.

- 463 Jacobs, M.A., Alwood, A., Thaipisuttikul, I., Spencer, D., Haugen, E., Ernst, S., Will, O., Kaul,
464 R., Raymond, C., Levy, R., Chun-Rong, L., Guenthner, D., Bovee, D., Olson, M.V., and
465 Manoil C. (2003). Comprehensive transposon mutant library of *Pseudomonas aeruginosa*.
466 Proc. Natl. Acad. Sci. U. S. A. 100, 14339-44.
- 467 Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E. (2015). The Phyre2
468 web portal for protein modeling, prediction and analysis. Nat. Protocols 10, 845-858.
- 469 Liu, Y., Zhao, L.Y., Yang, M.J., Yin, K.Y., Zhou, X.H., Leung, K.Y., Liu, Q., Zhang, Y.X., and
470 Wang, Q.Y. (2017). Transcriptomic dissection of the horizontally acquired response
471 regulator EsrB reveals its global regulatory roles in the physiological adaptation and
472 activation of T3SS and the cognate effector repertoire in *Edwardsiella piscicida* during
473 infection toward turbot. Virulence 8, 1355-77.
- 474 Liu, Y.G., and Chen, Y. (2007). High-efficiency thermal asymmetric interlaced PCR for
475 amplification of unknown flanking sequences. BioTechniques 43, 649-56.
- 476 Okuda, J., Kiriyama, M., Suzuki, E., Kataoka, K., Nishibuchi, M., and Nakai, T. (2009).
477 Characterization of proteins secreted from a type III secretion system of *Edwardsiella*
478 *tarda* and their roles in macrophage infection. Dis. Aquat. Organ. 84, 115-21.
- 479 Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
480 Doncheva, N.T., Morris, J.H., Bork, P., Jensen, L.J., and von Mering, C. (2019). STRING
481 v11: protein-protein association networks with increased coverage, supporting functional
482 discovery in genome-wide experimental datasets. Nucleic Acids Res. 47, D607-613.
- 483 Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. (2013). MEGA6: molecular
484 evolutionary genetics analysis Version 6.0. Mol. Biol. Evol. 30, 2725-2729.
- 485 Tjaden, B. (2015). De novo assembly of bacterial transcriptomes from RNA-seq data. Genome
486 Biol. 16, 1.
- 487 Vray, B., and Plasman, N. (1994). Separation of murine peritoneal macrophages using Percoll
488 density gradients. J. Immunol. Methods 174, 53-9.
- 489 Wang, Q.Y., Yang, M.J., Xiao, J.F., Wu, H.Z., Wang, X., Lv, Y.Z., Xu, L.L., Zheng, H.J., Wang,
490 S.Y., Zhao, G.P., Liu, Q., and Zhang, Y.X. (2009). Genome sequence of the versatile fish
491 pathogen *Edwardsiella tarda* provides insights into its adaptation to broad host ranges and
492 intracellular niches. PLoS ONE 4, e7646.
- 493 Yang, G.H., Billings, G., Hubbard, T.P., Park, J.S., Leung, K.Y., Liu Q., Davis, B.M., Zhang,
494 Y.X., Wang, Q.Y., and Waldor, M.K. (2017). Time-resolved transposon insertion
495 sequencing reveals genome-wide fitness dynamics during infection. mBio. 8, e1517-81.

- 496 Yin, K.Y., Guan, Y.P., Ma, R.Q., Wei, L.F., Liu, B., Liu, X.H., Zhou, X.S., Ma, Y., Zhang, Y.X.,
497 Waldor, M.K., and Wang, Q.Y. (2018). Critical role for a promoter discriminator in RpoS
498 control of virulence in *Edwardsiella piscicida*. PLoS Pathogen. 14, e1007272.
- 499 Zheng, J., and Leung, K.Y. (2007). Dissection of a type VI secretion system in *Edwardsiella*
500 *tarda*. Mol. Microbiol. 66, 1192-206.
- 501 Zhou, P., She, Y., Dong, N., Li, P., He, H., Borio, A., Wu, Q., Lu, S., Ding, X., Cao, Y., Xu, Y.,
502 Gao, W., Dong, M., Ding, J., Wang, D.C., Zamyatina, A., and Shao, F. (2018).
503 Alpha-kinase 1 is a cytosolic innate immune receptor for bacterial ADP-heptose. Nature
504 561, 122-126.
- 505