

Supplemental Material

Table S1. Predictors of Procedural Complications.

Variable	Univariate Logistic Regression			Multivariate Logistic Regression		
	OR	95%CI	P	OR	95%CI	P
Age (per year)	1.02	1.00-1.04	0.012	1.02	1.01-1.04	0.007
Female sex	1.20	0.84-1.70	0.318	-	-	-
AF duration (per year)	1.02	0.99-1.01	0.102	-	-	-
Paroxysmal AF	0.91	0.66-1.27	0.591	-	-	-
Congestive heart failure	1.17	0.66-2.07	0.583	-	-	-
Hypertension	0.81	0.58-1.13	0.205	-	-	-
Diabetes mellitus	0.59	0.3-1.17	0.133	-	-	-
Stroke or TIA	1.36	0.78-2.37	0.276	-	-	-
Vascular disease	0.91	0.49-1.66	0.748	-	-	-
Obstructive Sleep apnea	1.24	0.69-2.24	0.475	-	-	-
CHA ₂ DS ₂ -VASc	1.06	0.94-1.19	0.351	-	-	-
BMI (per Kg/m ²)	0.98	0.94-1.01	0.187	-	-	-
eGFR (per ml/min)	0.99	0.98-0.99	0.040	-	-	-
Indexed LA volume (per mL/m ²)	0.99	0.98-1.00	0.196	-	-	-
LVEF (per %)	1.01	0.99-1.03	0.394	-	-	-
NOAC	0.64	0.43-0.97	0.033	0.61	0.40-0.92	0.018
Center Case-Load (per 100 ablations/year)	1.10	1.01-1.20	0.030	-	-	-

HR – hazard ratio; CI – confidence interval; AF - atrial fibrillation; TIA - transitory ischemic attack; CHA₂DS₂-VASc - cardiac failure or dysfunction, hypertension, age ≥75 years [doubled], diabetes, stroke [doubled] - vascular disease, age 65–74 years, sex category [female]; BMI - body mass index; LA - left atrium; LVEF - left ventricular ejection fraction.