

OMTN, Volume 18

Supplemental Information

**circIFT80 Functions as a ceRNA of miR-1236-3p
to Promote Colorectal Cancer Progression**

Wenming Feng, Hui Gong, Yongchun Wang, Guoliang Zhu, Tao Xue, Yao Wang, and Ge Cui




Fig. S1 A. The qRT-PCR assay indicated the expression level of circIFT80 in HCT116 cells treated with si-circIFT80. Data are the means \pm SD of three experiments, $**P<0.01$; B. The qRT-PCR assay indicated the expression level of IFT80 in HCT116 cells treated with si-circIFT80. Data are the means \pm SD of three experiments; C. The qRT-PCR assay indicated the expression level of circIFT80 in HCT116 cells treated with si-IFT80. Data are the means \pm SD of three experiments; D. The qRT-PCR assay indicated the expression level of IFT80 in HCT116 cells treated with si-IFT80. Data are the means \pm SD of three experiments, $**P<0.01$; E. The qRT-PCR assay indicated the expression level of circIFT80 in HT-29 cells infected with circIFT80 overexpression plasmid, $**P<0.01$; F. The qRT-PCR assay indicated the expression level of IFT80 in HT-29 cells infected with circIFT80 overexpression plasmid, $**P<0.01$; G. circIFT80 was resistant to RNase R digestion in HCT116 cells. Data are the means \pm SD of three experiments, $**P<0.01$; H. circIFT80 was resistant to RNase R digestion in DLD1 cells. Data are the means \pm SD of three experiments, $**P<0.01$;