


SUPPLEMENTAL MATERIAL

**Alleviation of murine osteoarthritis by deletion of the focal  
adhesion mechanosensitive adapter, Hic-5**


Aya Miyauchi<sup>1</sup>, Joo-ri Kim-Kaneyama<sup>1\*</sup>, Xiao-Feng Lei<sup>1</sup>, Song Ho Chang<sup>2</sup>, Taku Saito<sup>2</sup>, Shogo Haraguchi<sup>1</sup>, Takuro Miyazaki<sup>1</sup> and Akira Miyazaki<sup>1</sup>

<sup>1</sup>Department of Biochemistry, Showa University School of Medicine, Tokyo, Japan


<sup>2</sup>Sensory & Motor System Medicine, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan




Supplementary Figure S1. Uncropped scans of Western blots. (Fig. 4A)




Supplementary Figure S2. Uncropped scans of Western blots. (Fig. 4B)





Supplementary Figure S3. Uncropped scans of Western blots. (Fig. 4D)



Supplementary Figure S4. Uncropped scans of Western blots. (Fig. 4E)



Supplementary Figure S5. Uncropped scans of Western blots. (Fig. 5A)



Supplementary Figure S6. Uncropped scans of Western blots. (Fig. 5C)