

Dopamine agonists in prolactinoma

Table S3. Anthropometric measures before and after treatment with dopamine receptor antagonists in patients with prolactinomas.

Study (Ref.nr.)	Weight baseline, Kg (SD)	Weight follow-up, Kg (SD)	BMI baseline (SD)	BMI follow-up (SD)	Waist-hip ratio baseline (SD)	Waist-hip ratio follow-up (SD)	WC baseline, cm (SD)	WC follow-up, cm (SD)
Schwetz (25)	-	-	27.9 (5.9)	28.6 (5.6)	-	-	-	-
Auriemma II (26)	96.1 (12.2)	88.8 (10.3)	31.7 (3.9)	29.3 (3.4)	-	-	105(10.1)	101 (10.3)
Medic (31)	-	-	24 (6.4)	22.8 (6.1)	-	-	-	-
Pala (27)	57.3 (9.5)	54.8 (9.2)	24.2 (4.0)	23.2 (3.9)	0.90 (0.06)	0.90 (0.07)	85.3 (12.0)	82.3 (10.0)
Barbosa (33)	-	-	29.3 (15.4)	28.4 (14.5)	-	-	88.3 (11.8)	89.8 (11.3)
Auriemma (5)	75.6 (18.4)	69.6 (15.6)	27.6 (5.3)	24.3 (4.7)	-	-	88.7 (14.6)	73.4 (5.9)
Ciresi (4)	-	-	25.6 (5.2)	25.4 (5.0)	-	-	88.8 (10.5)	87.1 (10.2)
Inancli (29)	-	-	27.1 (5.9)	26.7 (5.6)	-	-	-	-
Berinder (6)	78.5 (28.5)	75.5 (23.3)	25.8 (7.8)	25.3 (7.2)	0.87 (0.10)	0.87 (0.10)	90.2 (12.5)	87.7 (8.8)
Silva (8)	-	-	29.2 (15.3)	28.4 (14.5)	-	-	87.3 (12.5)	84.3 (11.2)
Serri (7)	82.5 (20.5)	82.5 (20.5)	29.0 (6.0)	-	-	-	-	-
Yavuz (28)	-	-	26.3 (5.3)	25.0 (5.3)	0.83 (0.06)	0.82 (0.06)	-	-
Doknic (30)	85.8 (17.4)	80.8 (16)	27.5 (3.4)	26.0 (3.1)	-	-	-	-
Iglesias (32)	-	-	31.3 (5.1)	-	-	-	-	-

(-) Not reported, WC: waist circumference