

eTable 1. ICD-10 codes used to define maternal drug-related deaths.

ICD-10 code	Diagnosis
Accidental poisoning	X40-X44
Poisoning of undetermined intent by exposure to drugs	Y10-Y14
Drug-induced diseases	D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, K85.3, L10.5, L27.0, L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2
Drugs in the blood	R78.1-R78.5
Mental/behavioral disorders due to drugs	F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7-F15.9, F16.0-F16.5, F16.7-F16.9, F18.0-F18.5, F18.7-F18.9, F19.0-F19.5, F19.7-F19.9

eTable 2. Sociodemographic characteristics of delivery hospitalizations excluded from analytic sample, California, 2010-2012.

Sociodemographic characteristic	N (%)
Age in years	
<20 years	33,056 (8.2%)
20-35 years	317,199 (78.9%)
≥35 years	51,995 (12.9%)
Race/ethnicity	
White, non-Hispanic	56,437 (14.0%)
Black, non-Hispanic	10,832 (2.7%)
Hispanic	280,223 (69.7%)
Asian/Pacific Islander	39,113 (9.7%)
Other	15,645 (3.9%)
Insurance/payer	
Private	96,378 (24.0%)
Medicaid	281,063 (69.9%)
Self-pay	20,021 (5.0%)
Other	4,788 (1.2%)