

**Supplementary Table S1. Clinical parameters of pancreatic cancer patients in TMAs**

<b>Variable</b>	<b>No. of Patients (%)</b>
<b>Sex</b>	
Men	60 (65.9)
Women	31 (34.1)
<b>Age</b>	
≤65	46 (50.5)
>65	45 (49.5)
<b>Tumor location</b>	
Head	55 (60.4)
Neck	7 (7.7)
Body/tail	16 (17.6)
Uncinate process	13 (14.3)
<b>Tumor size, cm</b>	
≤3	48 (52.7)
>3	43 (47.3)
<b>Margin status</b>	
R0	64 (70.3)
R1	23 (25.3)
R2	4 (4.4)
<b>Tumor grade</b>	
Poorly differentiated	17 (18.7)
Moderately differentiated	50 (54.9)
Well differentiated	24 (26.4)
<b>Stage</b>	
I	11 (12.1)
II	74 (81.3)
III	4 (4.4)
IV	2 (2.2)
<b>Recurrence status</b>	
Yes	61 (67.0)
No	30 (33.0)
<b>Metastasis status</b>	
Yes	68 (74.7)
No	23 (25.3)
<b>Lymph node status</b>	
Negative	44 (48.4)
Positive	47 (51.6)
<b>Adjuvant therapy</b>	
Yes	37 (40.7)
No	54 (59.3)
<b>CA19-9, U/mL</b>	
≤37	19 (20.9)
>37	72 (79.1)