

Supplementary Table 1. Primer sequences used in this study

Gene names	Forward primer (5'-3')	Reverse primer (5'-3')
mFabp4	CCGCAGACGACAGGA	CTCATGCCCTTCATAAACT
mPparg	GGAAGACCCTCGCATTCCCTT	GTAATCAGCAACCATTGGGTCA
mCebpa	GCGGGAACGCAACAAACATC	GTCACTGGTCAACTCCAGCAC
mGlut4	ACACTGGTCCTAGCTGTATTCT	CCAGGCCACGTTGCATTGTA
mNos2	CCAAGCCCTCACCTACTTCC	CTCTGAGGGCTGACACAAGG
mF4/80	CTTTGGCTATGGGCTTCCAGTC	GCAAGGAGGACAGAGTTATCGTG
mArg1	CTCCAAGCCAAAGTCCTTAGAG	AGGAGCTGTCATTAGGGACATC
mTnfa	ACGGCATGGATCTCAAAGAC	AGATAGCAAATCGGCTGACG
mChi313	AGAAGGGAGTTCAAACCTGGT	GTCTTGCTCATGTGTGTAAGTGA
mGapdh	TGAAGCAGGCATCTGAGGG	CGAAGGTGGAAGAGTGGGAG

mFabp4, mouse fatty acid-binding protein 4; mPparg, mouse peroxisome proliferator-activator- γ ; mCebpa, mouse CCAAT/enhancer binding protein- α ; mGlut4, mouse glucose transporter type 4; mNos2, mouse nitric oxide synthase 2; mArg1, mouse arginase 1; mTnfa, mouse tumor necrosis factor- α ; mChi313, mouse chitinase-like lectin; mGapdh, mouse glyceraldehyde 3-phosphate dehydrogenase.