Appendix 1. Prevalence Rates* of Depressive Disorders Recorded per 1,000 Delivery Hospitalizations, by State and Year; National Inpatient Sample and State Inpatient Databases, Healthcare Cost and Utilization Project, 2000–2015†

State	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	AARC‡
National	4.1	5.5	7.3	9.6	12.1	13.9	14.0	15.6	18.1	17.6	21.8	22.1	21.6	23.6	26.4	28.7	1.6¶
Arizona	2.8	4.0	6.1	8.0	10.1	10.5	10.9	10.7	13.0	13.8	15.9	17.6	17.4	19.0	20.1	21.4	1.2¶
Arkansas					5.5	8.4	11.5	10.3	9.6	13.7	13.7	13.9	15.1	18.3	20.3		1.2¶
California	2.3	3.1		5.2	6.5	7.6	7.8	8.6	9.5	10.6	12.5	14.0					1.0¶
Colorado	4.9	8.7	12.7	14.9	19.1	22	23.4	25.2	28.9	31.2	33.5	36.5	31.6	34.3	37.4	39.8	2.4¶
D.C.														16.9	19.0		\$
Florida	2.8	4.3	5.6	7.1	9.2	9.3	10.2	10.7	11.9	12.8	13.6	14.0	13.7	14.3	16.2		1.0¶
Georgia													13.1	12.8	15.0		0.9
Hawaii	2.5	3.7	3.6	6.0	6.1	6.0	7.6	5.2	7.8	9.0	9.3	11.8	9.5	12.7	12.6		0.7¶
Iowa	5.2	5.5	8.8	12.4	17	17.1	17.5	21.3	21.4	24.1	25.8	30.8	29.1	33.1	34.1	36.8	2.2¶
Kentucky	3.5	4.4	6.0	9.0	11.4	11.9	12.0	12.1	15.3	19.4	21.3	22.7	23.8	24.6	26.9	27.6	1.7¶
Maine	15.2	20.7	27.8	41.3			42.4	50.7	53.9	59.0	55.9	53.4	56.6				3.8¶
Maryland	5.6	9.0	13.2	15.5	21.5	24.2	28.2	30.5	31.5	33.4	37.5	38.9	39.7	41.4	45.4		2.9¶
Massachusetts	9.7	11.5	15.2	18.5	21.4	23.3	25.6	28.6	27.7	32.5	36.9	40.1	37.8	40.3			2.5¶
Michigan	4.9	6.1	9.3	11.1	14.6	19.4	21.1	22.4	25.4	27.8	29.2	30.7	30.7	35.8	39.3		2.5¶
Minnesota															51.1	53.8	\$
Mississippi											8.9	9.0					§
Nebraska		6.3	9.6	14.2	14.2	13.5	14.8	17.2	21.9	23.0	24.0	26.0	22.5	28.7	29.9	33.8	1.7¶
Nevada			2.3	3.7	5.0	4.9	5.6	5.8	6.4	7.1	7.7	9.1	8.8	8.4	13.8		0.7¶
New Jersey	3.1	3.5	5.3	6.8	8.2	9.5	11.5	15.9	19.2	17.4	18.6	17.5	16.1	18.6	19.4		1.4¶
New Mexico									15.4	19.1	20.8	26.2	25.7	27.1	29.2		2.3¶
New York	3.6	4.7	6.6	8.6	10.0	11.4	12.7	13.7	15.6	15.8	18.4	20.1	19.3	20.1	22.5		1.4¶

Haight SC, Byatt N, Moore Simas TA, Robbins CL, Ko JY. Recorded diagnoses of depression during United States delivery hospitalizations, 2000–2015. Obstet Gynecol 2019;133.

The authors provided this information as a supplement to their article.

©2019 American College of Obstetricians and Gynecologists.

North Carolina	4.5	6.5	9.3	13.0	15.4	16.2	18.0	17.5	18.4	20.6	22.5	24.1	25.1	28.9	35.8	39.7	1.9 [¶]
Oregon	7.6	11.9	15.2	16.4	19.8	21.3	21.2	23.7	28.6	30.3	34.3	36.7	40.2	41.3	49.8		2.6¶
Rhode Island			25.9	25.4	30.7	34.5	32.3	33.4	34.6	31.7	37.5	49.1	40.7	38.4	45.3		1.5¶
South Carolina	3.6	4.6	7.2	9.5	11.3	11.7	11.6	12.4	16.5	20.3	25.0	26.2	25.9	28.2			1.8¶
South Dakota								20.3	21.3	28.7	28.8	32.3	34.3	38.7	43.4		3.2¶
Utah	5.0	7.4	11.7	14.5	17.6	17.3	17.7	20.5	24.6	29.9	29.4	32.0	31.4	36.8	37.5		2.3¶
Vermont		8.8	8.8	11.6	18.6	26.4	31.6	48.7	43.3	47.5	56.4	52.1	56.4	75.4	76.3		5.3¶
Washington	6.0	7.4	10.1	11.9	13.4	13.4	15.0	15.7	18.2	22.4	24.0	25.6	26.6	32.0	37.8		1.9¶
West Virginia	5.1	8.0	8.9	11.4	13.8	16.7	19.7	21.5	19.2	20.9	22.4	20.7	26.3	28.7	31.6		1.7¶
Wisconsin	7.6	9.9	13.2	18.9	23.4	26.1	29.0	32.5	33.8	35.3	40.0	41.8	39.2	43.4	49.3	52.0	3.0 [¶]

⁻⁻⁻ Data not available

^{*}Prevalence rate numerator consisted of depressive disorder ICD-9-CM codes (291.89, 292.84, 293.83, 296.2-296.26, 296.3-296.36, 300.4, and 311), prevalence rate denominator consisted of delivery hospitalization discharges

[†]Rates are for 2000 through the third quarter of 2015.

[‡]Estimated average annual rate change

[§] Insufficient data (<3 years) to assess linear trend.

[¶] Trend significantly increasing at P < 0.05.