

Supplemental Table 2 Results from simple logistic regression analysis of dichotomous PCS and MCS ten years after a *Giardia*-outbreak with respect to eight-categories exposure (IBSxCFxExposure status) after three, six and ten years, respectively

PCS	OR for PCS below 45			P-value			N			
	3 years	6 years	10 years	3 years	6 years	10 years	3 years	6 years	10 years	
Exposure status										
Neither condition among controls	ref	ref	ref	ref	ref	ref	364	385	532	
Neither condition among exposed	1.3	1.0	1.0	.52	.97	.99	163	226	280	
IBS-only among controls	2.4	2.0	4.4	.079	.13	<.001	41	46	75	
IBS-only among exposed	1.9	2.6	2.4	.11	.003	.003	85	99	146	
CF-only among controls	4.6	3.4	6.1	.006	.005	<.001	22	36	50	
CF-only among exposed	4.7	8.8	14.0	<.001	<.001	<.001	68	44	45	
IBS and CF among controls	10.5	12.9	6.9	<.001	<.001	<.001	20	12	18	
IBS and CF among exposed	9.7	8.1	17.4	<.001	<.001	<.001	142	97	101	
							Total N	905	945	1247

MCS	OR for MCS below 45			P-value			N			
	3 years	6 years	10 years	3 years	6 years	10 years	3 years	6 years	10 years	
Exposure status										
Neither condition among controls	ref	ref	ref	ref	ref	ref	368	390	536	
Neither condition among exposed	1.1	1.4	1.6	.74	.14	.030	164	226	282	
IBS-only among controls	1.5	0.3	1.7	.38	.077	.091	42	46	75	
IBS-only among exposed	2.3	2.0	2.1	.004	.015	.002	85	99	146	
CF-only among controls	2.8	2.9	9.6	.036	.006	<.001	22	36	51	
CF-only among exposed	3.9	4.2	7.6	<.001	<.001	<.001	68	44	45	
IBS and CF among controls	2.3	1.9	15.9	.11	.36	<.001	20	12	19	
IBS and CF among exposed	4.7	7.4	11.4	<.001	<.001	<.001	142	97	101	
							Total N	911	950	1255

PCS = Physical component summary (0-100); MCS = Mental component summary (0-100); IBS = Irritable bowel syndrome; CF = Chronic fatigue
 Non-significant p-values are highlighted