

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Exposure to Bisphenol A and Bisphenol S and Incident Type 2 Diabetes: A Case–Cohort Study in the French Cohort D.E.S.I.R.

Fanny Rancière, Jérémie Botton, Rémy Slama, Marlène Z. Lacroix, Laurent Debrauwer, Marie Aline Charles, Ronan Roussel, Beverley Balkau, Dianna J. Magliano, and the D.E.S.I.R. Study Group

Table of Contents

Table S1. Baseline characteristics of D.E.S.I.R. individuals included and not included in the random subcohort after applying exclusion criteria.

Table S2. Adjusted associations between exposure to BPA and BPS and the risk of type 2 diabetes in the D.E.S.I.R. cohort (models including both BPA and BPS).

Table S3. Adjusted associations between average BPA exposure and average BPS detection and the risk of type 2 diabetes among men and women in the D.E.S.I.R. cohort.

Table S4. Adjusted associations between average BPS detection and the risk of type 2 diabetes according to baseline BMI in the D.E.S.I.R. cohort.

Figure S1. Flow chart of the case-cohort study nested in the French prospective cohort study D.E.S.I.R. when considering exposure at year 3 in relation to incident T2D.

Figure S2. Relationship between log-transformed BPA-G concentration at baseline and adjusted hazard ratio (HR) of type 2 diabetes in the D.E.S.I.R. case-cohort study fitted with restricted cubic splines (three knots placed at the 10th, 50th and 90th percentiles).