

Gene	Females		Males	
	+/+	Δ PMT	+/+	Δ PMT
<i>Il-1β</i>	2.1 \pm 0.31 (6)	2.9 \pm 0.71 (6) ^a	2.2 \pm 1.70 (6)	0.3 \pm 0.06 (6)
<i>Il1ra</i>	0.9 \pm 0.16 (6)	3.8 \pm 1.63 (5)	1.4 \pm 0.35 (6)	1.5 \pm 0.79 (6)
<i>Il-6</i>	0.8 \pm 0.13 (5)	8.6 \pm 4.68 (4)	2.3 \pm 1.97 (6)	1.9 \pm 0.61 (6)
<i>Tnf-α</i>	<i>n.p.</i>	1.9 \pm 0.83 (4)	5.0 \pm 4.11 (6)	0.1 \pm 0.02 (6)
<i>Hmox1</i>	<i>n.p.</i>	1.5 \pm 0.53 (4)	0.9 \pm 0.20 (4)	7.4 \pm 1.20 (4) ^c
<i>Nqo1</i>	<i>n.p.</i>	1.7 \pm 0.34 (4)	1.0 \pm 0.25 (4)	1.7 \pm 0.23 (4)
<i>Apcs</i>	5.1 \pm 1.44 (4)	6.9 \pm 2.09 (4)	2.3 \pm 0.71 (4)	1.3 \pm 0.73 (4)
<i>Gpx</i>	<i>n.p.</i>	0.33 \pm 0.09 (4) ^b	<i>n.p.</i>	<i>n.p.</i>
<i>Lpin</i>	<i>n.p.</i>	0.5 \pm 0.08 (4)	<i>n.p.</i>	<i>n.p.</i>

S2 Table. An analysis of hepatic mRNA transcripts demonstrates no consistent damage response in the liver after acute acrylamide exposure.