

Supporting Information

Article title: **Tomato's green gold: Bioeconomy potential of residual tomato leaf biomass as a novel source for the secondary metabolite rutin**

Laura V. Junker-Frohn*^{1,6}, Manuel Lück^{2,6}, Simone Schmittgen^{3,6}, Joana Wensing^{4,6}, Laura Carraresi^{4,6}, Björn Thiele^{1,6}, Tanja Groher^{3,6,a}, Julia J. Reimer^{5,6}, Stefanie Bröring^{4,6}, Georg Noga^{3,6}, Andreas Jupke^{2,6}, Ulrich Schurr^{1,6}, Björn Usadel^{1,5}, Anika Wiese-Klinkenberg^{1,6}, Alexandra Wormit^{5,6}

¹ Institute of Bio and Geosciences, Plant Sciences (IBG-2), Forschungszentrum Jülich GmbH, Jülich, Germany.

² Chair of Fluid Process Engineering (AVT.FVT), RWTH Aachen University, Aachen, Germany.

³ Institute of Plant Sciences and Resource Conservation (INRES), Horticultural Sciences, University of Bonn, Bonn, Germany.

⁴ Institute for Food and Resource Economics (ILR), Chair for Technology and Innovation Management in Agribusiness, University of Bonn, Bonn, Germany.

⁵ Institute for Biology I - Botany, RWTH Aachen University, Aachen, Germany.

⁶ Bioeconomy Science Center, c/o Forschungszentrum Jülich, 52425 Jülich, Germany.

The following Supporting Information is available for this article:

^a Current address: Agroscope, Taenikon 1, 8356 Ettenhausen, Switzerland

Figure S1:




Figure S1: Effect of stress treatments on young tomato plants over 16 days of nitrogen deficiency in combination with increased light intensities and/or chilling temperatures compared to control plants. a) Plant biomass as fresh weight (g). b) Total leaf area (cm²). Data represent mean of N=4 plants (\pm SD).