

Supplementary Table S4. Histopathologic data and stage according to sex

Characteristic	Men (n=113)	Women (n=76)	p-value
Tumor size (cm)			.537
Mean (95% CI)	2.6 (2.3–2.8)	2.8 (2.3–3.4)	
Depth of invasion (mm)			.503
Mean (95% CI)	10.5 (9.1–11.9)	11.1 (9.5–12.7)	
Histologic grade			.704
Well differentiated	56 (49.6)	33 (43.4)	
Moderately differentiated	43 (38.1)	32 (42.1)	
Poorly differentiated	14 (12.4)	11 (14.5)	
Lymphovascular invasion			.281
Present	30 (26.5)	15 (19.7)	
Absent	83 (73.5)	61 (80.3)	
Perineural invasion			.352
Present	40 (35.4)	32 (42.1)	
Absent	73 (64.6)	44 (57.9)	
Tumor infiltrating lymphocytes			.107
Low	66 (58.4)	42 (55.3)	
Intermediate	34 (30.1)	17 (22.4)	
High	13 (11.5)	17 (22.4)	
T category			.465
T1	27 (23.9)	12 (15.8)	
T2	39 (34.5)	30 (39.5)	
T3	46 (40.7)	34 (44.7)	
T4	1 (0.9)	0	
N category			.816
N0	58 (51.3)	42 (55.3)	
N1	17 (15.0)	13 (17.1)	
N2	24 (21.2)	12 (15.8)	
N3	14 (12.4)	9 (11.8)	

Values are presented as number (%) unless otherwise indicated.
CI, confidence interval.