

**Supplementary Table S1.** A literature review of clinicopathologic features in 10 cases (including the present case) with concurrent anti-GBM nephritis and IgA nephropathy

Reference, year	Country	Number	Sex (M/F)	Age (yr)	Present illness	Hypertension	Diabetes	History of upper respiratory tract infection	Previous history of proteinuria and hematuria	Hemoptysis	Oliguria	Nephrotic syndrome	HBV infection	24 Urine proteinuria (g/day) or spot urine	Creatinine (mg/dl)	Anti-GBM Ab	Crescent (%)	Treatments	Plasmapheresis	Prognosis	Hemodialysis	Other specific feature
Trpkov et al. (1998) <sup>1</sup>	Canada	1	M	54	Anorexia Malaise Lethargy	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Negative	≥2	8.1	Postive (not described)	25%	IV Metyl-PD+IV cyclophosphamide	Performed	Improved	Death due to GI bleeding	Transplant due to Goodpasture disease
Cui et al. (2006) <sup>2</sup>	China	1	F	27	Oliguria	Negative	Negative	Positive	Negative	Negative	Positive	Negative	Negative	Postive (not described)	15.2	134%	100%	IV Metyl-PD+ Oral-PD+IV cyclophosphamide	Not performed	Not improved	Dependent	
Wechsler et al. (2008) <sup>3</sup>	USA	1	M	55	Gross hematuria	Positive	Positive	Negative	Positive	Negative	Negative	Negative	Negative	2.8	3.5	8.6 EU/mL	20%	IV Metyl-PD+IVIG + Mycophenolate mofetil + Rituximab	Not performed	Improved	Independent	Acquired immune deficiency syndrome
Wang et al. (2012) <sup>4</sup>	China	1	F	31	Edema Gross hematuria	164/74 mm Hg Negative	Negative	Negative	Negative	Negative	Negative	Positive	Negative	3.8	3.2	93.5 RU/mL	2/10 cellular 100%	IV Metyl-PD+ Oral-PD+IV cyclophosphamide	Performed	Not improved	Independent	Septic arthritis
Yamaguchi et al. (2013) <sup>5</sup>	Japan	1	M	46	Azotemia Fever	Positive, but controlled 134/83 mm Hg	Positive	Negative	Positive Hematuria, proteinuria	Negative		Negative	Negative	2.9	5.2	214 EU/mL	15/16 cellular 94%	IV Metyl-PD+Oral-PD	Performed Failed due to allergy	Not improved	Dependent	
Gao et al. (2014) <sup>6</sup>	China	1	F	38	Anorexia Fatigue	Negative	Negative	Positive	Negative	Negative	Negative	Negative	Negative	2.2	2.8	237.6 EU/mL	28%	IV Metyl-PD+ Oral-PD+IV cyclophosphamide	Not performed	Improved	Independent	
Troxell et al. (2016) <sup>7</sup>	USA	1	F	60	Fever Dysphagia Weight loss	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Negative	0.4	0.73	120–150%	8%	IV Metyl-PD+ Oral-PD+ Methotrexate	Not performed	Improved	Independent	
Ge et al. (2015) <sup>8</sup>	China	1	M	24	Edema Headache	Positive	Negative	Negative	Negative	Negative	Negative	Positive	Positive	7.04	3.7	Postive (not described)	58%	IV Metyl-PD+ Oral-PD+IV cyclophosphamide	Not performed	Not improved	Dependent	Reversible posterior leukoencephalopathy syndrome
Xu et al. (2016) <sup>9</sup>	China	1	F	50	Nocturia Gross hematuria	Negative	Negative	Negative	Positive Hematuria	Negative	Negative	Negative	Negative	0.41	2.6	258 EU/mL	18/31 cellular 89%	IV Metyl-PD+ Oral-PD+ Mycophenolate mofetil	Not performed	Improved	Independent	
This case	Korea	1	F	38	Fever	Negative	Negative	Negative	Positive	Negative	Negative	Negative	Negative	1.4	5.4	187.2 U/mL	63%	IV Metyl-PD+ Oral-PD+IV cyclophosphamide	Not performed	Improved	Independent	

GBM, glomerular basement membrane; HBV, hepatitis B virus; PD, prednisolone; GI, gastrointestinal.

#### REFERENCES

1. Trpkov K, Abdulkareem F, Jim K, Solez K. Recurrence of anti-GBM antibody disease twelve years after transplantation associated with de novo IgA nephropathy. *Clin Nephrol* 1998; 49: 124-8.
2. Cui Z, Zhao MH, Wang SX, Liu G, Zou WZ, Wang HY. Concurrent antiglomerular basement membrane disease and immune complex glomerulonephritis. *Ren Fail* 2006; 28: 7-14.
3. Wechsler E, Yang T, Jordan SC, Vo A, Nast CC. Anti-glomerular basement membrane disease in an HIV-infected patient. *Nat Clin Pract Nephrol* 2008; 4: 167-71.
4. Wang A, Wang Y, Wang G, Zhou Z, Xun Z, Tan X. Mesangial IgA deposits indicate pathogenesis of anti-glomerular basement membrane disease. *Mol Med Rep* 2012; 5: 1212-4.
5. Yamaguchi H, Takizawa H, Ogawa Y, Takada T, Yamaji I, Ura N. A case report of the anti-glomerular basement membrane glomerulonephritis with mesangial IgA deposition. *CEN Case Rep* 2013; 2: 6-10.
6. Gao B, Li M, Xia W, Wen Y, Qu Z. Rapidly progressive glomerulonephritis due to anti-glomerular basement membrane disease accompanied by IgA nephropathy: a case report. *Clin Nephrol* 2014; 81: 138-41.
7. Troxell ML, Houghton DC. Atypical anti-glomerular basement membrane disease. *Clin Kidney J* 2016; 9: 211-21.
8. Ge YT, Liao JL, Liang W, Xiong ZY. Anti-glomerular basement membrane disease combined with IgA nephropathy complicated with reversible posterior leukoencephalopathy syndrome: an unusual case. *Am J Case Rep* 2015; 16: 849-53.
9. Xu D, Wu J, Wu J, et al. Novel therapy for anti-glomerular basement membrane disease with IgA nephropathy: a case report. *Exp Ther Med* 2016; 11: 1889-92.