

## SM3. Baseline characteristics of 243 patients on enrolment into Mwesio IDC-OPD service

Variable	Total	(%)	DM-1	(%)	DM-2	(%)	Other/NA	(%)
<b>Patients (N)</b>	243		62		125		56	
<b>Age</b>								
<18	24	(9.9%)	19	(30.6%)	0	(0.0%)	5	<b>(8.9%)</b>
18-40	74	(30.5%)	39	(62.9%)	20	(16.0%)	15	<b>(26.8%)</b>
41-60	112	(46.1%)	3	(4.8%)	80	(64.0%)	29	<b>(51.8%)</b>
>60	33	(13.6%)	1	(1.6%)	25	(20.0%)	7	<b>(12.5%)</b>
<b>Sex</b>								
Male	134	(55.1%)	34	(54.8%)	70	(56.0%)	30	<b>(53.6%)</b>
Female	108	(44.4%)	28	(45.2%)	55	(44.0%)	25	<b>(44.6%)</b>
No data	1	(0.4%)	0	(0.0%)	0	(0.0%)	1	<b>(1.8%)</b>
<b>Occupation</b>								
Labourer	121	(49.8%)	29	(46.8%)	85	(68.0%)	7	<b>(12.5%)</b>
Office worker	19	(7.8%)	2	(3.2%)	11	(8.8%)	6	<b>(10.7%)</b>
Student	8	(3.3%)	8	(12.9%)	0	(0.0%)	0	<b>(0.0%)</b>
Unable to work	11	(4.5%)	7	(11.3%)	3	(2.4%)	1	<b>(1.8%)</b>
Other	48	(19.8%)	15	(24.2%)	26	(20.8%)	7	<b>(12.5%)</b>
No data	36	(14.8%)	1	(1.6%)	0	(0.0%)	35	<b>(62.5%)</b>
<b>Medical History</b>								
<b>CVD</b>								
Yes	5	(2.1%)	1	(1.6%)	3	(2.4%)	1	<b>(1.8%)</b>
No data	24	(9.9%)	0	(0.0%)	0	(0.0%)	24	<b>(42.9%)</b>
<b>Hypertension</b>								
Yes	20	(8.2%)	0	(0.0%)	18	(14.4%)	2	<b>(3.6%)</b>
No data	25	(10.3%)	0	(0.0%)	1	(0.8%)	24	<b>(42.9%)</b>
<b>Tuberculosis</b>								
Yes	6	(2.5%)	1	(1.6%)	5	(4.0%)	0	<b>(0.0%)</b>
No data	24	(9.9%)	0	(0.0%)	0	(0.0%)	24	<b>(42.9%)</b>
<b>Malnutrition</b>								
Yes	19	(7.8%)	15	(24.2%)	2	(1.6%)	2	<b>(3.6%)</b>
No data	24	(9.9%)	0	(0.0%)	0	(0.0%)	24	<b>(42.9%)</b>
<b>Family Hx of DM</b>								
Yes	35	(14.4%)	6	(9.7%)	23	(18.4%)	6	<b>(10.7%)</b>
No data	24	(9.9%)	0	(0.0%)	0	(0.0%)	24	<b>(42.9%)</b>

<b>Patients &gt;=18 years(N)*</b>	219		43		125		51	
<b>Smoking</b>								
Yes	18	(8.2%)	1	(2.3%)	16	(12.8%)	1	<b>(2.0%)</b>
No data	21	(9.6%)	0	(0.0%)	0	(0.0%)	21	<b>(41.2%)</b>
<b>Alcohol</b>								
Yes	50	(22.8%)	12	(27.9%)	32	(25.6%)	6	<b>(11.8%)</b>
No data	33	(15.1%)	0	(0.0%)	1	(0.8%)	32	<b>(62.7%)</b>
<b>BMI</b>								
<18.5	68	(31.1%)	23	(53.5%)	34	(27.2%)	11	<b>(21.6%)</b>
18.5-24.9	96	(43.8%)	15	(34.9%)	59	(47.2%)	22	<b>(43.1%)</b>
25.0-29.9	41	(18.7%)	5	(11.6%)	27	(21.6%)	9	<b>(17.6%)</b>
>=30	7	(3.2%)	0	(0.0%)	5	(4.0%)	2	<b>(3.9%)</b>
No data	7	(3.2%)	0	(0.0%)	0	(0.0%)	7	(13.7%)

Note: For smoking, alcohol and BMI, denominators included only patients aged 18 years or above.



