

***Schinus terebinthifolia* leaf lectin (SteLL) has anti-infective action and modulates
the response of *Staphylococcus aureus*-infected macrophages**

Isana Maria de Souza Feitosa Lima¹, Adrielle Zagnignan¹, Deivid Martins Santos¹,
Hermerson Sousa Maia¹, Lucas dos Santos Silva¹, Brenda da Silva Cutrim¹, Silvamara
Leite Vieira¹, Clovis Macêdo Bezerra Filho², Eduardo Martins de Sousa¹, Thiago
Henrique Napoleão², Karen Angeliki Krogfelt^{3,4}, Anders Løbner-Olesen⁵, Patrícia
Maria Guedes Paiva², Luís Cláudio Nascimento da Silva^{1*}

¹Programas de Pós-Graduação, Universidade Ceuma, São Luís, Maranhão, Brazil;

²Departamento de Bioquímica, Universidade Federal de Pernambuco, Recife,
Pernambuco, Brazil;

³Department of Viral and Microbial Diagnostics, Statens Serum Institut, Copenhagen,
Denmark;

⁴Department of Science and Environment, Roskilde University, 4000 Roskilde, Denmark;

⁵Department of Biology, Section for Functional Genomics, University of Copenhagen,
Copenhagen, Denmark.

*Corresponding author:


Luís C. N. da Silva

Programa de Pós-Graduação em Biologia Microbiana, Universidade Ceuma

Rua Josué Montello, nº 1, Renascença II, São Luís - MA, CEP 65.075-120

E-mail: luiscn.silva@ceuma.br; luisclaudionsilva@yahoo.com.br

Supplementary Information



Supplementary Figure 1: Effects of SteLL on bactericidal activity of mice peritoneal macrophages towards extracellular and (ExSa) intracellular (IntraSa) *S. aureus* ATCC 6538. M1 macrophages: macrophages treated with LPS + INF- γ . M2 macrophages: macrophages treated with IL-4 + IL-13. Con: untreated cells; Sa: *S. aureus*. (*) Indicates significant differences in relation to *S. aureus*-infected cells without SteLL treatment ($p < 0.05$).