

Additional File 4 - Cerebral Amyloid Angiopathy (CAA) quantification

Brain Region: Hippocampus			
Braak Stage	I-II	III-IV	V-VI
CAA Score^a	Grade 0 (3)	Grade 0 (4)	Grade 0 (4)
	Grade 1 (3)	Grade 1 (0)	Grade 1 (0)
	Grade 2 (1)	Grade 2 (5)	Grade 2 (0)
	Grade 3 (0)	Grade 3 (1)	Grade 3 (2)
	Grade 4 (0)	Grade 4 (0)	Grade 4 (2)
Brain Region: Cortex			
Braak Stage	I-II	III-IV	V-VI
CAA Score	Grade 0 (10)	Grade 0 (7)	Grade 0 (5)
	Grade 1 (1)	Grade 1 (2)	Grade 1 (0)
	Grade 2 (1)	Grade 2 (4)	Grade 2 (3)
	Grade 3 (0)	Grade 3 (0)	Grade 3 (2)
	Grade 4 (0)	Grade 4 (0)	Grade 4 (0)

a: Capillary CAA was absent in the sample set. CAA scores show the prevalence and severity of CAA-laden arteries. CAA Score: Grade 0 = none, Grade 1 = mild, Grade 2 = moderate, Grade 3 = severe, Grade 4 = very severe (Adapted from Arvanitakis Z, Leurgans SE, Wang Z, Wilson RS, Bennett DA, Schneider JA (2011) Cerebral amyloid angiopathy pathology and cognitive domains in older persons. *Annals of Neurology* 69:320–327. doi: 10.1002/ana.22112)