

Table 2. Comparison of estimates from analyses using multiply imputed (MI) datasets and analyses in the complete cases

Outcome	Intervention	Control		Risk difference (95% CI)	Relative risk (95% CI)	P value
Completion of any STI test at 6 weeks	45.3% (n=244)	24.1% (n=284)	Weighted MI	21.2% (12.5, 29.8)	1.88 (1.47, 2.40)	<0.001
			Unweighted MI	21.3% (12.5, 30.2)	1.89 (1.43, 2.52)	<0.001
	44.1% (94/213)	19.6% (44/224)	<i>Weighted Complete Cases</i>	<i>24.9% (16.7, 33.0)</i>	<i>2.26 (1.59, 2.93)</i>	<i><0.001</i>
Completion of chlamydia and gonorrhoea tests at 6 weeks	44.3% (n=244)	24.1% (n=284)	Weighted MI	20.2% (11.7, 28.8)	1.84 (1.44, 2.36)	<0.001
			Unweighted MI	20.4% (11.6, 29.2)	1.85 (1.39, 2.46)	<0.001
	43.2% (92/213)	19.6% (44/224)	<i>Weighted Complete Cases</i>	<i>24.0% (15.9, 32.1)</i>	<i>2.22 (1.56, 2.88)</i>	<i><0.001</i>