

Supplementary Online Content

Dey AK, Gaddipati R, Elnabawi YA, et al. Association between soluble lectinlike oxidized low-density lipoprotein receptor-1 and coronary artery disease in psoriasis. *JAMA Dermatol*. Published online November 20, 2019. doi:10.1001/jamadermatol.2019.3595

eTable. Characteristics of psoriasis patients at baseline

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Characteristics of psoriasis patients at baseline.

Variable	Psoriasis (N=175)
Demographic and clinical characteristics	
Age, years	49.7 ± 12.6 (47.84-51.61)
Males	96 (55)
Hypertension	49 (28)
Hyperlipidemia	78 (45)
Type-2 diabetes	18 (10)
Body mass index, kg/m ²	29.6 ± 6.0 (28.66-30.46)
Current smoker	14 (8)
Statin use	55 (32)
Clinical and lab values	
Total cholesterol, mg/dL	183.4 ± 37.1 (177.86-188.95)
HDL cholesterol, mg/dL	56.0 ± 17.2 (53.58-58.53)
LDL cholesterol, mg/dL	102.9 ± 30.0 (98.49-107.38)
Triglycerides, mg/dL	122.0 ± 74.3 (110.83-133.07)
Framingham risk score	2.0 (1.0-6.0)
High sensitivity c-reactive protein, mg/L	2.0 (0.8-4.8)
Soluble LOX1*, pg/mL	210.3 (110.9-336.2)
Psoriasis severity	
Psoriasis area severity index score	5.6 (3.0-9.2)
Biologic treatment	37 (21)
Coronary artery characterization	
Non-calcified coronary burden, mm ² (x100)	1.10 ± 0.45 (1.05-1.14)
Dense-calcified coronary burden, mm ² (x100)	0.019 (0.006-0.063)

Values reported in the table as Mean ± SD (95% CI) or Median (IQR) for continuous data and N (%) for categorical data; *LOX1: Lectin-like oxidized low-density lipoprotein receptor-1.