

S1 Table. Average body weight of binge drinking C57BL/6 mice intranasally infected with different bacterial doses. Mice were administered alcohol (4.4 g/kg) and 30 min later intranasally infected with various bacterial doses. Mice were weighed before infection and 24 h post infection. PBS (control) indicates mice that were not infected or administered alcohol. Alcohol (control) indicates mice that were administered alcohol and not infected. (*) indicates statistical comparison between pre-infection and post-infection (24 h) per group by Student's unpaired *t*-test, *, $p \leq 0.05$, **, $p \leq 0.01$.

Body Weight (g)			
<i>B. thailandensis</i> (CFUs)	Pre-Infection	Post-Infection (24 h)	Decrease (%)
PBS (control)	19.8 ± 0.012 (n = 6)	19.8 ± 0.015 (n = 6)	0.0
Alcohol (control)	19.7 ± 0.121 (n = 6)	19.8 ± 0.132 (n = 6)	0.0
3 x 10 ⁵	19.3 ± 0.327 (n = 6)	17.6 ± 0.230 (n = 6)	8.8 **
5 x 10 ⁴	19.6 ± 0.528 (n = 6)	17.9 ± 0.478 (n = 6)	8.7 *
8 x 10 ³	19.7 ± 0.448 (n = 6)	17.6 ± 0.412 (n = 6)	10.7 **
500	19.5 ± 0.351 (n = 6)	17.1 ± 0.444 (n = 6)	12.3 **

Values are means \pm SEM; n, number of mice