animals

Supplementary Materials: Improved Post-Thaw Quality of Canine Semen after Treatment with Exosomes from Conditioned Medium of Adipose-Derived Mesenchymal Stem Cells

Ahmad Yar Qamar [1], Xun Fang [1], Min Jung Kim [2,*] and Jongki Cho [1,*]

[1] College of Veterinary Medicine, Chungnam National University, Daejeon 34134, Korea; drahmadqamar@gmail.com (A.Y.Q.); fx2442@gmail.com (X.F.)

[2] Department of Theriogenology and Biotechnology, College of Veterinary Medicine, Seoul National University, Seoul 08826, Korea

[*] Correspondence: tinia19@snu.ac.kr (M.J.K.); cjki@cnu.ac.kr (J.C.); Tel.: +82-42-821-6788 (J.C.) Fax: +82-42-821-8903 (J.C.)

Table S1. Motility characteristics of dog sperm cooled at 4 °C for 8–10 h.

Groups	Progressive motility (%)	VCL (µm/sec)	VSL (µm/sec)	VAP (µm/sec)	BCF (Hz)
Control	40.5 ± 0.4 [a,b,c]	98.5 ± 0.8 [b]	31.1 ± 0.6	50.9 ± 0.6	5.5 ± 0.2
25 µg/ mL	39.5 ± 0.3 [b]	96.0 ± 1.1 [b]	29.9 ± 0.6	49.0 ± 0.6	5.4 ± 0.2
50 µg/ mL	41.6 ± 0.7 [a]	103.1 ± 1.1 [a]	31.4 ± 0.3	52.7 ± 0.8	6.1 ± 0.3
100 µg/ mL	38.6 ± 0.4 [c]	96.8 ± 1.1 [b]	29.8 ± 0.5	50.2 ± 0.6	5.4 ± 0.6

Values with different superscripts letters in a column differ significantly ($p < 0.05$, $n = 4$). VCL, Curvilinear velocity; VSL, Straight-line velocity; VAP, Average path velocity; BCF, Beat cross frequency.

Table S2. Effect of exosomal treatment on the progressive motility and velocities of post-thaw dog semen.

Group	Progressive motility (%)	VCL (µm/sec)	VSL (µm/sec)	VAP (µm/sec)	BCF (Hz)
Control	20.9 ± 0.6 [b]	50.1 ± 0.9 [b]	17.7 ± 0.7 [b]	27.7 ± 0.3 [b]	2.5 ± 0.0 [b]
Treatment (50 µg/mL)	25.3 ± 0.8 [a]	62.6 ± 1.1 [a]	20.6 ± 0.5 [a]	32.1 ± 0.5 [a]	3.2 ± 0.1 [a]

Values with different superscripts letters in a column differ significantly ($p < 0.05$, $n = 4$). VCL, Curvilinear velocity; VSL, Straight-line velocity; VAP, Average path velocity; BCF, Beat cross frequency.