

SUPPLEMENTARY TABLE S1. PRIMERS USED FOR QUANTITATIVE REVERSE TRANSCRIPTION PCR

<i>Target</i>	<i>Forward primer (5'-3')</i>	<i>Reverse primer (5'-3')</i>
GAPDH	CATCACCATCTTCCAGGAGCGA	GTGGTTCACACCCATCACAAACA
DES	CAGTATGAGACCATCGCGGCTA	GCAGGTGTAGGACTGGATCTGG
MSTN	ACAGCCTGAATCCAACCTTAGGCA	CTCATCGCAGTCAAGCCCAAAG
MYF5	ATGTAACAGCCCTGTCTGGTCC	GAGAGAGGGAAGCTGTGTCCTG
MYF6	TTCTTGAGGGTGCGGATTCCT	GGGAGTTGCGTTCCTCTGAAG
MYH1	TGCAGAGCAAGATTGAGGACGA	CCAGCTTCTTCCAGCCTCTCG
MYOD	TCCAACCTGCTCTGATGGCATGA	CCTCACTGTAGTAGGCGGTGTC
MYOG	CAGTGAATGCAACTCCCACAGC	CTCAGTTGGGCATGGTTTCGTC
TNN1	GGTGGATGAGGAGCGCTATGAT	AGATCCATGGACACCTTGTGCT