

## Supplementary Data

SUPPLEMENTARY TABLE S1. PREOPERATIVE OSTEOPOROSIS AND CARDIOVASCULAR RISK CATEGORIES

	<i>High risk</i>	<i>Intermediate risk</i>	<i>Low risk</i>
Atrial fibrillation risk	Clinical heart disease Age >75 years Comorbidities: <ul style="list-style-type: none"> <li>• Valvular heart disease</li> <li>• TIA or stroke</li> <li>• Vascular disease</li> <li>• Renal failure</li> </ul>	Age >60 years HTN Symptomatic thyrotoxicosis Cardiovascular risk factors: <ul style="list-style-type: none"> <li>• Diabetes</li> <li>• HTN</li> <li>• Hyperlipidemia</li> <li>• Cigarette smoking</li> </ul>	Age <60 years No comorbidities
Osteoporosis risk	Postmenopausal (age >50 years or oophorectomy) Prior radiotherapy	Perimenopausal women (age 40–50 years) Osteopenia Osteoporosis risk factors: <ul style="list-style-type: none"> <li>• Cigarette smoking</li> <li>• Gastric bypass</li> <li>• Anorexia</li> <li>• Family history</li> </ul>	Age <40 years No comorbidities No osteoporosis risk factors