

SUPPLEMENTARY TABLE S2. ASSOCIATIONS BETWEEN UIC GROUPS AND ALL-CAUSE, CARDIOVASCULAR, AND CANCER MORTALITY IN THE NHANES III AFTER FURTHER ADJUSTMENT FOR TOTAL SODIUM INTAKE AND FAT/CALORIE RATIO<sup>a,b</sup>

	<i>Very low UIC</i>	<i>Low UIC</i>	<i>Normal UIC (reference)</i>	<i>High UIC</i>	<i>Very high UIC</i>
All-cause mortality	0.94 [0.77–1.14]	0.95 [0.84–1.09]	1	0.92 [0.68–1.23]	1.21 [1.05–1.39]
Cardiovascular mortality	0.84 [0.63–1.12]	0.86 [0.68–1.07]	1	0.72 [0.40–1.29]	1.11 [0.84–1.48]
Cancer mortality	1.38 [0.96–2.00]	1.15 [0.88–1.51]	1	1.14 [0.71–1.85]	1.48 [0.99–2.20]

<sup>a</sup>The reference is the normal UIC group.

<sup>b</sup>HR adjusted for age, sex, race/ethnicity, education status, active smoking, diabetes, hypertension, hypercholesterolemia, previous cardiovascular disease, previous cancer, BMI, eGFR, total sodium intake (quintiles), and fat/calorie ratio (quintiles).