

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Socioeconomic and behavioral determinants of overweight/obesity in Botswana: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029570
Article Type:	Research
Date Submitted by the Author:	31-Jan-2019
Complete List of Authors:	Keetile, Mpho; University of Botswana, Population Studies Navaneetham, Kannan; University of Botswana, Population Studies Letamo, Gobopamang; University of Botswana, Population Studies Bainame, Kenabetsho; University of Botswana, Population Studies Rakgoasi, Serai; University of Botswana, Population Studies Gabaitiri, Lesego; University of Botswana, Statistics Masupe, Tiny; University of Botswana, School of Medicine Molebatsi, Robert; University of Botswana, Sociology
Keywords:	Overweight, Obesity, Socioeconomic, Behavioural, Botswana

SCHOLARONE™
Manuscripts

1
2
3
4
5
6 **Socioeconomic and behavioural determinants of overweight/obesity in**
7 **Botswana: A cross-sectional study**
8
9

10 Mpho Keetile¹, Kannan Navaneetham¹, Gobopamang Letamo¹, Kenabetsho
11 Bainame¹, Serai Daniel Rakgoasi¹, Lesego Gabaitiri², Tiny Masupe³, Robert
12 Molebatsi⁴.
13
14

15
16 **Author Affiliations**
17

18 ¹Department of Population Studies, University of Botswana, Gaborone, Botswana

19 ²Department of Statistics, University of Botswana, Gaborone, Botswana

20 ³School of Medicine, University of Botswana, Gaborone, Botswana

21 ⁴Department of Sociology, University of Botswana, Gaborone
22
23
24
25

26 **Corresponding author**
27

28 Mpho Keetile, Department of Population Studies, Private Bag 00705, University of
29 Botswana, Gaborone, Botswana. email: mphokeet@yahoo.com. Tel: +26771231375
30
31
32
33
34
35
36

37 **Word count: 3507**
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective

To undertake a comprehensive assessment of socioeconomic and behavioural determinants of overweight/obesity in Botswana.

Design

The study adopted a cross-sectional design and was conducted in selected urban and rural areas of Botswana. Rural areas constituted 15 urban villages and 15 rural areas, while urban areas constituted 1 city and 2 towns. Body Mass Index (BMI) was categorized into; underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), normal weight ($18.5 \leq \text{BMI} < 25 \text{ kg/m}^2$), overweight ($25 \leq \text{BMI} < 30 \text{ kg/m}^2$) and obese ($\text{BMI} \geq 30 \text{ kg/m}^2$). We used binary logistic regression analysis to estimate determinants of overweight/obesity because overweight and obese were coded as: being overweight and obese ($\text{BMI} \geq 25$) = 1; not overweight and obese = 0 ($\text{BMI} < 25$). Analysis of questionnaire responses used logistic regression models.

Setting

Botswana

Participants

1178 males and females aged 15 years and above from the survey on 'Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course', conducted in March 2016.

Outcome Measures

Adjusted ORs for socioeconomic and behavioural determinants of overweight/obesity.

Results

Prevalence of overweight/obesity in the study population was estimated at 41%. The adjusted odds of overweight/obesity were highest among women (AOR=2.74, 95% C.I. =1.92-3.90), in ages 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), individuals with secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary or higher education (AOR=1.99, 95% C.I. =1.16-3.38), smokers (AOR=2.16, 95% C.I. = 1.22-3.83) and poor physically active people (AOR=1.46, 95% C.I. =1.03-3.24).

Conclusion

Being female, elderly, of high education levels, a smoker and having poor physical activity were identified as key factors for prevalence of overweight/obesity. These findings suggest the need for broad based strategies encouraging physical activity among different socioeconomic groups.

Key words

Overweight, Obesity, socioeconomic, behavioral, Botswana

Strengths and Limitations of the study

- To our knowledge this is the first cross-sectional study in Botswana to comprehensively assess both the socioeconomic and behavioral determinants of overweight/obesity among adults.
- Data were collected from a large and randomly selected representative population. The data contained information on potential confounding factors, with a low proportion of missing information.
- A limitation for this study was that since a cross sectional design was employed, data on each participant was recorded only once making it difficult to infer the temporal association between an explanatory and an outcome variable.
- Only an association, and not causation, can be inferred from this study.

Introduction

The disease profile in Botswana is changing at an alarming rate, with most deaths and disability in the foreseeable future likely to be accounted for by the ominous epidemics of non-communicable diseases (NCDs) such as heart disease, stroke, cancer and other chronic diseases. Children, adults and the elderly are all vulnerable to the risk factors that contribute to NCDs especially overweight/obesity, due to unhealthy diets, low physical activity, exposure to tobacco smoke and/or the effects of the harmful use of alcohol¹⁻³. The key drivers of NCDs are ageing, rapid unplanned urbanisation, and the globalisation of unhealthy lifestyles. For example, globalization of unhealthy lifestyles like unhealthy diets may show up in individuals as raised blood pressure, increased blood glucose, elevated blood lipids, and overweight/ obesity¹.

Several studies have recently shown that many low income and middle-income countries (LMICs) are experiencing an increase in the prevalence of overweight/obesity⁴⁻⁶. This prevalence is predicted to continue to increase in the future. Much of the increase in prevalence levels for overweight/obesity is expected in countries such as India, some parts of South-East Asia, China, most of South America and some parts of Sub Saharan Africa⁷⁻⁸. Until recently, SSA was minimally affected by the obesity/overweight epidemic due to under-nutrition and a major burden of HIV and tuberculosis⁹. Meanwhile research on overweight/obesity has started to gain significance in sub Saharan Africa¹⁰⁻¹⁴ due to its increasing magnitude.

It has been shown by previous studies that socioeconomic and behavioural factors have a significant association with overweight/obesity¹⁵⁻¹⁶. These studies have

1
2
3 shown that socioeconomic factors such as sex, age, education level, work status,
4 place of residence and wealth status are significant determinants of
5 overweight/obesity. It has also been found that women are more likely to be
6 overweight/obese than men, while the odds of being overweight/obesity has been
7 seen to increase with age¹⁷⁻¹⁹. Other studies have shown that being employed in a
8 high income job, residing in urban areas, having high education and wealth status is
9 significantly associated with overweight/obesity²⁰.

10
11
12
13
14
15
16
17
18
19
20
21 There is also available evidence which suggest significant correlation between
22 behavioural factors and overweight/obesity. For example, there is significant
23 evidence of a positive association between physical inactivity and the prevalence of
24 overweight/obesity²¹. It has also been suggested that the relation between physical
25 activity and health outcomes such as overweight/obesity may be moderated by a
26 number of lifestyle factors, especially sedentary lifestyle^{15, 20, 21}. In many developing
27 countries, there has been a substantial shift from jobs with high-energy expenditure
28 such as farming, mining, and forestry, and employment is inclined towards the more-
29 sedentary sectors of manufacturing, services and office based-work²². Moreover,
30 there have been changes in lifestyle behaviours condoning alcohol consumption,
31 poor diets and smoking. These are also key risk factors of overweight/obesity²³⁻²⁴.

32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49 Despite evidence indicating increasing burden of overweight/obesity in Botswana,
50 there is little evidence of socioeconomic and behavioural determinants of
51 overweight/obesity. To the best of our knowledge this is the first cross sectional
52 study which takes into account both socioeconomic and behavioural factors for
53 understanding the heterogeneity of overweight/obesity in Botswana. The reduction
54
55
56
57
58
59
60

1
2
3 of overweight/obesity prevalence should be a key goal of health-care policy because
4 of its association with many NCDs. The main objective of this study was to assess
5 socioeconomic and behavioural determinants associated with overweight/obesity
6 prevalence in Botswana.
7
8
9
10
11
12
13
14

15 **Methods**

16
17
18 The study used cross sectional design by selecting respondents in three cities and
19 towns, fifteen urban villages and fifteen rural areas across Botswana using a
20 multistage probability sampling technique. The survey was carried out in March
21 2016. The survey collected self-reported data on several NCDs as classified by the
22 WHO classification of diseases and their risk factors. Information collected from
23 these respondents included social and behavioural characteristics, and
24 anthropometric measurements (height and weight). Body Mass Index (BMI) was
25 used to classify overweight/obesity in adults. BMI was derived from weight and
26 height: $\text{weight (kg)} / (\text{height (m)} \times \text{height (m)})$. Participants removed their shoes and
27 heavy outer clothing before weight and height were measured. Weight was
28 measured, to the nearest 0.1kg, while height was measured in metres. BMI was
29 categorized into; underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), normal weight ($18.5 \leq \text{BMI} < 25$
30 kg/m^2), overweight ($25 \leq \text{BMI} < 30 \text{ kg/m}^2$) and obese ($\text{BMI} \geq 30 \text{ kg/m}^2$). Overweight and
31 obese were used to create a binary outcome variable which was coded as: being
32 overweight/obese ($\text{BMI} \geq 25$) =1; not overweight/obese =0 ($\text{BMI} < 25$).
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Patient and Public involvement

Since this was a cross-sectional study respondents were drawn from the general population rather than from healthcare settings, thereby providing access to all eligible people aged 15 years and above. As part of the study protocol all respondents were requested to sign a consent form and were told that participation in the study was voluntary. It was also highlighted to respondents that only the general findings of the study shall be shared to insure confidentiality.

Measurement of variables

Information for NCD risk factors was collected through self-reports. For tobacco smoking - the survey question asked respondent: 'Do you currently smoke any tobacco products such as cigarettes, cigars or pipes? This variable was coded such that yes=1 and no=0. Alcohol consumption was derived from the question: 'Have you ever consumed alcohol in the past 30 days?' This variable was used because in Botswana, it has been observed that alcohol consumers are heavy drinkers²⁵. The resultant variable was yes=1 and no=0. For poor physical activity, four key features of the quality of physical activity measures (e.g. activity type, intensity, frequency, and duration) are usually considered when choosing one for a research study. For this study the following two questions asking respondents about moderate and vigorous intensity physical activity were used. Activity at work asked the following questions: "Does your work involve vigorous-intensity activity that causes large increases in breathing or heart rate like [examples] for at least 10 minutes continuously?" and "Does your work involve moderate-intensity activity that causes small increases in breathing or heart rate such as brisk walking for at least 10 minutes continuously?". Different types of responses to the two levels of physical

1
2
3 activities have been grouped together and given a value based on the intensity of
4 the activity. The resultant variable was coded such that those who responded
5 'yes'=1 (physically active) and those who said 'no' were coded=0 (or poor physically
6 active).
7
8
9
10
11
12
13
14
15

16 For the measurement of insufficient or non-consumption of fruits and vegetables the
17 following question was asked to assess fruit and vegetable consumption in the
18 population; "How many servings of fruits do you eat on one of those days (on a
19 typical day)?" and "How many servings of vegetables do you eat on one of those
20 days (On a typical day)". The resultant variable was coded such that people who
21 reported to have taken more than 5 servings of fruit and vegetables were given a
22 code, sufficient intake=0 and those who reported to have taken less than 5 servings
23 were given a code insufficient intake=1. This was done based on the general
24 recommendation by the WHO panel on diet, nutrition and chronic disease prevention
25 that considers poor fruit/vegetables intake as having less than 5 servings of fruits
26 and vegetables in a day.
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 A wealth index (WI) was constructed to be a measure of wealth status. WI is a
42 composite measure of, typically, indicators of ownership of consumer durables,
43 housing characteristics, and access to public services. Information on a range of
44 durable assets was collected during the survey (e. g. car, refrigerator, television,),
45 housing characteristics (e. g. material of dwelling floor and roof, main cooking fuel),
46 access to basic services (e. g. electricity supply, source of drinking water, sanitation
47 facilities) and ownership of livestock (e.g. cattle, goats, sheep, horses, chickens).
48 Further to collection of information on durable assets, information on land and
49 livestock ownership was collected. Principal component analysis was employed to
50
51
52
53
54
55
56
57
58
59
60

1
2
3 derive the wealth index variable, which had five categories from the 1st to the 5th
4
5 quintile (poorest to richest).
6
7

8 **Control variables**

9
10
11 Age and place of residence were used as control variables. These variables were
12
13 conceptualized to have an association with the outcome variable. Therefore to hold
14
15 their likely association with the outcome variable, they are included in the combined
16
17 effects model, so that the association between the independent variables becomes
18
19 isolated and discernible.
20
21
22

23 **Statistical analysis**

24
25
26 Data analysis involved descriptive and multivariable analysis. Descriptive analysis
27
28 provided some basis for further exploration of key determinants of overweight and
29
30 obesity in Botswana. In the binary logistic regressions, outcome variables were
31
32 coded as '1' if respondent was overweight/obese and '0' if they were not
33
34 overweight/obese. Results of logistic regression analysis were presented together
35
36 with adjusted odds ratios (AOR), 95% confidence intervals (CI) and significance
37
38 levels. Data analysis was done using SPSS version 25 program. In order to control
39
40 for cluster effects complex samples module in SPSS has been used since the NCDs
41
42 survey used multistage sampling technique.
43
44
45
46

47 **Results**

48 ***Sample characteristics***

49
50
51
52
53 Table 1 presents results on the sample description for the respondents. The results
54
55 show that a total of 1,178 respondents between the ages of 15 years and above
56
57 were successfully interviewed. The sample constituted a high proportion of females
58
59
60

1
2
3 (69.1%) than males (30.9%). The sample age distribution suggests a relatively
4 young study population, with over half (59%) of the sample being less than 39 years
5 of age, and almost three quarters (73.5%) being less than fifty years of age. As age
6 increases, the sample sex distribution becomes more skewed in favour of females,
7 from just fewer than 6 in every ten among those in the 20-24 years age group, to
8 over 7 and 8 out of every ten among respondents over 40 years of age.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1: Sample Characteristics

Variable	%	N
Sex		
Male	30.9	364
Female	69.1	813
Total		1177
Age in years		
<24	26.4	270
25– 34	29.5	302
35 – 44	19.2	196
45– 54	12.7	130
55 – 64	7.3	75
65+ years	4.9	50
Total	100	1023
Locality Type		
Cities/Towns	30.2	355
Urban Villages	45.4	534
Rural Settlements	24.5	288
Total	100.0	1177
Marital Status		
Never Married	73.8	864
Currently married	17	199
Formerly married	9.2	108
Total	100.0	1171
Highest Level of Education Attained		
Primary or Less	35.5	410
Junior Secondary	27.2	314
Senior Secondary	17.3	200
Tertiary & Over	19.9	230
Total	100.0	1154
Work Status in past 12 months		
Public Sector	10.5	122
Private Sector	15.7	182
Self Employed	11.2	130
Not Employed	37.5	436
Homemaker-Student	18.8	218
Retired-Other	6.4	74
Total	100.0	1162
Wealth status		
Lowest	19.9	234
Second	20.1	237
Middle	19.9	235
Fourth	20.1	237
Highest	19.9	235
Total	100.0	1178

1
2
3
4
5 More than two fifths (45.4%) of the population resided in urban villages; just under a
6
7 third (30.2%) resided in cities and towns while a quarter (24.5%) resided in rural
8
9 areas and settlements. Almost three quarters (73.8%) of respondents were never
10
11 married; over a third (35.5%) had primary education or less; over a quarter (27.2%)
12
13 had junior secondary education while just under a fifth had senior secondary
14
15 education (17.3%) and tertiary education and over (19.9%). Close to two fifth
16
17 (37.5%) of respondents were not employed; while over a quarter were employed in
18
19 either the public (10.5%) or private sector (15.7%). Just over one in every ten
20
21 (11.2%) were self-employed, while close to a fifth (18.8%) were either home makers
22
23 or students; while only 6.4 per cent were retired from work.
24
25
26
27
28
29

30 Table 2 below shows behavioural characteristics of the study population. It shows
31
32 that prevalence of alcohol consumption was estimated at 17% while tobacco
33
34 smoking was estimated at 11.5%. However, since prevalence rate for smoking
35
36 represents active smokers and not passive (second-hand smokers) it may be a
37
38 misrepresentation of the effects of tobacco smoking in the general population. Poor
39
40 physical activity was also high in the study population with 48.9% of respondents
41
42 indicating that they were physically inactive. Poor fruit and vegetable (FV)
43
44 consumption calculated based on the general recommendation by the WHO panel
45
46 on diet, nutrition and chronic disease prevention that considers poor fruit/vegetables
47
48 intake as having less than 5 servings of fruits and vegetables in a week was
49
50 estimated at 82.5%.
51
52
53
54
55
56
57
58
59
60

Table 2: Behavioural characteristics of the study population

Variable	%	N
Smoking?		
Yes	11.5	136
No	88.5	1042
Total		1178
Alcohol consumption		
Yes	17.3	204
No	82.7	974
Total		1178
Poor physical activity		
Yes	48.9	576
No	51.1	602
Total		1147
Fruit and vegetable intake?		
Poor fruit/vegetable intake	82.5	1045
Sufficient intake	17.5	133
Total		1178

Prevalence of overweight/obesity

Estimated prevalence of overweight/obesity in the sampled study population was 41.3% (Table 3). There were significant gender differences in prevalence of overweight/obesity. It was found that women than men (48.6% in women, 25.5% in men) were found to be overweight/obese. It was observed that individuals residing in rural areas (46.8%), public sector employee (55.2%) and those with lower education (45%) showed high prevalence rates for overweight/obesity. However, it was found that wealth status was not significantly linked with overweight/obesity. Among

1
2
3 behavioural factors this study offers evidence that non-smoker (42.9%) showed high
4 prevalence rates of overweight/obesity. Meanwhile alcohol consumption, poor
5 physical activity and poor FV consumption did not show any significant association
6 with overweight and obesity.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 3: Prevalence of overweight/obesity by socioeconomic and behavioural characteristics of the study population

Variable	%	N	P-value
Sex			
Male	25.5	364	0.000**
Female	48.6	813	
Age			0.000**
≤24	21.8	270	
25-34	36.8	302	
35-44	53	196	
45-54	54.5	130	
55-64	61.8	75	
65+	43.8	50	
Marital status			0.000**
Never-married	35.5	864	
Currently-married	62.2	199	
Formerly-married	54.1	108	
Education			0.019**
Primary or less	45.4	410	
Secondary	36.7	514	
Tertiary or higher	44.6	230	
Residence			0.008**
Cities and towns	37.9	355	
Urban villages	41	534	
Rural villages	46.8	288	
Work status			0.000**
Public sector	52.2	122	
Private sector	36.8	182	
self-employed	55	130	
Not employed	45.8	436	
Home-maker/student	29.4	218	
Retired/other	40.3	74	
Wealth status			0.765
Lowest	38.8	234	
second	43.2	237	
middle	41.1	235	
Fourth	39.7	237	
Highest	44.1	235	
Alcohol consumption			0.445
No	33.3	766	
yes	37.7	412	
Smoking			0.004**
Yes	29.7	136	

No	42.9	1042	
Poor physical activity			0.547
Yes	41.5	376	
No	39.6	771	
Poor fruit/vegetable consumption			0.832
No	41.9	133	
yes	43	1045	
Overall	41.3	1178	

Note: **Statistically significant at $p \leq 0.05$.

Determinants of overweight/obesity

Table 4 indicates results of the logistic regression model for overweight/obesity with socioeconomic and behavioural factors. One unadjusted and two adjusted models were run. Model I assessed the association between overweight/obesity and each of the socioeconomic variables and behavioural factors, Model II assessed the association between overweight/obesity adjusting for socioeconomic factors, while Model III assess the association between overweight/obesity adjusting for both socioeconomic and behavioural factors. It was observed that gender was a significant correlate of overweight/obesity when controlling for both socioeconomic and behavioural factors. Women were observed to be 2 times (AOR=2.74, 95% C.I. =1.92-3.90) more likely to be overweight/obese than men.

It was also found that the odds of being overweight/obese increased with age after controlling for socioeconomic and behavioural factors and were highest among individuals aged 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), but declined at ages 65+ years (AOR=2.88, 95% C.I. = 1.21-6.86). Education was also a found to be a significant factor associated with overweight/obesity in the study population

1
2
3 after controlling for socioeconomic covariates. It was found that individuals who had
4 secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary or higher (AOR=1.99, 95%
5 C.I. =1.16-3.38) education were more likely to be overweight/obese than individuals
6 with primary or less education. Meanwhile, there was no significant relationship
7 observed between work status, wealth status and overweight/obesity in the general
8 study population.
9
10
11
12
13
14
15
16
17
18
19

20 It was found that in the crude model for the association between a risk factor and
21 overweight/obesity only smoking (OR=0.56, C.I. =0.37-0.83) and alcohol
22 consumption (OR=0.66, C.I. =0.47-0.91) were found to be negatively associated with
23 overweight/obesity. Meanwhile, poor physical activity and fruit and vegetable
24 consumption did not show any significant association with overweight/obesity.
25 Adjustment for other socioeconomic variables and risk factors suggested that
26 variations in smoking accounted for part of the socioeconomic variation in
27 overweight/obesity. For instance it was noted the odds of overweight/obesity
28 increased two times (AOR=2.16, 95% C.I. = 1.22-3.83) among smokers than non-
29 smokers. Overweight/obesity was also significantly associated with poor physical
30 activity (AOR=1.46, 95% C.I. =1.03-3.24) after adjustment for both socioeconomic
31 and behavioural factors.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 4: Odds ratios for the association between overweight/obesity, socioeconomic and behavioural variables in the study population, NCD study 2016

Variable	Model I		Model II		Model III	
	Exp	C.I.	Exp B	C.I.	Exp B	C.I.
Sex						
Male	1.00		1.00		1.00	
Female	2.75**	(2.08-3.64)	3.01**	(2.12-4.26)	2.74**	(1.92-3.90)
Age						
≤24	1.00		1.00		1.00	
25-34	2.08**	(1.42-3.03)	1.77**	(1.12-2.77)	1.83**	(1.17-2.88)
35-44	4.02**	(2.67-6.08)	3.70**	(2.23-6.12)	3.87**	(2.34-6.42)
45-54	4.28**	(2.70-6.78)	3.04**	(1.66-5.55)	3.26**	(1.78-5.98)
55-64	5.78**	(3.26-10.2)	4.91**	(2.35-10.2)	5.53**	(2.62-11.6)
65+	2.78**	(1.46-5.28)	2.74**	(1.15-6.50)	2.88**	(1.21-6.86)
Marital status						
Never-married	0.46**	(0.30-0.71)	0.68	(0.34-1.34)	0.66	(0.33-1.32)
Currently-married	1.39	(0.84-2.29)	1.34	(0.66-2.71)	1.28	(0.63-2.61)
Formerly-married	1.00		1.00		1.00	
Education						
Primary or less	1.00		1.00		1.00	
Secondary	0.69**	(0.53-0.91)	0.49**	(0.29-0.83)	1.70**	(1.11-2.61)
Tertiary or higher	0.96	(0.69-1.35)	0.87	(0.58-1.32)	1.99**	(1.16-3.38)
Residence						
Cities and towns	1.00		1.00		1.00	
Urban villages	1.13	(0.86-1.50)	1.06	(0.74-1.50)	1.07	(0.75-1.52)
Rural villages	1.44**	(1.04-1.99)	1.37	(0.88-2.12)	1.37	(0.88-2.12)
Work status						
Public sector	1.61	(0.89-2.93)	1.30	(0.64-2.65)	1.22	(0.59-2.50)
Private sector	0.86	(0.49-1.51)	1.20	(0.61-2.36)	1.19	(0.60-2.35)
self-employed	1.18	(0.65-2.13)	1.24	(0.61-2.51)	1.29	(0.63-2.62)
Not employed	1.25	(0.75-2.08)	1.30	(0.70-2.40)	1.30	(0.70-2.41)
Home-maker/student	0.61	(0.35-1.07)	0.96	(0.48-1.92)	0.96	(0.48-1.92)
Retired/other	1.00		1.00		1.00	
Wealth status						
Lowest	0.80	(0.55-1.17)	0.60	(0.34-1.05)	1.03	(0.64-1.68)
second	0.96	(0.66-1.40)	0.67	(0.40-1.11)	1.12	(0.68-1.85)
middle	0.88	(0.60-1.28)	0.73	(0.45-1.19)	1.20	(0.71-2.01)
Fourth	0.83	(0.57-1.21)	0.79	(0.50-1.23)	1.53	(0.87-2.69)
Highest	1.00		1.00		1.00	

Smoking		N.I	
Yes	0.56* (0.37-0.83)		2.16** (1.22-3.83)
No	1.00		1.00
Poor physical activity		N.I	
Yes	1.08 (0.83-1.40)		1.46** (1.03-3.24)
No	1.00		1.00
Poor fruit/vegetable consumption		N.I	
Yes	0.82 (0.55-1.22)		0.80 (0.28-2.24)
no	1.00		1.00
Alcohol consumption		N.I	
Yes	0.66* (0.47-0.91)		1.23 (0.53-2.83)
No	1.00		1.00

N.I- not included

Model I: Crude model; Model II: socioeconomic variables included; Model III: socioeconomic + behavioural characteristics included. **statistically significant at $p \leq 0.05$.

Discussion

Estimated prevalence of overweight/obesity in the study population was 41.3%. This high prevalence is similar to the growing prevalence of overweight/obesity sweeping across Southern Africa, with national prevalence rates ranging between 30 and 60% among populations over the age of 15 years in most countries^{10,26-31}. The high prevalence of overweight/obesity observed in Botswana may be attributed to: dietary shift away from high-fibre, low-calorie diets rich in fruits and vegetables towards refined, energy-dense foods high in fat, calories, sweeteners and salt²⁷ and to low physical activity due to transition from agrarian to a sedentary industrial society³².

The existence of socioeconomic differences in overweight/obesity prevalence is a well-established finding, and has been previously confirmed³³⁻³⁴. This study adds to those findings. In particular, this analysis suggests that more women than men were overweight/obese. For example, women were found to be two times more likely to be overweight/obese than men. Literature yields mounting evidence on the gender differential for overweight/obesity across and within countries³⁵⁻³⁷. While in some developed countries such as the US and the UK men have been observed to be more overweight than women³⁸, in developing countries women are noted to be overweight than men^{35,39}. However, although gender differences for overweight and obesity vary greatly within and between countries, overall women are likely to be overweight and obese than men. Similarly in this data it was noted that gender disparities in overweight and obesity were exacerbated among women, like elsewhere in developing countries, especially in the Middle East and SSA³⁵.

Gender differences in overweight/obesity observed between developed and developing countries with weight gain high among men in developed countries and

1
2
3 high among women in developing countries indicate that gender disparities in excess
4 weight gain are explained by myriad sociocultural dynamics. For instance,
5 acculturation, through complex sociocultural pathways, affects weight gain among
6 both men and women. Moreover, the nutrition transition taking place in many
7 developing countries has also affected excess weight gain among both genders, but
8 has had an even greater impact on the physical activity levels of women⁴⁰.
9
10 Furthermore, in some countries, cultural values favour larger body size among
11 women or men as a sign of fertility, healthfulness, or prosperity especially in SSA⁴¹.
12 Other studies suggest that contextual factors drive gender differences in food
13 consumption, and women often report consuming healthier foods, yet may consume
14 more sugar-laden foods, than men⁴².

15
16 Similarly, in Botswana gender differences in overweight/obesity can be explained by
17 a wide array of sociocultural dynamics. Firstly, although women are more likely to
18 report eating or wanting to eat “healthier” foods, they seem to prefer and consume
19 more foods high in added sugars than men including energy-dense processed foods
20 such as cookies, chocolate, and ice cream. Secondly, overweight among women
21 may also be attributed to comparatively low levels of physical activity⁴³. Moreover,
22 like in other African settings, being overweight/obese among Botswana women is
23 considered as a sign of social status, fertility, good health and prosperity³⁹.

24
25 The odds of being overweight/obese increased with age after controlling for
26 socioeconomic and behavioural factors. This is quite indicative that age is a
27 significant correlate of overweight/obesity independent of other covariates. It was
28 found that odds of being overweight and obese increased with age and were highest
29 among individuals aged 55-64 years but declined at ages 65+ years. This finding
30 corroborates data from large population studies⁴⁴⁻⁴⁶ which shows that mean body

1
2
3 weight and BMI gradually increases during most of adult life and reach peak values
4
5 at 50–59 years of age in both men and women and after the age of 60 years, mean
6
7 body weight and BMI tend to decrease.
8
9

10 Clinically it has been found that aging is associated with considerable body changes
11
12 in body composition⁴⁷. It has been suggested that after 20–30 years of age, fat-free
13
14 mass (FFM) progressively decreases, whereas fat mass increases⁴⁸. Aging is also
15
16 associated with a redistribution of both body fat and FFM whereby there is a greater
17
18 relative increase in intra-abdominal fat than in subcutaneous or total body fat, and
19
20 there is a greater relative decrease in peripheral than in central FFM because of the
21
22 loss of skeletal muscle⁴⁹. This possibly explains the observed pattern noted in this
23
24 data of having high odds of overweight/obesity among the elderly than the young
25
26 population.
27
28
29

30
31 Education differences were also observed for overweight/obesity. It was found that
32
33 individuals who had secondary and tertiary education were more likely to be
34
35 overweight/obese than individuals with primary or less education. This finding
36
37 corroborates other studies which have shown that education may be associated with
38
39 overweight/obesity via socioeconomic status, literacy and health behaviors⁵⁰.
40
41 Furthermore, studies which have used education as a proxy for socioeconomic
42
43 status in developing countries have shown that individuals with high educational
44
45 attainment were more likely to be obese⁵¹. The relationship between educational
46
47 attainment and obesity also depends on the individual's level of development, such
48
49 that positive associations are more common in more educated groups⁵².
50
51
52
53

54
55 There was no significant relationship observed between overweight/obesity and work
56
57 status and wealth status in the general study population. The observed pattern of no
58
59 association between overweight/obesity and work and wealth status may suggest
60

1
2
3 that in Botswana, there is no diet differences based on the wealth and work status of
4 individuals. Moreover, Botswana has over the decades moved from lower human
5 development index (HDI) to middle HDI, consequently leading to nutrition transition
6 where both the poor and non-poor have access to diets with more fat, more meat,
7 added sugars and bigger portion sizes, and lower physical activity.
8
9

10
11
12 Adjustment for known risk factors suggested that variations in smoking accounted for
13 part of the socioeconomic variation in overweight/obesity. It was noted the odds of
14 overweight/obesity increased two times among smokers than non-smokers. Chiolero
15 *et al.*⁴¹ based on literature review observed that heavy smokers tend to have greater
16 body weight than do light smokers or non-smokers, which likely reflect a clustering of
17 other risky behaviours (e.g. low level of physical activity, poor diet, and smoking) that
18 is conducive to weight gain²³⁻²⁴. In this dataset it was observed that smoking, among
19 men clustered with poor diet and alcohol consumption.
20
21
22

23
24
25 Overweight/obesity was significantly associated with poor physical activity. A decline
26 in physical activity leads to adiposity. This has been established in literature⁵³⁻⁵⁴. It
27 has been shown that although there is genetic predisposition to overweight/obesity,
28 overweight/obesity can also result from an energy imbalance between calories
29 consumed and calories expended⁵⁵. This may be due to lack of physical activity
30 ultimately leading to decreases in energy expenditure (perhaps creating a chronic
31 energy imbalance), than to increases in energy intake, strongly implicating physical
32 inactivity in the aetiology of overweight/obesity. The relationship between overweight
33 and obesity and poor physical activity can be bidirectional, whereby poor physical
34 activity leads to overweight/obesity and also overweight/obesity may lead to further
35 poor physical activity hence predisposing individuals to chronic diseases such as
36 hypertension and diabetes²⁹.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusion

Findings from this study indicate high levels of overweight/obesity in Botswana. Key socioeconomic factors associated with overweight/obesity are being a woman, increasing age, and high education (secondary and tertiary level education). Behavioral factors found to be positively associated with overweight/obesity were smoking and poor physical activity. These results demonstrate that socioeconomic and behavioural factors play an important role in prevalence of overweight/obesity in Botswana. Consequently any programs and interventions for reduction and management of overweight/obesity should focus on women, adults, individuals with high education, smokers and physically inactive people.

Acknowledgments

We wish to thank University of Botswana for affording us the time and resources to undertake this study. We also wish to thank the Office of Research of Development (ORD) at University of Botswana for providing funding for the survey on 'Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course' (NCD study) from which data for this article was derived.

Contributors

MK conceived and undertook the study as part of his PhD research. KN was the lead investigator, while MK, GL, KB, SDR, LG, TM and RM were co-investigators and takes responsibility for the integrity of the data. KN, KB, and MK oversaw the NCD study design and data analysis. MK wrote the first draft of the article. All authors reviewed and approved the final draft of the article.

Funding

The 'Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course' survey was supported by Office of Research and Development, Round 28 Funding.

Disclaimer

The funding body had no role in the design of the study, analyses, interpretation or decision to submit the manuscript for publication.

Competing Interests

None declared

Participant consent

Obtained

Ethics Approval

All ethical clearance formalities were completed before the start of the study. The study proposal along with the necessary documents was submitted to Institutional Review Board of the University of Botswana and Ministry of Health and Wellness for ethical clearance.

Data sharing statement

The dataset used and/or analysed during the current study is available from the corresponding author (Mpho Keetile) on reasonable request from the Office of Research and Development, University of Botswana.

References

1. World Health Organization, The global status of overweight and obesity. Fact Sheet. (2016), Geneva.
2. Musaiger AO, Al-Hazzaa HM. Prevalence and risk factors associated with nutrition-related noncommunicable diseases in the Eastern Mediterranean region. *Int J Gen Med*. 2012;5:199-217.
3. Baldwin W, Kaneda T, Amato L, et al. Noncommunicable diseases and youth: a critical window of opportunity for Latin America and the Caribbean. Washington, DC: Population Reference Bureau; 2013. Used with permission.
4. Bhurosy T, Jeewon R. Overweight and obesity epidemic in developing countries: a problem with diet, physical activity, or socioeconomic status?. *ScientificWorldJournal*. 2014;2014:964236.
5. Ford N.D., Patel S.A, Narayan K.M.V, Obesity in Low- and Middle-Income Countries: Burden, Drivers, and Emerging Challenges. *Annu. Rev. Public Health* 2017. 38:145–64
6. Amugsi DA, Dimbuene ZT, Mberu B, et al. Prevalence and time trends in overweight and obesity among urban women: an analysis of demographic and health surveys data from 24 African countries, 1991–2014. *BMJ*;7:e017344. doi:10.1136/bmjopen-2017-017344.
7. Hruby A, Hu FB. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*. 2015;33(7):673-89.
8. Mamun A.A , Finlay J.E, Shifting of under-nutrition to over-nutrition and its determinants among women of reproductive ages in the 36 low to medium income countries. *Obes Res Clin Pract*. 2015; 9: 75-86.

- 1
2
3 9. Micklesfield LK, Lambert EV, Hume DJ, et al. Socio-cultural, environmental
4 and behavioural determinants of obesity in black South African women.
5
6 *Cardiovasc J Afr.* 2013;24(9-10):369-75.
7
8
- 9
10 10. Biadgilign S, Mgutshini T, Haile D, Gebremichael B, Moges Y, Tilahun K.
11
12 Epidemiology of obesity and overweight in sub-Saharan Africa: a protocol for a
13
14 systematic review and meta-analysis. *BMJ Open.* 2017;7(11):e017666.
15
16 Published 2017 Nov 25. doi:10.1136/bmjopen-2017-017666.
17
18
- 19
20 11. Ball GD, Mushquash AR, Keaschuk RA, et al. Using intervention mapping to
21
22 develop the Parents as Agents of Change (PAC) intervention for managing
23
24 pediatric obesity. *BMC Res Notes* 2017;10:43.
25
26
- 27
28 12. Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents:
29
30 identification, associated health risks, and treatment approaches: a scientific
31
32 statement from the American Heart Association. *Circulation* 2013;128:1689–
33
34 712.
35
36
- 37
38 13. Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents:
39
40 identification, associated health risks, and treatment approaches: a scientific
41
42 statement from the American Heart Association. *Circulation* 2013;128:1689–
43
44 712
45
46
- 47
48 14. World Health Organization (WHO). Obesity and overweight. Geneva: World
49
50 Health Organization, 2016.
51
- 52
53 15. Levine J.A., Poverty and obesity in the U.S. *Diabetes* 2011;60:2667-2668.
54
- 55
56 16. Kim J, Sharma SV, Park SK. Association between socioeconomic status and
57
58 obesity in adults: evidence from the 2001 to 2009 Korea national health and
59
60 nutrition examination survey. *J Prev Med Public Health.* 2014;47(2):94-103.

- 1
2
3 17. You J, Choo J. Adolescent Overweight and Obesity: Links to Socioeconomic
4 Status and Fruit and Vegetable Intakes. *Int J Environ Res Public Health*.
5 2016;13(3):307. Published 2016 Mar 9. doi:10.3390/ijerph13030307.
6
7
8
9
10 18. Maffetone PB, Laursen PB. The Prevalence of Overfat Adults and Children in
11 the US. *Front Public Health*. 2017;5:290. Published 2017 Nov 1.
12 doi:10.3389/fpubh.2017.00290.
13
14
15
16
17 19. Lumsden MA, Hor K. Impact of obesity on the health of women in midlife. *The*
18 *Obstetrician & Gynaecologist* 2015; 17:201–8.
19
20
21
22 20. Kim TJ, von dem Knesebeck O. Income and obesity: what is the direction of
23 the relationship? A systematic review and meta-analysis. *BMJ Open*.
24 2018;8(1):e019862. Published 2018 Jan 5. doi:10.1136/bmjopen-2017-019862
25
26
27
28
29 21. Hong I, Coker-Bolt P, Anderson KR, Lee D, Velozo CA. Relationship Between
30 Physical Activity and Overweight and Obesity in Children: Findings From the
31 2012 National Health and Nutrition Examination Survey National Youth Fitness
32 Survey. *Am J Occup Ther*. 2016;70(5):7005180060p1-8.
33
34
35
36
37
38
39 22. Popkin, B. M., & Slining, M. M. New dynamics in global obesity facing low- and
40 middle-income countries. *Obesity Reviews: An Official Journal of the*
41 *International Association for the Study of Obesity*, 2013; 14(0 2), 11–20.
42 <http://doi.org/10.1111/obr.12102>.
43
44
45
46
47
48 23. Dare, S., Mackay, D. F., & Pell, J. P. Relationship between Smoking and
49 Obesity: A Cross-Sectional Study of 499,504 Middle-Aged Adults in the UK
50 General Population. *PLoS ONE*, 2015; 10(4).
51
52
53
54
55
56 24. Tuovinen, E.-L., Saarni, S. E., Männistö, S., Borodulin, K., Patja, K., Kinnunen,
57 T. H. ... Korhonen, T. Smoking status and abdominal obesity among normal-
58
59
60

- 1
2
3 and overweight/obese adults: Population-based FINRISK study. Preventive
4
5 Medicine Reports, 2016;4, 324–330.
6
7
- 8 25. Keetile M., Letamo G., & Navaneetham K. Patterns and determinants of
9
10 hypertension in Botswana, J Public Health. 2015;, 23:311–318.
11
12
- 13 26. Peltzer, K., and Phaswana-Mafuya, N. Fruit and vegetable intake and
14
15 associated factors in older adults in South Africa. Global Health Action, 2012;
16
17 5, 10.3402/
18
19
- 20 27. Nnyepi MS, Gwisai N, Lekgoa M, Seru T. Evidence of nutrition transition in
21
22 Southern Africa. Proc Nutr .2015; Soc. 17:1–9.
23
24
- 25 28. Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents:
26
27 identification, associated health risks, and treatment approaches: a scientific
28
29 statement from the American Heart Association. Circulation 2013;128:1689–
30
31 712
32
33
- 34 29. World Health Organization (WHO). Obesity and overweight. Geneva: World
35
36 Health Organization, 2016.
37
38
- 39 30. GBD 2015 Risk Factors Collaborators. Global, regional, and national
40
41 comparative risk assessment of 79 behavioural, environmental and
42
43 occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic
44
45 analysis for the Global Burden of Disease Study 2015. Lancet 2016;388:1659–
46
47 724.
48
49
- 50 31. Agymang C, Boatemaa S, Frempong Agymang G, de Graft Aikins, Obesity in
51
52 Sub Saharan Africa. Metabolic Syndrome. Springer International Publishing,
53
54 Switzerland. Ed. Rexford Ahima. 2015
55
56
57
58
59
60

- 1
2
3 32. Amoah A.G. Sociodemographic variations in obesity among Ghanaian
4 adults. *Public Health Nutr.* 2003;6:751-757
5
6
7
8 33. Biadgilign, S., Mgutshini, T., Haile, D., Gebremichael, B., Moges, Y., & Tilahun,
9 K. Epidemiology of obesity and overweight in sub-Saharan Africa: a protocol
10 for a systematic review and meta-analysis. *BMJ Open*, 2017;7(11).
11
12
13
14
15 34. Cohen, A. K., Rai, M., Rehkopf, D. H., & Abrams, B. Educational attainment
16 and obesity: A systematic review. *Obesity Reviews: An Official Journal of the*
17 *International Association for the Study of Obesity*, 2013; 14(12), 989–1005.
18 <http://doi.org/10.1111/obr.12062>.
19
20
21
22
23
24
25 35. Kanter R & Caballero B. Global Gender Disparities in Obesity: A Review
26 *American Society for Nutrition. Adv. Nutr.* 2012;3: 491–498, 2012;
27 [doi:10.3945/an.112.002063](https://doi.org/10.3945/an.112.002063).
28
29
30
31
32 36. Li L, Gower BA, Shelton RC, Wu X. Gender-Specific Relationship between
33 Obesity and Major Depression. *Front Endocrinol (Lausanne)*. 2017;8:292.
34
35
36
37 37. Pinto KA, Griep RH, Rotenberg L, Almeida MdCC, Barreto RS, et al.
38 Correction: Gender, time use and overweight and obesity in adults: Results of
39 the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *PLOS ONE*
40 2018;13(8): e0198380.
41
42
43
44
45
46
47 38. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, et al.,
48 Overweight, obesity, and depression: a systematic review and meta-analysis
49 of longitudinal studies. *Arch Gen Psychiatry.* 2010; 67:220–9.
50
51
52
53
54 39. Letamo G. The prevalence of, and factors associated with, overweight and
55 obesity in Botswana. *J Biosoc Sci.* 2011; 43(1):75-84.
56
57
58
59
60

- 1
2
3 40. Brochu P, Bouchard M, Haddad S. Physiological daily inhalation rates for
4 health risk assessment in overweight/obese children, adults, and elderly. *Risk*
5 *Anal.* 2013. doi: 10.1111/risa.12125.
6
7
8
9
10 41. Chielero A, Faeh D, Paccaud F, Cornuz J. "Consequences of smoking for
11 body weight, body fat distribution, and insulin resistance." *American Journal of*
12 *Clinical Nutrition* 2008; 87: 801-809.
13
14
15
16
17 42. Coulson T, Bruce E. Kendall, Julia Barthold, Floriane Plard, Susanne
18 Schindler, Arpat Ozgul, and Jean-Michel Gaillard, "Modelling Adaptive and
19 Nonadaptive Responses of Populations to Environmental Change," *The*
20 *American Naturalist* ,2017;190(3). 313-336.
21
22
23
24
25
26
27 43. Malete L, Mokgathe L, Nnyepi M, et al. Effects of a High Protein Food
28 Supplement on Physical Activity, Motor Performance and Health Related
29 Quality of Life of HIV Infected Botswana Children on Anti-Retroviral Therapy
30 (ART). *AIMS Public Health.* 2017;4(3):258-277.
31
32
33
34
35
36
37 44. Flegal KM, Carroll MD, Kuczmarski RJ, Johnson CL. Overweight and obesity in
38 the United States: prevalence and trends, 1960–1994. *Int J*
39 *ObesRelatMetabDisord*; 1998; 22:39–47.
40
41
42
43
44 45. Villareal D.T., Apovian C.M., Kushner R.F., Klein S., Obesity in older adults:
45 technical review and position statement of the American Society for Nutrition
46 and NAASO, The Obesity Society, *The American Journal of Clinical Nutrition*,
47 2005 82 (5), 923–934
48
49
50
51
52
53
54 46. Sperrin M, Marshall AD, Higgins V, Renehan AG, Buchan IE. Body mass index
55 relates weight to height differently in women and older adults: serial cross-
56
57
58
59
60

- 1
2
3 sectional surveys in England (1992-2011). *J Public Health (Oxf)*.
4
5 2016;38(3):607-613.
6
7
- 8 47. Gabriela Vazquez, Sue Duval, David R. Jacobs, Karri Silventoinen;
9
10 Comparison of Body Mass Index, Waist Circumference, and Waist/Hip Ratio in
11
12 Predicting Incident Diabetes: A Meta-Analysis, *Epidemiologic Reviews*, 2007
13
14 29,(1).115–128.
15
16
- 17 48. Emerging Risk Factors Collaboration, Wormser D, Kaptoge S, et al. Separate
18
19 and combined associations of body-mass index and abdominal adiposity with
20
21 cardiovascular disease: collaborative analysis of 58 prospective studies.
22
23 *Lancet*. 2011;377(9771):1085-95.
24
25
- 26 49. Morio, B., Barra, V., Ritz, P. et al. Benefit of endurance training in elderly
27
28 people over a short period is reversible. *Eur J Appl Physiol* (2000) 81: 329.
29
30
- 31 50. Fletcher J, Frisvold D. Higher education and health investments: Does more
32
33 schooling affect preventive health care use. *Journal Capital*. 2009; 3(2):144–
34
35 176.
36
37
- 38 51. Cohen AK, Rai M, Rehkopf DH, Abrams B. Educational attainment and
39
40 obesity: a systematic review. *Obes Rev*. 2013;14(12):989-1005.
41
42
- 43 52. Devaux, Marion, et al. (2011), “Exploring the Relationship Between Education
44
45 and Obesity”, *OECD Journal: Economic Studies* , Vol. 2011/1.
46
47
- 48 53. Traversy G. & Chaput Jean-Philippe (2015), Alcohol Consumption and
49
50 Obesity: An Update, *Curr Obes Rep* (2015) 4:122–130.
51
52
- 53 54. Fiscella, K., & Kitzman, H. Disparities in Academic Achievement and Health:
54
55 The Intersection of Child Education and Health Policy. *Pediatrics*, 2009; 123,
56
57 1073-1080.
58
59
60

1
2
3 55. Ford ND, Patel SA, & Narayan KM. Obesity in Low- and Middle-Income
4 Countries: Burden, Drivers, and Emerging Challenges. *Annu Rev Public*
5 *Health*. 2017;20;38:145-164.
6
7
8
9

10 56. Weichenthal, S., Hoppin, J. A., & Reeves, F. Obesity and the cardiovascular
11 health effects of fine particulate air pollution. *Obesity (Silver Spring, Md.)*,
12 2014. 22(7), 1580–1589. <http://doi.org/10.1002/oby.20748>
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open

Socioeconomic and behavioural determinants of overweight/obesity among adults in Botswana: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029570.R1
Article Type:	Original research
Date Submitted by the Author:	10-Jun-2019
Complete List of Authors:	Keetile, Mpho; University of Botswana, Population Studies Navaneetham, Kannan; University of Botswana, Population Studies Letamo, Gobopamang; University of Botswana, Population Studies Bainame, Kenabetsho; University of Botswana, Population Studies Rakgoasi, Serai; University of Botswana, Population Studies Gabaitiri, Lesego; University of Botswana, Statistics Masupe, Tiny; University of Botswana, School of Medicine Molebatsi, Robert; University of Botswana, Sociology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Global health
Keywords:	Overweight, Obesity, Socioeconomic, Behavioural, Botswana

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Socioeconomic and behavioural determinants of overweight/obesity**
6 **among adults in Botswana: A cross-sectional study**
7
8

9 Mpho Keetile¹, Kannan Navaneetham¹, Gobopamang Letamo¹, Kenabetsho
10 Bainame¹, Serai Daniel Rakgoasi¹, Lesego Gabaitiri², Tiny Masupe³, Robert
11 Molebatsi⁴.
12
13

14
15 **Author Affiliations**
16

17 ¹Department of Population Studies, University of Botswana, Gaborone, Botswana

18 ²Department of Statistics, University of Botswana, Gaborone, Botswana

19 ³School of Medicine, University of Botswana, Gaborone, Botswana

20 ⁴Department of Sociology, University of Botswana, Gaborone
21
22
23
24

25 **Corresponding author**
26

27 Mpho Keetile, Department of Population Studies, Private Bag 00705, University of
28 Botswana, Gaborone, Botswana. email: mphokeet@yahoo.com. Tel: +26771231375
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Word count: 3923

Abstract

Objective

To undertake a comprehensive assessment of socioeconomic and behavioural determinants of overweight/obesity among adult population in Botswana.

Design

The study adopted a cross-sectional design by selecting adult respondents in three cities and towns, fifteen urban villages and fifteen rural areas across Botswana using a multistage probability sampling technique.

Setting

The study was conducted in selected rural and urban areas of Botswana.

Participants

The study sample consisted of 1178 adult males and females aged 15 years and above.

Primary Outcome Measures

Objectively measured overweight/obesity.

Results

Prevalence of overweight/obesity in the study population was estimated at 41%. The adjusted odds ratio (AOR) of overweight/obesity were highest among women (AOR=2.74, 95% C.I. =1.92-3.90), in ages 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), among individuals with secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary education (AOR=1.99, 95% C.I. =1.16-3.38), smokers (AOR=2.16, 95% C.I. = 1.22-3.83) and people with poor physically activity (AOR=1.46, 95% C.I. =1.03-3.24). These were statistically significant at 5% level.

Conclusion

Being female, elderly, of high education levels, a smoker and having poor physical activity were identified as key factors for prevalence of overweight/obesity. These findings suggest the need for broad based strategies encouraging physical activity among different socioeconomic groups.

Key words

Overweight, Obesity, socioeconomic, behavioural, Botswana

Strengths and Limitations of the study

- To our knowledge this is the first cross-sectional study in Botswana to comprehensively assess both the socioeconomic and behavioural determinants of overweight/obesity among adults.
- Data were collected from a large and randomly selected representative population. The data contained information on potential confounding factors, with a low proportion of missing information making the study more comparable.
- As the cross sectional design was employed, the causal relationship between an explanatory and an outcome variable was not established.

Introduction

The disease profile in Botswana is changing, with most deaths and disability in the foreseeable future likely to be accounted for by the ominous epidemics of non-communicable diseases (NCDs) such as heart disease, stroke, cancer and other chronic diseases. Children, adults and the elderly are all vulnerable to the risk factors that contribute to NCDs especially overweight/obesity, due to unhealthy diets, low physical activity, exposure to tobacco smoke and/or the effects of the harmful use of alcohol¹⁻³. The key drivers of NCDs are ageing, rapid unplanned urbanisation, and the globalisation of unhealthy lifestyles. For example, globalization of unhealthy lifestyles like unhealthy diets may show up in individuals as raised blood pressure, increased blood glucose, elevated blood lipids, and overweight/ obesity¹.

Several studies have recently shown that many low income and middle-income countries (LMICs) are experiencing an increase in the prevalence of overweight/obesity⁴⁻⁶. This prevalence is predicted to continue to increase in the future. Much of the increase in prevalence levels for overweight/obesity is expected in countries such as India, some parts of South-East Asia, China, most of South America and some parts of Sub Saharan Africa (SSA)⁷⁻⁸. Until recently, SSA was minimally affected by the overweight/obesity epidemic due to under-nutrition and a major burden of Human Immuno Deficiency Virus (HIV) and tuberculosis (TB)⁹. Meanwhile research on overweight/obesity has started to gain significance in SSA¹⁰⁻¹⁴ due to its increasing magnitude.

It has been shown by previous studies that socioeconomic and behavioural factors have a significant association with overweight/obesity¹⁵⁻¹⁶. These studies have shown that socioeconomic and demographic factors such as sex, age, education level, work

1
2
3 status, place of residence and wealth status are significant determinants of
4
5 overweight/obesity. It has also been found that women are more likely to be
6
7 overweight/obese than men, while the odds of being overweight/obesity has been
8
9 seen to increase with age¹⁷⁻¹⁹. Other studies have shown that being employed in a
10
11 high income job, residing in urban areas, having high education and wealth status is
12
13 significantly associated with overweight/obesity²⁰.
14
15
16
17
18

19
20 There is also available evidence which suggest significant correlation between
21
22 behavioural factors and overweight/obesity. For example, there is significant evidence
23
24 of a positive association between physical inactivity and the prevalence of
25
26 overweight/obesity²¹. It has also been suggested that the relation between physical
27
28 activity and health outcomes such as overweight/obesity may be moderated by a
29
30 number of lifestyle factors, especially sedentary lifestyle^{15, 20, 21}. In many developing
31
32 countries, there has been a substantial shift from jobs with high-energy expenditure
33
34 such as farming, mining, and forestry, and employment is inclined towards the more-
35
36 sedentary sectors of manufacturing, services and office based-work²². Moreover,
37
38 there have been changes in lifestyle behaviours condoning alcohol consumption, poor
39
40 diets and smoking. These are also key risk factors of overweight/obesity²³⁻²⁴.
41
42
43
44
45
46

47
48 The Botswana WHO STEPS surveys conducted in 2007 and 2014 have shown
49
50 increasing levels of overweight/obesity in the general adult population of Botswana.
51
52 Prevalence of overweight/obesity was 18.7% in 2007²⁵ and increased to 38.6 % in
53
54 2014²⁶. Despite evidence of the high prevalence of overweight/obesity in Botswana,
55
56 there is little evidence of socioeconomic and behavioural determinants of
57
58 overweight/obesity. To the best of our knowledge this is the first cross sectional study
59
60

1
2
3 which takes into account both socioeconomic and behavioural factors for
4 understanding the heterogeneity of overweight/obesity in Botswana. The reduction of
5 overweight/obesity prevalence should be a key goal of health-care policy because of
6 its association with many NCDs. The main objective of this study was to assess
7 socioeconomic and behavioural determinants associated with overweight/obesity
8 prevalence in Botswana.
9
10
11
12
13
14
15
16
17
18

19 **Methods**

20 ***Sampling, recruitment and ethics***

21
22
23 The study used cross sectional design by selecting respondents in three cities and
24 towns, fifteen urban villages and fifteen rural areas across Botswana using a
25 multistage probability sampling technique. A list of districts, localities and enumeration
26 areas (EAs) together with their households was made based on the 2011 Botswana
27 Population and Housing census to make sampling frame. At the first stage the
28 population was stratified into cities and towns, urban villages and rural settlements. A
29 listing of all 26 census districts in each strata was made and from these districts a total
30 of all 4845 EAs were listed for rural and urban localities. At the second stage, localities
31 in urban and rural districts were randomly selected. At the third and fourth stage a
32 random selection of EAs and households, respectively was made. Lastly, individuals
33 aged 15 years and over were selected for interview from the list of households with
34 persons 15 years and over. The NCD study questionnaire was developed using the
35 instruments from the WHO Study on Global Ageing and Adult Health (SAGE), and
36 WHO STEPS Surveys.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 All ethical clearance formalities were completed before the start of the study. The study
4 proposal along with the necessary documents were submitted to and approved by the
5 Institutional Review Board of the University of Botswana. Approval for conducting the
6 study was also obtained from the Government of Botswana through the Ministry of
7 Health and Wellness. Privacy and confidentiality of highest standard were maintained
8 by treating all respondents as anonymous and no names of respondents are
9 mentioned or implied when presenting findings of the study.
10
11
12
13
14
15
16
17
18
19

20 ***Patient and Public involvement***

21
22
23 No patients were involved in developing the research question, outcome measures
24 and overall design of the study. Since this was a cross-sectional study, respondents
25 were drawn from the general population rather than from healthcare settings, thereby
26 providing access to all eligible people aged 15 years and above. As part of the study
27 protocol, all respondents were requested to sign a consent form and were told that
28 participation in the study was voluntary. For participants aged 15-17 years both
29 consent and assent were obtained. All individuals aged 15 years and above who had
30 successfully completed the NCD study questionnaire were included in the analysis for
31 this study.
32
33
34
35
36
37
38
39
40
41
42
43
44

45 ***Definitions and Measurement of variables***

46
47
48 The outcome variable for the study is overweight/obesity. Body Mass Index (BMI) was
49 used to classify overweight/obesity. BMI was derived from weight and height: weight
50 (kg) / (height (m) x height (m))¹. The Charder MS7301 250Kg digital scale and the
51 Muac measuring tape were used for anthropometric measurements. Weight was
52 measured, to the nearest 0.1kg, while height was measured in metres. BMI was
53 categorized into: underweight (BMI < 18.5 kg/m²), normal weight (18.5 ≤ BMI <25
54
55
56
57
58
59
60

1
2
3 kg/m²), overweight ($25 \leq \text{BMI} < 30 \text{ kg/m}^2$) and obese ($\text{BMI} \geq 30 \text{ kg/m}^2$)¹. Overweight and
4
5 obese were used to create a binary outcome variable which was coded as: being
6
7 overweight/obese ($\text{BMI} \geq 25 \text{ kg/m}^2$) =1; not overweight/obese =0 ($\text{BMI} < 25 \text{ kg/m}^2$).
8
9

10
11 Information on NCD risk factors was collected through self-reports. Respondents were
12
13 asked questions on NCD risk factors such as tobacco use, alcohol consumption,
14
15 physical activity and fruit and vegetable consumption. For tobacco use, respondents
16
17 were asked whether they currently smoke tobacco products and their response was
18
19 coded as yes=1 and no=0. Alcohol consumption was measured based on the intensity
20
21 of alcohol consumed in the past 30 days. Respondents who had consumed alcohol in
22
23 the past 30 days were asked about the number of standard alcohol drinks they had
24
25 each day in the past 7 days and if they reported to have had three or more drinks per
26
27 day (of approximately 60g alcohol) it was considered to be excessive drinking=1 and
28
29 0=otherwise.
30
31
32

33
34 Based on WHO recommendations poor physical activity was calculated using an
35
36 average typical types of activity undertaken¹³. It was calculated based on the time
37
38 taken while doing physical activity in the past 7 days. Respondents were asked
39
40 whether they do any moderate to rigorous intensity activities for at least 10 minutes
41
42 continuously. This was considered for the domains of work and walking (includes at
43
44 work and at home, walking to travel from place to place, and any other walking for
45
46 recreation, sport, exercise, or leisure). Based on the time taken on work and walking
47
48 the respondents were grouped into four categories of: no activity, low, moderate and
49
50 high activity levels to show the intensity of their physical activity. The resultant variable
51
52 was coded such that no and low activity (<10 minutes of physical activity) =1 and
53
54 moderate and high activity (≥ 10 minutes of physical activity) were coded=0.
55
56
57
58
59
60

Poor fruit and vegetables consumption was created when an individual reported daily consumption of less than the recommended 5 servings of fruit and vegetables. Respondents reported the number of servings for fruits/vegetables they had in a typical day, and if the servings were less than 5 in a day, they were considered to be having poor fruit/vegetable consumption^{13, 27, 28}.

A wealth index (WI) was constructed to be a measure of wealth status. WI is a composite measure of indicators of ownership of consumer durables, housing characteristics, and access to public services¹⁶. Information on a range of durable assets was collected during the survey (e. g. car, refrigerator, television,), housing characteristics (e. g. material of dwelling floor and roof, main cooking fuel), access to basic services (e. g. electricity supply, source of drinking water, sanitation facilities) and ownership of livestock (e.g. cattle, goats, sheep, horses, chickens). Further to collection of information on durable assets, information on ownership of land was also collected. Principal component analysis was employed to derive the wealth index variable, which had five categories from the 1st to the 5th quintile (poorest to richest).

Control variables

Age and place of residence were used as control variables. These variables were conceptualized to have an association with the outcome variable. Therefore, as control variables they are included in the combined effects model, so that the association between the independent variables of interest becomes isolated and discernible.

Statistical analysis

Data analysis involved descriptive and multivariable analysis. Descriptive analysis provided some basis for further exploration of key determinants of overweight and obesity in Botswana. In the multivariable binary logistic regression, outcome variable

was coded as '1' if respondents were overweight/obese and '0' if they were not overweight/obese. Results of logistic regression analysis were presented together with adjusted odds ratios (AOR), 95% confidence intervals (CI) and 5% statistical significance level. Chi-square test was used for bivariate analysis. Data analysis was done using SPSS version 25 program. Since this was a population based study representative of the population, selection bias was minimal. Moreover the use of systematic random sampling across the population meant that selection bias was minimised further. In order to control for cluster effects complex samples module in SPSS has been used since the NCDs survey used multistage sampling technique.

Results

Sample characteristics

Table 1 presents results on the sample description for the respondents. The results show that a total of 1,178 respondents between the ages of 15 years and above were successfully interviewed. The sample constituted a high proportion of females (69.1%) than males (30.9%). The sample age distribution suggests a relatively young population, with over half (59%) of the sample being less than 39 years of age, and almost three quarters (73.5%) being less than fifty years of age. As age increases, the sample skewed in favour of females.

Table 1: Socio demographic characteristics

Variable	N	%
Sex		
Male	364	30.9
Female	813	69.1
Total	1177	100

Age in years		
<24	270	26.4
25– 34	302	29.5
35 – 44	196	19.2
45– 54	130	12.7
55 – 64	75	7.3
65+ years	50	4.9
Total	1023	100
Locality Type		
Cities/Towns	355	30.2
Urban Villages	534	45.4
Rural Settlements	288	24.5
Total	1177	100
Marital Status		
Never Married	864	73.8
Currently married	199	17.0
Formerly married	108	9.2
Total	1171	100
Highest Level of Education Attained		
Primary or Less	410	35.5
Junior Secondary	314	27.2
Senior Secondary	200	17.3
Tertiary & Over	230	19.9
Total	1154	100
Work Status in past 12 months		
Public Sector	122	10.5
Private Sector	182	15.7
Self Employed	130	11.2
Not Employed	436	37.5
Homemaker-Student	218	18.8
Retired-Other	74	6.4
Total	1162	100
Wealth status		
Lowest	234	19.9
Second	237	20.1
Middle	235	19.9
Fourth	237	20.1
Highest	235	19.9
Total	1178	100

More than two fifths (45.4%) of the population resided in urban villages; while the remaining proportion resided in cities and towns (30.2%), and rural areas (24.5%).

1
2
3 Almost three quarters (73.8%) of respondents were never married; over a third
4
5 (35.5%) had primary education or less; over a quarter (27.2%) had junior secondary
6
7 education while just under a fifth had senior secondary education (17.3%) and tertiary
8
9 education (19.9%). Close to two fifth (37.5%) in the sample were not employed; over
10
11 a quarter were employed in either the public or private sector (26.2%) while the
12
13 remaining proportion were either self-employed (11.2%) , home makers or students
14
15 (18.2%) and retirees (6.4%).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2 below shows behavioural characteristics of the study population. It shows that prevalence of alcohol consumption was estimated at 17.3% while tobacco smoking was estimated at 11.5%. Poor physical activity was also high in the study population with 48.9% of respondents indicating poor physical activity. Poor fruit and vegetable intake was estimated at 82.5%.

Table 2: Behavioural characteristics of the study population

Variable	N	%
Smoking		
Yes	136	11.5
No	1042	88.5
Total	1178	100
Alcohol consumption		
Yes	204	17.3
No	974	82.7
Total	1178	100
Poor physical activity		
Yes	576	48.9
No	602	51.1
Total	1178	100
Poor fruit/vegetable intake		
Yes	1045	82.5
No	133	17.5
Total	1178	100

Prevalence of overweight/obesity

Estimated prevalence of overweight/obesity in the study population was 41.3% (Table 3). There were significant gender differences in prevalence of overweight/obesity. It was found that women than men (48.6% among women, 25.5% among men) were overweight/obese. It was observed that individuals residing in rural areas (46.8%), public sector employees (55.2%) and those with lower education (45%) showed high prevalence rates for overweight/obesity. Meanwhile it was found that wealth status was not significantly linked with overweight/obesity. Among behavioural factors this

1
2
3 study offers evidence that non-smoker (42.9%) showed high prevalence rates of
4
5 overweight/obesity. Alcohol consumption, poor physical activity and poor fruits and
6
7 vegetable consumption did not show any significant association with
8
9
10 overweight/obesity.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 3: Prevalence of overweight/obesity by socioeconomic and behavioural characteristics of the study population

Variable	N	%	P-value
Sex			
Male	364	25.5	0.000**
Female	813	48.6	
Age			0.000**
≤24	270	21.8	
25-34	302	36.8	
35-44	196	53.0	
45-54	130	54.5	
55-64	75	61.8	
65+	50	43.8	
Marital status			0.000**
Never-married	864	35.5	
Currently-married	199	62.2	
Formerly-married	108	54.1	
Education			0.019**
Primary or less	410	45.4	
Secondary	514	36.7	
Tertiary or higher	230	44.6	
Residence			0.008**
Cities and towns	355	37.9	
Urban villages	534	41.0	
Rural villages	288	46.8	
Work status			0.000**
Public sector	122	52.2	
Private sector	182	36.8	
self-employed	130	55.0	
Not employed	436	45.8	
Home-maker/student	218	29.4	
Retired/other	74	40.3	
Wealth status			0.765
Lowest	234	38.8	
second	237	43.2	

middle	235	41.1	
Fourth	237	39.7	
Highest	235	44.1	
Alcohol consumption			0.445
No	766	33.3	
yes	412	37.7	
Smoking			0.004**
Yes	136	29.7	
No	1042	42.9	
Poor physical activity			0.547
Yes	376	41.5	
No	771	39.6	
Poor fruit/vegetable consumption			0.832
No	133	41.9	
yes	1045	43.0	
Overall	1178	41.3	

Note: **Chi-square test statistically significant at $p < 0.05$.

Determinants of overweight/obesity

Table 4 indicates results of the logistic regression model for overweight/obesity with socioeconomic and behavioural factors. One unadjusted and two adjusted models were run. Model I assessed the association between overweight/obesity and each of the socioeconomic variables and behavioural factors, Model II assessed the association between overweight/obesity adjusting for socioeconomic factors, while Model III assess the association between overweight/obesity adjusting for both socioeconomic and behavioural factors. It was observed that gender had a significant association with overweight/obesity when controlling for both socioeconomic and behavioural factors. Women were observed to have more than 2 times (AOR=2.74, 95% C.I. =1.92-3.90) greater odds for overweight/obesity than men.

1
2
3 The odds of being overweight/obese increased with age after controlling for
4 socioeconomic and behavioural factors and were highest among individuals aged 55-
5
6
7
8 64 years (AOR=5.53, 95% C.I. = 2.62-11.6), but declined at ages 65+ years
9
10 (AOR=2.88, 95% C.I. = 1.21-6.86). Education was also found to be a significant factor
11
12 associated with overweight/obesity in the study population after controlling for
13
14 socioeconomic covariates. Further, individuals who had secondary (AOR=1.70, 95%
15
16 C.I. = 1.11-2.61) and tertiary or higher education (AOR=1.99, 95% C.I. = 1.16-3.38)
17
18 were found to have higher probability of being overweight/obese than individuals with
19
20 primary or less education. Meanwhile, there was no significant relationship observed
21
22 between work status, wealth status and overweight/obesity in the general study
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The association between smoking and alcohol consumption with overweight/obesity was found to be negative in the crude model. However, controlling for other socioeconomic and demographic factors, the odds of overweight/obesity increased two times (AOR=2.16, 95% C.I. = 1.22-3.83) among smokers than non-smokers. Overweight/obesity was also significantly associated with poor physical activity (AOR=1.46, 95% C.I. = 1.03-3.24) after adjusting for socioeconomic and behavioural factors.

Table 4: Odds ratios (OR) for the association of Overweight/obesity with socioeconomic and behavioural variables factors in the study population, NCD study 2016.

Variable	Model I		Model II		Model III	
	OR	C.I.	OR	C.I.	OR	C.I.
Sex						
Male	1.00		1.00		1.00	
Female	2.75**	(2.08-3.64)	3.01**	(2.12-4.26)	2.74**	(1.92-3.90)
Age						
≤24	1.00		1.00		1.00	
25-34	2.08**	(1.42-3.03)	1.77**	(1.12-2.77)	1.83**	(1.17-2.88)
35-44	4.02**	(2.67-6.08)	3.70**	(2.23-6.12)	3.87**	(2.34-6.42)
45-54	4.28**	(2.70-6.78)	3.04**	(1.66-5.55)	3.26**	(1.78-5.98)
55-64	5.78**	(3.26-10.2)	4.91**	(2.35-10.2)	5.53**	(2.62-11.6)
65+	2.78**	(1.46-5.28)	2.74**	(1.15-6.50)	2.88**	(1.21-6.86)
Marital status						
Never-married	0.46**	(0.30-0.71)	0.68	(0.34-1.34)	0.66	(0.33-1.32)
Currently-married	1.39	(0.84-2.29)	1.34	(0.66-2.71)	1.28	(0.63-2.61)
Formerly-married	1.00		1.00		1.00	
Education						
Primary or less	1.00		1.00		1.00	
Secondary	0.69**	(0.53-0.91)	0.49**	(0.29-0.83)	1.70**	(1.11-2.61)
Tertiary or higher	0.96	(0.69-1.35)	0.87	(0.58-1.32)	1.99**	(1.16-3.38)
Residence						
Cities and towns	1.00		1.00		1.00	
Urban villages	1.13	(0.86-1.50)	1.06	(0.74-1.50)	1.07	(0.75-1.52)
Rural villages	1.44**	(1.04-1.99)	1.37	(0.88-2.12)	1.37	(0.88-2.12)
Work status						
Public sector	1.61	(0.89-2.93)	1.30	(0.64-2.65)	1.22	(0.59-2.50)
Private sector	0.86	(0.49-1.51)	1.20	(0.61-2.36)	1.19	(0.60-2.35)
self-employed	1.18	(0.65-2.13)	1.24	(0.61-2.51)	1.29	(0.63-2.62)
Not employed	1.25	(0.75-2.08)	1.30	(0.70-2.40)	1.30	(0.70-2.41)
Home-maker/student	0.61	(0.35-1.07)	0.96	(0.48-1.92)	0.96	(0.48-1.92)
Retired/other	1.00		1.00		1.00	
Wealth status						
Lowest	0.80	(0.55-1.17)	0.60	(0.34-1.05)	1.03	(0.64-1.68)
second	0.96	(0.66-1.40)	0.67	(0.40-1.11)	1.12	(0.68-1.85)
middle	0.88	(0.60-1.28)	0.73	(0.45-1.19)	1.20	(0.71-2.01)

Fourth	0.83 (0.57-1.21)	0.79 (0.50-1.23)	1.53 (0.87-2.69)
Highest	1.00	1.00	1.00
Smoking		N.I	
Yes	0.56* (0.37-0.83)		2.16** (1.22-3.83)
No	1.00		1.00
Poor physical activity		N.I	
Yes	1.08 (0.83-1.40)		1.46** (1.03-3.24)
No	1.00		1.00
Poor fruit/vegetable consumption		N.I	
Yes	0.82 (0.55-1.22)		0.80 (0.28-2.24)
no	1.00		1.00
Alcohol consumption		N.I	
Yes	0.66* (0.47-0.91)		1.23 (0.53-2.83)
No	1.00		1.00

N.I- not included, Model I: Crude model; Model II: socioeconomic variables included; Model III: socioeconomic + behavioural characteristics included. **statistically significant at $p < 0.05$.

Discussion

The prevalence of overweight/obesity in the study population was 41.3%. Botswana is also joined with other Southern African countries where the national prevalence rates range between 30% and 60% among populations over the age of 15 years in most countries^{10, 26-31}. Botswana is currently undergoing a nutrition transition from high-fibre, low-calorie diets rich in fruits and vegetables to refined, energy-dense foods high in fat, calories, sweeteners and salt¹² and to low physical activity due to transition from agrarian to a sedentary industrial society³² contributing to a greater risk of overweight/obesity.

The observed socioeconomic differences in overweight/obesity prevalence is well-established finding, and consistent with most previous studies³³⁻³⁴. For example, women were found to have higher odds of being overweight/obese than men. Similar findings were observed in most studies in developing countries³⁵⁻³⁷. However, in some developed countries such as US and UK, it was found that adult men tend to be overweight/obese than adult women³⁸⁻⁴². Contrastingly in developing countries such as Botswana adult women are generally noted to be overweight/obese than men at all ages^{26, 35, 39-40}.

In the context of Botswana, observed gender differences in overweight/obesity in this study may be explained by sociocultural dynamics, whereby acculturation, through complex sociocultural pathways encourages weight gain among women^{26, 40}. For example, in Botswana cultural values favour larger body size among women as a sign of fertility, healthfulness, or prosperity²⁶. One indicative finding from this study was that women reported low levels of physical activity which may also contribute to observed high prevalence of overweight/obesity among them.

1
2
3 The study found that age is associated with overweight/obesity even after controlling
4 for socioeconomic and behavioural factors. Overweight/obesity was highest among
5 individuals aged 55-64 years but declined at ages 65+ years. This finding corroborates
6 data from large population studies⁴³⁻⁴⁶ which show that mean body weight and BMI
7 gradually increases during most of adult life and reach peak values at 50–59 years of
8 age in both men and women and after the age of 60 years, mean body weight and
9 BMI tend to decrease⁴⁷⁻⁴⁹. In South Africa similar findings have been observed that
10 overweight/obesity tends to increase with age²¹.

11
12
13
14
15
16
17
18
19
20
21
22 Education differences were also observed for overweight/obesity. It was found that
23 individuals who had high educational attainment had higher odds of overweight/obese
24 than those with low education. This finding corroborates other studies which have
25 shown that education may be associated with overweight/obesity via socioeconomic
26 status, literacy and health behaviors⁵⁰⁻⁵². There was no significant relationship
27 observed between overweight/obesity and work and wealth status. Other studies in
28 neighbouring countries with comparable Human Development Index (HDI) as
29 Botswana including Namibia and South Africa²⁷⁻²⁸ have also shown no wealth or work
30 status differences in overweight/obesity. There are several reasons to such scenario.
31 These reasons include; rapid and disorganized urbanization, increasing sedentary
32 lifestyles, easy access to and consumption of unhealthy food and high energy drinks
33 among both the poor and non-poor in these countries^{21, 27-28}. Further to this a general
34 observation has been made by Ford et al.⁵² that overweight/obesity is no longer an
35 issue of affluence in majority of low and middle income countries.

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Further, the study noted that the odds of overweight/obesity were two times greater
55 among smokers than non-smokers. Tuovinen *et al.*⁴¹ based on literature review
56 observed that heavy smokers tend to have greater body weight than do light smokers
57
58
59
60

1
2
3 or non-smokers, which likely reflect a clustering of other risky behaviours (e.g. low
4 level of physical activity and poor diet) that is conducive to weight gain²³⁻²⁴. In this
5 study, it was observed that smoking, especially among men clustered with poor diet
6 and alcohol consumption. On the other hand some studies have shown that smoking
7 tends to suppress appetite leading to low BMI among smokers^{10, 21, 30}.

8
9
10
11
12
13
14
15 Other important finding in this study is that overweight/obesity was significantly
16 associated with poor physical activity. This is consistent with other studies which have
17 shown that a decline in physical activity leads to adiposity⁵². The interpretation of this
18 finding is that Botswana has undergone a transition from an agrarian to industrial
19 society due to urbanization subsequently leading to industrial modernity.
20 Consequently this has led to an increase in poor physical activity in the general
21 population. Poor physical activity leads to overweight/obesity through energy
22 imbalance between calories consumed and calories expended⁵². Generally formal
23 physical activity that is planned, structured, repetitive, and purposeful is not common
24 in Botswana. Consequently in order to increase physical activity and reduce
25 overweight/obesity there is need for a population-based, multi-sectoral, multi-
26 disciplinary, and culturally relevant approach.

27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 **Conclusion**

44 Findings from this study indicate higher prevalence of overweight/obesity in Botswana.
45 Key socioeconomic factors associated with overweight/obesity are being a woman,
46 older, and with higher education (secondary and tertiary level education). Behavioral
47 factors such as smoking and poor physical activity were found to be positively
48 associated with overweight/obesity. These results demonstrate that socioeconomic
49 and behavioural factors play an important role in prevalence of overweight/obesity in
50 Botswana. Botswana's multisectoral national strategic plan implementation framework

1
2
3 for combating NCDs should focus on developing a community based and people
4 centered awareness about the harmful impact of smoking, alcohol, physically inactivity
5 and poor quality of diet facilitated by comprehensive national monitoring and evaluation
6 framework.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Acknowledgments

We wish to thank University of Botswana for affording us the time and resources to undertake this study. We also wish to thank the Office of Research of Development (ORD) at University of Botswana for providing funding for the survey on 'Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course' (NCD study) from which data for this article was derived. We also wish to thank the study participants.

Contributors

MK conceived and undertook the study as part of his PhD research. KN was the lead investigator, while MK, GL, KB, SDR, LG, TM and RM were co-investigators and takes responsibility for the integrity of the data. KN, KB, and MK oversaw the NCD study design and data analysis. MK wrote the first draft of the article. All authors reviewed and approved the final draft of the article.

Funding

The 'Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course' survey was supported by Office of Research and Development, Round 28 Funding.

Disclaimer

The funding body had no role in the design of the study, analyses, interpretation or decision to submit the manuscript for publication.

Competing Interests

Nil

Participant consent

Obtained

Ethics Approval

All ethical clearance formalities were completed before the start of the study. The study proposal along with the necessary documents was submitted to and approved by the Institutional Review Board of the University of Botswana (Ref #: UBR/RES/IRB/1583) and the Ministry of Health and Wellness (Ref #: HPDME: 13/18/1 Vol. X (130)). Privacy and confidentiality of highest standard were maintained by treating all respondents as anonymous and no names of respondents are mentioned or implied when presenting findings of the study.

Data sharing statement

The dataset used and/or analysed during the current study is available from the corresponding author (Mpho Keetile) on reasonable request from the Office of Research and Development, University of Botswana.

References

1. World Health Organization, The global status of overweight and obesity. *Fact Sheet*. Geneva. 2016.
2. Musaiger AO, Al-Hazzaa HM. Prevalence and risk factors associated with nutrition-related noncommunicable diseases in the Eastern Mediterranean region. *Int J Gen Med*. 2012;5:199-217.
3. Baldwin W, Kaneda T, Amato L, et al. Noncommunicable diseases and youth: a critical window of opportunity for Latin America and the Caribbean. Washington, DC: *Population Reference Bureau*; 2013.
4. Bhurosy T, Jeewon R. Overweight and obesity epidemic in developing countries: a problem with diet, physical activity, or socioeconomic status?. *ScientificWorldJournal*. 2014;;964236. doi:10.1155/2014/964236
5. Ford ND., Patel SA, Narayan KMV. Obesity in Low- and Middle-Income Countries: Burden, Drivers, and Emerging Challenges. *Annu. Rev. Public Health*. 2017; 38:145–64
6. Amugsi DA, Dimbuene ZT, Mberu B, et al. Prevalence and time trends in overweight and obesity among urban women: an analysis of demographic and health surveys data from 24 African countries, 1991–2014. *BMJ*;7:e017344. doi:10.1136/bmjopen-2017-017344.
7. Hruby A, Hu FB. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*. 2015;33(7):673-89.
8. Mamun A.A , Finlay J.E. Shifting of under-nutrition to over-nutrition and its determinants among women of reproductive ages in the 36 low to medium income countries. *Obes Res Clin Pract*. 2015; 9: 75-86.

- 1
2
3 9. Micklesfield LK, Lambert EV, Hume DJ, et al. Socio-cultural, environmental and
4 behavioural determinants of obesity in black South African women. *Cardiovasc*
5
6 *J Afr.* 2013;24(9-10):369-75.
7
8
9
10 10. Biadgilign S, Mgotshini T, Haile D, Gebremichael B, Moges Y, Tilahun K.
11
12 Epidemiology of obesity and overweight in sub-Saharan Africa: a protocol for a
13
14 systematic review and meta-analysis. *BMJ Open.* 2017;7(11):e017666.
15
16
17 11. Ball GD, Mushquash AR, Keaschuk RA, et al. Using intervention mapping to
18
19 develop the Parents as Agents of Change (PAC) intervention for managing
20
21 pediatric obesity. *BMC Res Notes* 2017;10:43.
22
23
24 12. Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents:
25
26 identification, associated health risks, and treatment approaches: a scientific
27
28 statement from the American Heart Association. *Circulation.* 2013;128:1689–
29
30 712.
31
32
33 13. World Health Organization (WHO). Obesity and overweight. *World Health*
34
35 *Organization, Geneva.* 2016.
36
37
38 14. Levine J.A. Poverty and obesity in the U.S. *Diabetes.* 2011; 60:2667-2668.
39
40
41 15. Kim J, Sharma SV, Park SK. Association between socioeconomic status and
42
43 obesity in adults: evidence from the 2001 to 2009 Korea national health and
44
45 nutrition examination survey. *J Prev Med Public Health.* 2014;47(2):94-103.
46
47
48 16. You J, Choo J. Adolescent Overweight and Obesity: Links to Socioeconomic
49
50 Status and Fruit and Vegetable Intakes. *Int J Environ Res Public Health.*
51
52 2016;13(3):307.
53
54
55 17. Maffetone PB, Laursen PB. The Prevalence of Overfat Adults and Children in
56
57 the US. *Front Public Health.* 2017;5:290.
58
59
60

- 1
2
3 18. Lumsden MA, Hor K. Impact of obesity on the health of women in midlife. *The*
4 *Obstetrician & Gynaecologist*. 2015; 17 (1):201–8.
5
6
7
8 19. Kim TJ, von dem Knesebeck O. Income and obesity: what is the direction of the
9 relationship? A systematic review and meta-analysis. *BMJ Open*.
10 2018;8(1):e019862.
11
12
13 20. Hong I, Coker-Bolt P, Anderson KR, Lee D, Velozo CA. Relationship Between
14 Physical Activity and Overweight and Obesity in Children: Findings From the
15 2012 National Health and Nutrition Examination Survey National Youth Fitness
16 Survey. *Am J Occup Ther*. 2016; 70(5):7005180060p1-8.
17
18
19 21. Popkin BM, Slining MM. New dynamics in global obesity facing low- and middle-
20 income countries. *Obesity Reviews: An Official Journal of the International*
21 *Association for the Study of Obesity*. 2013; 14(02), 11–20.
22 <http://doi.org/10.1111/obr.12102>.
23
24
25 22. Dare S, Mackay DF, Pell JP. Relationship between Smoking and Obesity: A
26 Cross-Sectional Study of 499,504 Middle-Aged Adults in the UK General
27 Population. *PLoS ONE*. 2015; 10(4).
28
29
30 23. Tuovinen EL, Saarni SE, Männistö S, et al. Smoking status and abdominal
31 obesity among normal- and overweight/obese adults: Population-based
32 FINRISK study. *Preventive Medicine Reports*. 2016;4(1), 324–330.
33
34
35 24. Keetile M, Letamo G, Navaneetham K. Patterns and determinants of
36 hypertension in Botswana, *J Public Health*. 2015; 23(1),311–318.
37
38
39 25. Republic of Botswana. Chronic Disease Risk Surveillance Report 2010, Ministry
40 of Health & WHO, *Government Printers*. 2010.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 26. Letamo G. The prevalence of, and factors associated with, overweight and
4 obesity in Botswana. *J Biosoc Sci.* 2011; 43(1):75-84.
5
6
7 27. Peltzer K, Phaswana-Mafuya N. Fruit and vegetable intake and associated
8 factors in older adults in South Africa. *Global Health Action*, 2012; 5(1), 1-8.
9 doi:10.3402/gha.v5i0.18668
10
11 28. Nnyepi MS, Gwisai N, Lekgoa M, et al. Evidence of nutrition transition in
12 Southern Africa. *Proc Nutr Soc.* 2015; 17 (1):1–9. doi:
13 10.1017/S0029665115000051
14
15
16
17
18 29. World Health Organization (WHO). Obesity and overweight. Geneva: *World*
19 *Health Organization*, 2015.
20
21
22
23 30. GBD 2015 Risk Factors Collaborators. Global, regional, and national
24 comparative risk assessment of 79 behavioural, environmental and
25 occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic
26 analysis for the Global Burden of Disease Study 2015. *Lancet.* 2016;388
27 (1):1659–724.
28
29
30
31
32
33
34
35 31. Agymang C, Boatemaa S, Frempong-Agymang G, et al. Obesity in Sub Saharan
36 Africa. *Metabolic Syndrome.* Springer International Publishing, Switzerland. Ed.
37 Rexford Ahima. 2015
38
39
40
41
42 32. Amoah AG. Socio-demographic variations in obesity among Ghanaian adults.
43 *Public Health Nutr.* 2003;6 (1):751-757
44
45
46
47 33. Biadgilign S, Mgutshini T, Haile D., et al. Epidemiology of obesity and overweight
48 in sub-Saharan Africa: a protocol for a systematic review and meta-analysis.
49 *BMJ Open*, 2017; 7(11).
50
51
52
53
54 34. Cohen AK, Rai M, Rehkopf DH. et al. Educational attainment and obesity: A
55 systematic review. *Obesity Reviews: An Official Journal of the International*
56
57
58
59
60

- 1
2
3 *Association for the Study of Obesity*, 2013; 14(12), 989–1005.
4
5 <http://doi.org/10.1111/obr.12062>.
6
7
8 35. Kanter R, Caballero B. Global Gender Disparities in Obesity: A Review of
9 American Society for Nutrition. *Adv. Nutr.* 2012;3 (1): 491–498, 2012;
10 doi:10.3945/an.112.002063.
11
12 36. Li L, Gower BA, Shelton RC, Wu X. Gender-Specific Relationship between
13 Obesity and Major Depression. *Front Endocrinol (Lausanne)*. 2017;8 (1):292.
14
15 37. Pinto KA, Griep RH, Rotenberg L, Almeida MDCC. et al. Correction: Gender,
16 time use and overweight and obesity in adults: Results of the Brazilian
17 Longitudinal Study of Adult Health (ELSA-Brasil). *PLOS ONE* 2018;13(8):
18 e0198380.
19
20 38. Luppino FS, de Wit LM, Bouvy PF. et al. Overweight, obesity, and depression:
21 a systematic review and meta-analysis of longitudinal studies. *Arch Gen*
22 *Psychiatry*. 2010; 67(1):220–9.
23
24 39. The GBD 2015 Obesity Collaborators. Health Effects of Overweight and Obesity
25 in 195 Countries over 25 Years. *N Engl J Med* 2017; 377:13-27.
26
27 40. Kamadjeu RM, Edwards R, Atanga JS, Kiawi EC, Unwin N, Mbanya J-C.
28 Anthropometry measures and prevalence of obesity in the urban adult
29 population of Cameroon: an update from the Cameroon Burden of Diabetes
30 Baseline Survey. *BMC Public Health*. 2006;6.
31
32 41. Tuovinen EL, Saarni SE, Männistö S, et al. Smoking status and abdominal
33 obesity among normal- and overweight/obese adults: Population-based
34 FINRISK study. *Prev Med Rep*. 2016; 4:324–330.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 42. Flegal KM, Carroll MD, Kuczmarski RJ, et al. Overweight and obesity in the
4 United States: prevalence and trends, 1960–1994. *Int J Obes Relat Metab*
5 *Disord*; 1998; 22 (1):39–47.
6
7
8
9
10 43. Villareal DT, Apovian CM, Kushner RF, et al. Obesity in older adults: technical
11 review and position statement of the American Society for Nutrition and NAASO,
12 The Obesity Society, *The American Journal of Clinical Nutrition*, 2005 82 (5),
13 923–934
14
15
16
17
18
19
20 44. Sperrin M, Marshall AD, Higgins V. et al. Body mass index relates weight to
21 height differently in women and older adults: serial cross-sectional surveys in
22 England (1992-2011). *J Public Health (Oxf)*. 2016;38(3):607-613.
23
24
25
26
27 45. Vazquez G, Duval S, Jacobs DR, et al. Comparison of Body Mass Index, Waist
28 Circumference, and Waist/Hip Ratio in Predicting Incident Diabetes: A Meta-
29 Analysis, *Epidemiologic Reviews*, 2007; 29(1).115–128.
30
31
32
33
34 46. Emerging Risk Factors Collaboration, Wormser D, Kaptoge S, et al. Separate
35 and combined associations of body-mass index and abdominal adiposity with
36 cardiovascular disease: collaborative analysis of 58 prospective studies. *Lancet*.
37 2011;377(9771):1085-95.
38
39
40
41
42
43
44 47. Morio B, Barra V, Ritz P. et al. Benefit of endurance training in elderly people
45 over a short period is reversible. *Eur J Appl Physiol* 2000; 81(1): 329.
46
47
48
49 48. Fletcher J, Frisvold D. Higher education and health investments: Does more
50 schooling affect preventive health care use. *Journal Capital*. 2009; 3(2):144–
51 176.
52
53
54
55
56 49. Cohen AK, Rai M, Rehkopf DH, et al. Educational attainment and obesity: a
57 systematic review. *Obes Rev*. 2013;14(12):989-1005.
58
59
60

- 1
2
3 50. Devaux M, Sassi F, Church J, et al. (2011), "Exploring the Relationship between
4 Education and Obesity", *OECD Journal: Economic Studies*, Vol. 2011/1.
5
6
7
8 51. Traversy G, Jean-Philippe C. Alcohol Consumption and Obesity: An Update,
9
10 *Curr Obes Rep.* 2015; 4 (1):122–130.
11
12
13 52. Ford ND, Patel SA, Narayan KM. Obesity in Low- and Middle-Income Countries:
14
15 Burden, Drivers, and Emerging Challenges. *Annu Rev Public Health.*
16
17 2017;20(38):145-164.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

Title and abstract	Item No	Recommendation
	1	<p>(a) Socioeconomic and behavioural determinants of overweight/obesity among adults in Botswana: A cross-sectional study</p>
		<p>(b) Objective</p>
		<p>To undertake a comprehensive assessment of socioeconomic and behavioural determinants of overweight/obesity among adult population in Botswana.</p>
		<p>Design</p>
		<p>The study adopted a cross-sectional design by selecting adult respondents in three cities and towns, fifteen urban villages and fifteen rural areas across Botswana using a multistage probability sampling technique.</p>
		<p>Setting</p>
		<p>The study was conducted in selected rural and urban areas of Botswana.</p>
		<p>Participants</p>
		<p>The study sample consisted of 1178 adult males and females aged 15 years and above.</p>
		<p>Primary Outcome Measures</p>
		<p>Objectively measured overweight/obesity.</p>
		<p>Results</p>
		<p>Prevalence of overweight/obesity in the study population was estimated at 41%. The adjusted odds ratio (AOR) of overweight/obesity were highest among women (AOR=2.74, 95% C.I. =1.92-3.90), in ages 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), among individuals with secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary education (AOR=1.99, 95% C.I. =1.16-3.38), smokers (AOR=2.16, 95% C.I. = 1.22-3.83) and people with poor physically activity (AOR=1.46, 95% C.I. =1.03-3.24). These were statistically significant at 5% level.</p>
		<p>Conclusion</p>
		<p>Being female, elderly, of high education levels, a smoker and having poor physical activity were identified as key factors for prevalence of overweight/obesity. These findings suggest the need for broad based strategies encouraging physical activity among different socioeconomic groups.</p>
		<p>Key words</p>
		<p>Overweight, Obesity, socioeconomic, behavioural, Botswana</p>
		<p>Strengths and Limitations of the study</p>
		<ul style="list-style-type: none"> ○ To our knowledge this is the first cross-sectional study in Botswana to comprehensively assess both the socioeconomic and behavioural determinants of overweight/obesity among adults.
		<ul style="list-style-type: none"> ○ Data were collected from a large and randomly selected representative population. The data contained information on potential confounding factors, with a low proportion of missing information making the study more comparable.
		<ul style="list-style-type: none"> ○ As the cross sectional design was employed, the causal relationship between an explanatory and an outcome variable was not established.

Introduction

Background/rationale	2	The Botswana WHO STEPS surveys conducted in 2007 and 2014 have shown increasing levels of overweight/obesity in the general adult population of Botswana. Prevalence of overweight/obesity was 18.7% in 2007 and increased to 38.6 % in 2014. Despite evidence of the high prevalence of overweight/obesity in Botswana, there is little evidence of socioeconomic and behavioural determinants of overweight/obesity. To the best of our knowledge this is the first cross sectional study which takes into account both socioeconomic and behavioural factors for understanding the heterogeneity of overweight/obesity in Botswana. The reduction of overweight/obesity prevalence should be a key goal of health-care policy because of its association with many NCDs.
Objectives	3	The main objective of this study was to assess socioeconomic and behavioural determinants associated with overweight/obesity prevalence in Botswana.
Methods		
Study design	4	The study used cross sectional design, and employed stratified multistage probability sampling technique to select respondents in three cities and towns, fifteen urban villages and fifteen rural areas across Botswana.
Setting	5	Data was collected in March 2016, in cities/towns, urban and rural villages of Botswana.
Participants	6	A total sample of 1178 males and females aged 15 years were selected and included as sample for the survey using the Kish Grid Technique.
Variables	7	Overweight/obesity was the outcome variables, other NCDs risk factors such as alcohol consumption, smoking, poor fruit and vegetable consumption, and poor physical activity. Other socioeconomic covaraiates such as sex, age, work status, residence, education and marital status were used as controls.
Data sources/ measurement	8*	N/A
Bias	9	Validity of data collection tool was done through pretest prior to the survey
Study size	10	The sample size of the study was calculated using the sample size calculator and yielding a sample size of 1280. The sample size is powered to produce estimates for Botswana and for broad socioeconomic categories. The final selection of respondents for inclusion in the study was that 28.5% of the respondents were recruited from cities and towns, 47.3% from urban villages and 24.1% from rural villages. This is proportionate to the size of the population as estimated during the 2011 Botswana Population and Housing Census. Hence the total respondents from towns and cities= $28.5/100*1280=365$ respondents; $47.3/100*1280=606$ respondents from urban villages, and a total of $24.1/100*1280=309$ respondents from rural villages were targeted. Furthermore, the selection of the sample within each enumeration area is calculated in relation to the proportion of population in that district. E.g. for Gaborone = $178654/226649*365=288$ (see table1).

Table 1: Summary of sampled districts with their enumeration areas and proportion of study respondents.

Proposed Localities	Population aged 15-65 years (2011 Census)	Proposed Number of EA's	Proposed Number of Population Aged 15-65 years
---------------------	---	-------------------------	--

Cities and Towns (324291)	226049	18	365
Gaborone	178654	14	288
Selebi-Phikwe	34093	3	55
Jwaneng	13302	1	21
Urban Villages (538585)	271900	30	606
Kanye	31108	3	69
Letlhakeng	5439	1	12
Maitengwe	2969	0	7
Maun	40796	5	91
Mochudi	29331	3	65
Molepolole	43103	5	96
Serowe	35614	4	79
Tsabong	6131	1	14
Tlokeng	28211	3	63
Tutume	10626	1	24
Kopong	6856	1	15
Mmadimare	9192	1	20
Lerala	3943	0	9
Gumare	5180	1	12
Thamaga	13401	1	30
Rural Villages (274363)	21702	15	309
Nthanthe	1321	1	19
Ditshegwane	1209	1	17
Senete	1415	1	20
Sehithwa	2928	2	42
Mathubudukwane	1233	1	18
Serinane	358	0	5
Moiyabana	3014	2	43
Werda	1905	1	27
Dikwididi	225	0	3
Makuta	464	0	7
Kgope	521	0	7
Tobane	1455	1	21
Maunatlala	2871	2	41
Etsha 6	2783	2	40
Kubung	188	0	3

For each selected EA, 20 households were selected using systematic sampling method. This followed guidelines used in most Demographic Health Surveys where 20-25 households (hhs) were selected from the primary sampling units (PSUs). For instance, in the case of cities and town; $365/20=18$ EAs. The above procedure was followed for all districts, where each of the sampled EAs, 20 households were selected using systematic sampling method. The Kish grid was used to select the eligible respondents

from the selected households. Thus, once a household is selected, the interviewer created a listing (sampling frame) of all the persons in the household that are eligible for the interview process. This listing includes the name of the person, their gender, their relationship to the head of the household and their age. Once the listing was done, each eligible member was assigned a unique number. Then using a randomized response table a particular member was chosen for the interview.

Quantitative variables	11	Refer to 7 above
Statistical methods	12	<p>(a) Key two methods of data analysis were employed in the study: Bivariate analysis and logistic regression analysis to assess determinants of overweight/obesity</p> <p>(b) An analysis of the interaction effects was done before variables were included for analysis in logistic regression model.</p> <p>(c) None of the variables used in this study had a missing rate of more than 10%. This was within the cut-off from the literature regarding an acceptable percentage of missing data in a data set for valid statistical inferences (see, Schaffer 1999 & Bennett 2001 for example)</p> <p>(d) For sampling strategy, refer to attached file on the NCD study for methodology</p> <p>(e) N./A</p>
Results		
Participants	13*	<p>(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed-N/A</p> <p>(b) Give reasons for non-participation at each stage-N/A</p> <p>(c) Consider use of a flow diagram-N/A</p>
Descriptive data	14*	<p>(a) A total of 1,178 respondents between the ages of 15 years and above were successfully interviewed. The sample constituted a high proportion of females (69.1%) than males (30.9%). The sample age distribution suggests a relatively young population, with over half (59%) of the sample being less than 39 years of age, and almost three quarters (73.5%) being less than fifty years of age. As age increases, the sample skewed in favour of females.</p> <p>More than two fifths (45.4%) of the population resided in urban villages; while the remaining proportion resided in cities and towns (30.2%), and rural areas (24.5%). Almost three quarters (73.8%) of respondents were never married; over a third (35.5%) had primary education or less; over a quarter (27.2%) had junior secondary education while just under a fifth had senior secondary education (17.3%) and tertiary education (19.9%). Close to two fifth (37.5%) in the sample were not employed; over a quarter were employed in either the public or private sector (26.2%) while the remaining proportion were either self-employed (11.2%) , home makers or students (18.2%) and retirees (6.4%).</p> <p>(b) Indicate number of participants with missing data for each variable of interest-N/A</p>
Outcome data	15*	Report numbers of outcome events or summary measures-N/A
Main results	16	(a) It was observed that gender had a significant association with overweight/obesity when controlling for both socioeconomic and behavioural factors. Women were observed to have more than 2 times (AOR=2.74, 95% C.I. =1.92-3.90) greater odds for overweight/obesity than men.

The odds of being overweight/obese increased with age after controlling for socioeconomic and behavioural factors and were highest among individuals aged 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), but declined at ages 65+ years (AOR=2.88, 95% C.I. = 1.21-6.86). Education was also found to be a significant factor associated with overweight/obesity in the study population after controlling for socioeconomic covariates. Further, individuals who had secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary or higher education (AOR=1.99, 95% C.I. =1.16-3.38) were found to have higher probability of being overweight/obese than individuals with primary or less education. Meanwhile, there was no significant relationship observed between work status, wealth status and overweight/obesity in the general study population.

The association between smoking and alcohol consumption with overweight/obesity was found to be negative in the crude model. However, controlling for other socioeconomic and demographic factors, the odds of overweight/obesity increased two times (AOR=2.16, 95% C.I. = 1.22-3.83) among smokers than non-smokers. Overweight/obesity was also significantly associated with poor physical activity (AOR=1.46, 95% C.I. =1.03-3.24) after adjusting for socioeconomic and behavioural factors.

(b) Report category boundaries when continuous variables were categorized-N/A

(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period-N/A

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses-N/A
Discussion		
Key results	18	Key socioeconomic factors associated with overweight/obesity are being a woman, older, and with higher education (secondary and tertiary level education). Behavioral factors such as smoking and poor physical activity were found to be positively associated with overweight/obesity. These results demonstrate that socioeconomic and behavioural factors play an important role in prevalence of overweight/obesity in Botswana. Botswana's multisectoral national strategic plan implementation framework for combating NCDs should focus on developing a community based and people centered awareness about the harmful impact of smoking, alcohol, physical inactivity and poor quality of diet facilitated by comprehensive national monitoring and evaluation framework.
Limitations	19	The study employed a cross-sectional study design and the main limitation is that because data on each participant was recorded only once it was difficult to infer the temporal association between an explanatory and an outcome variable. Therefore, only an association, and not causation, can be inferred from this study.
Interpretation	20	Limitations of self-reported data such as under and over-reporting were borne in mind when interpreting results of the study
Generalisability	21	The NCDs study sample was not designed to be representative of Botswana, and we are cautious about generalizing these study findings
Other information		
Funding	22	The NCDs study was funded by Office of Research and Development at University of Botswana. However, the funder does not have any influence or say in the present article.

1
2 *Give information separately for exposed and unexposed groups.
3

4 **Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and
5 published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely
6 available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at
7 <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is
8 available at www.strobe-statement.org.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

Socioeconomic and behavioural determinants of overweight/obesity among adults in Botswana: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029570.R2
Article Type:	Original research
Date Submitted by the Author:	11-Sep-2019
Complete List of Authors:	Keetile, Mpho; University of Botswana, Population Studies Navaneetham, Kannan; University of Botswana, Department of Population Studies Letamo, Gobopamang; University of Botswana, Population Studies Bainame, Kenabetsho; University of Botswana, Population Studies Rakgoasi, Serai; University of Botswana, Population Studies Gabaitiri, Lesego; University of Botswana, Statistics Masupe, Tiny; University of Botswana Faculty of Health Sciences, School of Medicine Molebatsi, Robert; University of Botswana, Sociology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Global health
Keywords:	Overweight, Obesity, Socioeconomic, Behavioural, Botswana

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Socioeconomic and behavioural determinants of overweight/obesity**
6 **among adults in Botswana: A cross-sectional study**
7
8

9 Mpho Keetile¹, Kannan Navaneetham¹, Gobopamang Letamo¹, Kenabetsho
10 Bainame¹, Serai Daniel Rakgoasi¹, Lesego Gabaitiri², Tiny Masupe³, Robert
11 Molebatsi⁴.
12
13

14
15 **Author Affiliations**
16

17 ¹Department of Population Studies, University of Botswana, Gaborone, Botswana

18 ²Department of Statistics, University of Botswana, Gaborone, Botswana

19 ³School of Medicine, University of Botswana, Gaborone, Botswana

20 ⁴Department of Sociology, University of Botswana, Gaborone
21
22
23
24

25 **Corresponding author**
26

27 Mpho Keetile, Department of Population Studies, Private Bag 00705, University of
28 Botswana, Gaborone, Botswana. email: mphokeet@yahoo.com. Tel: +26771231375
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Word count: 3871

Abstract

Objective

To undertake a comprehensive assessment of socioeconomic and behavioural determinants of overweight/obesity among adult population in Botswana.

Design

The study adopted a cross-sectional design by selecting adult respondents in three cities and towns, fifteen urban villages and fifteen rural areas across Botswana using a multistage probability sampling technique.

Setting

The study was conducted in selected rural and urban areas of Botswana.

Participants

The study sample consisted of 1178 adult males and females aged 15 years and above.

Primary Outcome Measures

Objectively measured overweight/obesity.

Results

Prevalence of overweight/obesity in the study population was estimated at 41%. The adjusted odds ratio (AOR) of overweight/obesity were highest among women (AOR=2.74, 95% C.I. =1.92-3.90), in ages 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6), among individuals with secondary (AOR=1.70, 95% C.I. = 1.11-2.61) and tertiary education (AOR=1.99, 95% C.I. =1.16-3.38), smokers (AOR=2.16, 95% C.I. = 1.22-3.83) and people with poor physically activity (AOR=1.46, 95% C.I. =1.03-3.24). These were statistically significant at 5% level.

Conclusion

Women, older adults, people with high education level, smokers and people who reported poor physical activity were found to have higher odds of being overweight/obesity. These findings suggest the need for broad based strategies encouraging physical activity among different socioeconomic groups.

Key words

Overweight, Obesity, socioeconomic, behavioural, Botswana

Strengths and Limitations of the study

- To our knowledge this is the first cross-sectional study in Botswana to comprehensively assess both the socioeconomic and behavioural determinants of overweight/obesity among adults.
- Data were collected from a large and randomly selected representative population. The data contained information on potential confounding factors, with a low proportion of missing information making the study more comparable.
- As the cross sectional design was employed, the causal relationship between an explanatory and an outcome variable was not established.

Introduction

The disease profile in Botswana is changing, with most deaths and disability in the foreseeable future likely to be accounted for by non-communicable diseases (NCDs). Children, adults and the elderly are all vulnerable to the risk factors that contribute to NCDs especially overweight/obesity, due to unhealthy diets, low physical activity, exposure to tobacco smoke and/or the effects of the harmful use of alcohol¹⁻³. The key drivers of NCDs are ageing, rapid urbanisation and the globalisation of unhealthy lifestyles. Globalization of unhealthy lifestyles like unhealthy diets may show up in individuals as raised blood pressure, increased blood glucose, elevated blood lipids, and overweight/ obesity¹.

Several studies have recently shown that many low income and middle-income countries (LMICs) are experiencing an increase in the prevalence of overweight/obesity⁴⁻⁶. This prevalence is predicted to continue to increase in the future. Much of the increase in prevalence levels for overweight/obesity is expected in countries such as India, some parts of South-East Asia, China, most of South America and some parts of Sub Saharan Africa (SSA)⁷⁻⁸. Until recently, SSA was minimally affected by the overweight/obesity epidemic due to under-nutrition and a major burden of Human Immunodeficiency Virus (HIV) and tuberculosis (TB)⁹. Meanwhile research on overweight/obesity has started to gain significance in SSA¹⁰⁻¹⁴ due to its increasing magnitude.

It has been shown by previous studies that socioeconomic and behavioural factors have a significant association with overweight/obesity¹⁵⁻¹⁶. These studies have shown that socioeconomic and demographic factors such as sex, age, education level, work status, place of residence and wealth status are determinants of

1
2
3 overweight/obesity. Studies have found that women are more likely to be
4
5 overweight/obese than men, while the odds of being overweight/obesity has been
6
7 seen to increase with age¹⁷⁻¹⁹. It has also been observed that being employed
8
9 in a high income job, residing in urban areas, having high education and wealth
10
11 status are significantly associated with overweight/obesity²⁰.
12
13
14
15
16

17 The review of literature suggests significant association between a number of
18
19 behavioural factors and overweight/obesity. For example, significant positive
20
21 association was found between physical inactivity and overweight/obesity²¹. Further,
22
23 studies have noted that the association between physical inactivity and health
24
25 outcomes such as overweight/obesity is moderated by other lifestyle factors,
26
27 especially sedentary lifestyle^{15, 20, 21}. In most developing countries, there has been a
28
29 substantial shift from jobs with high-energy expenditure such as farming, mining and
30
31 forestry to more-sedentary sectors of manufacturing, services and office based-
32
33 work²². This shift has contributed to risky lifestyle behaviours condoning alcohol
34
35 consumption, poor diets and smoking²³⁻²⁴.
36
37
38
39
40
41

42 The Botswana World Health Organization (WHO) STEPS surveys conducted in 2007
43
44 and 2014 have shown rising levels of overweight/obesity among adult population of
45
46 Botswana. The estimated prevalence of overweight/obesity was 18.7% in 2007²⁵ and
47
48 increased to 38.6 % in 2014²⁶. Despite evidence of the high and rising prevalence of
49
50 overweight/obesity in Botswana, there is little evidence examining the association
51
52 between socioeconomic and behavioural factors, and overweight/obesity. To the
53
54 best of our knowledge, this is the first cross sectional study which takes into account
55
56 both socioeconomic and behavioural factors together for understanding the
57
58
59
60

1
2
3 determinants of overweight/obesity. The study would be significant in the context that
4
5 the reduction of overweight/obesity prevalence should be a key goal of health-care
6
7 policy because of its associated causes with many NCDs. Keeping this view, the
8
9 main aim of this study was to examine the association between socioeconomic and
10
11 behavioural factors, and overweight/obesity in Botswana.
12
13
14
15
16

17 **Methods**

18 ***Sampling, recruitment and ethics***

19
20
21
22
23 The study used cross sectional design by selecting respondents in three cities and
24
25 towns, fifteen urban villages and fifteen rural areas across Botswana using a
26
27 multistage probability sampling technique. A list of districts, localities and
28
29 enumeration areas (EAs) together with their households was prepared based on the
30
31 2011 Botswana Population and Housing census that form a sampling frame. At the
32
33 first stage, population was stratified into cities and towns, urban villages and rural
34
35 settlements. A listing of all 26 census districts in each stratum was made and from
36
37 these districts a total of all 4845 EAs were listed for cities and towns, urban villages
38
39 and rural settlements. At the second stage, localities in cities and towns, urban
40
41 villages and rural villages were randomly selected. At the third and fourth stage a
42
43 random selection of EAs and households, respectively was made. Lastly, individuals
44
45 aged 15 years and over were selected for interview from the list of households with
46
47 persons 15 years and over. The NCD study questionnaire was developed using the
48
49 instruments from the WHO Study on Global Ageing and Adult Health (SAGE), and
50
51 WHO STEPS Surveys.
52
53
54
55
56
57
58
59
60

1
2
3 All ethical clearance procedures were followed before the start of the study. The
4 study proposal along with the necessary documents were submitted to and approved
5 by the Institutional Review Board of the University of Botswana. Approval for
6 conducting the study was also obtained from the Government of Botswana through
7 the Ministry of Health and Wellness (MoHW).
8
9
10
11
12
13
14

15 As part of the study protocol, all respondents were requested to sign a consent form
16 and were told that participation in the study was voluntary. For participants aged 15-
17 17 years written consent was obtained from the parents or guardians. All individuals
18 aged 15 years and above who had successfully completed the NCD study
19 questionnaire were included in the analysis for this study. Privacy and confidentiality
20 of highest standard were maintained by treating all respondents as anonymous and
21 no names of respondents are mentioned or implied when presenting findings of the
22 study.
23
24
25
26
27
28
29
30
31
32
33

34 ***Definitions and Measurement of variables***

35
36
37 The outcome variable for the study is overweight/obesity. Body Mass Index (BMI)
38 was used to classify overweight/obesity. BMI was derived from weight and height:
39 $\text{weight (kg) / (height (m) x height (m))}^1$. The Charder MS7301 250Kg digital scale
40 and the Muac measuring tape were used for anthropometric measurements. Weight
41 was measured, to the nearest 0.1kg, while height was measured in metres. BMI was
42 categorized into: underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), normal weight ($18.5 \leq \text{BMI} < 25$
43 kg/m^2), overweight ($25 \leq \text{BMI} < 30 \text{ kg/m}^2$) and obese ($\text{BMI} \geq 30 \text{ kg/m}^2$)¹. Overweight
44 and obese were used to create a binary outcome variable which was coded as:
45 being overweight/obese ($\text{BMI} \geq 25 \text{ kg/m}^2$) =1; not overweight/obese =0 ($\text{BMI} < 25$
46 kg/m^2).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Information on NCD risk factors was collected through self-reports. Respondents
4 were asked questions on NCD risk factors such as tobacco use, alcohol
5 consumption, physical activity and fruit and vegetable consumption. For tobacco use,
6 respondents were asked whether they currently smoke tobacco products and their
7 response was coded as yes=1 and no=0. Alcohol consumption was measured based
8 on the intensity of alcohol consumed in the past 30 days. Respondents who had
9 consumed alcohol in the past 30 days were asked about the number of standard
10 alcohol drinks they had each day in the past 7 days and if they reported to have had
11 three or more drinks per day (of approximately 60g alcohol) it was considered to be
12 excessive drinking=1 and 0=otherwise.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Based on WHO recommendations poor physical activity was calculated using an
27 average typical types of activity undertaken¹³. It was calculated based on the time
28 taken while doing physical activity in the past 7 days. Respondents were asked
29 whether they do any moderate to rigorous intensity activities for at least 10 minutes
30 continuously. This was considered for the domains of work and walking (includes at
31 work and at home, walking to travel from place to place, and any other walking for
32 recreation, sport, exercise, or leisure). Based on the time taken on work and walking
33 the respondents were grouped into four categories of: no activity, low, moderate and
34 high activity levels to show the intensity of their physical activity. The resultant
35 variable was coded such that no and low activity (<10 minutes of physical activity) =1
36 and moderate and high activity (≥10 minutes of physical activity) were coded=0.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Poor fruit and vegetables consumption was created when an individual reported daily
53 consumption of less than the recommended 5 servings of fruits and vegetables.
54 Respondents reported the number of servings for fruits/vegetables they had in a
55
56
57
58
59
60

1
2
3 typical day, and if the servings were less than 5 in a day, they were considered to be
4
5 having poor fruit/vegetable consumption^{13, 27, 28}.
6
7

8 A wealth index (WI) was constructed to be a measure of wealth status. WI is a
9
10 composite measure of ownership of consumer durables, housing characteristics, and
11
12 access to public services¹⁶. Information on a range of durable assets was collected
13
14 during the survey (e. g. car, refrigerator, television), housing characteristics (e. g.
15
16 material of dwelling floor and roof, main cooking fuel), access to basic services (e. g.
17
18 electricity supply, source of drinking water, sanitation facilities) and ownership of
19
20 livestock (e.g. cattle, goats, sheep, horses, chickens). Further to collection of
21
22 information on durable assets, information on ownership of land was also collected.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Principal component analysis was employed to derive the wealth index variable,
which had five categories from the 1st to the 5th quintile (poorest to richest).

Control variables

Age and place of residence were used as control variables. These variables were conceptualized to have an association with the outcome variable. Therefore, as control variables they are included in the net effects model, so that the association between the independent variables of interest becomes isolated and discernible.

Statistical analysis

Descriptive and multivariable logistic analysis were used. Descriptive analysis provided bivariate association of explanatory and outcome variable for further exploration of key determinants of overweight and obesity in Botswana. In the multivariable binary logistic regression, the outcome variable was coded as '1' if respondents were overweight/obese and '0' if they were not overweight/obese. Results of logistic regression analysis were presented together with adjusted odd

1
2
3 ratios (AOR), 95% confidence intervals (CI) and 5% statistical significance level. Chi-
4 square test was used for bivariate analysis. Data analysis was done using SPSS
5 version 25 program. Since this was a population based study representative of the
6 population, selection bias was minimal and missing cases were few. In order to
7 control for cluster effects, complex sample module in SPSS has been used.
8
9
10
11
12
13

14 ***Patient and Public involvement***

15
16
17
18 No patients were involved in developing the research question, outcome measures
19 and overall design of the study.
20
21
22

23 **Results**

24 ***Sample characteristics***

25
26
27
28
29
30 Table 1 presents socioeconomic characteristics of the respondents. The sample
31 constituted a high proportion of females (69.1%) than males (30.9%). The sample
32 age distribution suggests a relatively young population, with over half (59%) of the
33 sample being less than 39 years of age, and almost three quarters (73.5%) being
34 less than fifty years of age.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Socio demographic characteristics

Variable	N	%
Sex		
Male	364	30.9
Female	813	69.1
Total	1177	100
Age in years		
<24	270	26.4
25– 34	302	29.5
35 – 44	196	19.2
45– 54	130	12.7
55 – 64	75	7.3
65+ years	50	4.9
Total	1023	100
Locality Type		
Cities/Towns	355	30.2
Urban Villages	534	45.4
Rural Settlements	288	24.5
Total	1177	100
Marital Status		
Never Married	864	73.8
Currently married	199	17.0
Formerly married	108	9.2
Total	1171	100
Highest Level of Education Attained		
Primary or Less	410	35.5
Junior Secondary	314	27.2
Senior Secondary	200	17.3
Tertiary & Over	230	19.9
Total	1154	100
Work Status in past 12 months		
Public Sector	122	10.5
Private Sector	182	15.7
Self Employed	130	11.2
Not Employed	436	37.5
Homemaker-Student	218	18.8
Retired-Other	74	6.4
Total	1162	100
Wealth status		
Lowest	234	19.9
Second	237	20.1
Middle	235	19.9
Fourth	237	20.1
Highest	235	19.9

Total	1178	100
--------------	-------------	------------

More than two fifths (45.4%) of the population resided in urban villages; while one – third of them resided in cities and towns (30.2%), and a quarter of them lived in rural areas (24.5%). Almost three-fourth (73.8%) of respondents was never married; over a third (35.5%) had primary education or less; over a quarter (27.2%) had junior secondary education while just under a fifth had senior secondary education (17.3%) and tertiary education (19.9%). Close to two fifth (37.5%) respondents were not employed; over a quarter were employed in either the public or private sector (26.2%).

Table 2 shows behavioural characteristics of the study population. It shows that prevalence of alcohol consumption was estimated at 17.3% while tobacco smoking was estimated at 11.5%. Poor physical activity was also high in the study population with 48.9% of respondents indicating poor physical activity. Poor fruit and vegetable intake was estimated at 82.5%.

Table 2: Behavioural characteristics of the study population

Variable	N	%
Smoking		
Yes	136	11.5
No	1042	88.5
Total	1178	100
Alcohol consumption		
Yes	204	17.3
No	974	82.7
Total	1178	100
Poor physical activity		
Yes	576	48.9
No	602	51.1
Total	1178	100
Poor fruit/vegetable intake		
Yes	1045	82.5
No	133	17.5
Total	1178	100

Prevalence of overweight/obesity

Estimated prevalence of overweight/obesity in the study population was 41.3% (Table 3). There were significant gender differences in prevalence of overweight/obesity. It was found that women than men (48.6% among women, 25.5% among men) were overweight/obese. Individuals residing in rural areas (46.8%), public sector employees (55.2%) and those with low education (45%) showed high prevalence of overweight/obesity. Meanwhile it was found that wealth status was not significantly associated with overweight/obesity. Among the

1
2
3 behavioural factors, the prevalence of overweight/obesity was greater among non-
4 smoker (42.9%). Alcohol consumption, poor physical activity and poor fruits and
5 vegetable consumption did not show any significant association with
6
7
8
9
10 overweight/obesity.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 3: Prevalence of overweight/obesity by socioeconomic and behavioural characteristics of the study population

Variable	N	%	P-value
Sex			
Male	364	25.5	≤0.001**
Female	813	48.6	
Age			≤0.001**
≤24	270	21.8	
25-34	302	36.8	
35-44	196	53.0	
45-54	130	54.5	
55-64	75	61.8	
65+	50	43.8	
Marital status			≤0.001**
Never-married	864	35.5	
Currently-married	199	62.2	
Formerly-married	108	54.1	
Education			0.019**
Primary or less	410	45.4	
Secondary	514	36.7	
Tertiary or higher	230	44.6	
Residence			0.008**
Cities and towns	355	37.9	
Urban villages	534	41.0	
Rural villages	288	46.8	
Work status			≤0.001**
Public sector	122	52.2	
Private sector	182	36.8	
self-employed	130	55.0	
Not employed	436	45.8	
Home-maker/student	218	29.4	
Retired/other	74	40.3	
Wealth status			0.765
Lowest	234	38.8	
second	237	43.2	

middle	235	41.1	
Fourth	237	39.7	
Highest	235	44.1	
Alcohol consumption			0.445
No	766	33.3	
yes	412	37.7	
Smoking			0.004**
Yes	136	29.7	
No	1042	42.9	
Poor physical activity			0.547
Yes	376	41.5	
No	771	39.6	
Poor fruit/vegetable consumption			0.832
No	133	41.9	
yes	1045	43.0	
Overall	1178	41.3	

Note: **Chi-square test statistically significant at $p < 0.05$.

Determinants of overweight/obesity

Three models were run to assess determinants of overweight/obesity (Table 4). Model I is the crude model and it assessed the association between overweight/obesity and each of the socioeconomic and demographic variables and behavioural factors independently, Model II assessed the association between overweight/obesity adjusting for socioeconomic factors only, while Model III examined the association between overweight/obesity adjusting for both socioeconomic and behavioural factors. It was observed that gender had a significant association with overweight/obesity when controlling for both socioeconomic and behavioural factors. Women were observed to have more than 2 times (AOR=2.74, 95% C.I. =1.92-3.90) greater odds for overweight/obesity than men.

1
2
3
4
5
6 The odds of being overweight/obese increased with age after controlling for
7 socioeconomic and behavioural factors and were highest among individuals aged
8 55-64 years (AOR=5.53, 95% C.I. = 2.62-11.6). However, it declined at ages 65+
9 years (AOR=2.88, 95% C.I. = 1.21-6.86). Education was also found to be a
10 significant factor associated with overweight/obesity when controlling for
11 socioeconomic and behavioural factors. Individuals who had secondary (AOR=1.70,
12 95% C.I. = 1.11-2.61) and tertiary or higher education (AOR=1.99, 95% C.I. =1.16-
13 3.38) were found to have higher probability of being overweight/obese than
14 individuals with primary or less education. Work status and wealth status were not
15 significantly associated with overweight/obesity.
16
17
18
19
20
21
22
23
24
25
26
27
28

29 The association between smoking and alcohol consumption with overweight/obesity
30 was found to be negative in the crude model. However, when controlling for other
31 socioeconomic and demographic factors, the odds of overweight/obesity increased
32 two times (AOR=2.16, 95% C.I. = 1.22-3.83) among smokers than non-smokers.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Table 4: Odds ratios (OR) from multivariable logistic regression for overweight/obesity among adults in the study population, Botswana.

Variable	Model I		Model II		Model III	
	OR	C.I.	OR	C.I.	OR	C.I.
Sex						
Male	1.00		1.00		1.00	
Female	2.75**	(2.08-3.64)	3.01**	(2.12-4.26)	2.74**	(1.92-3.90)
Age						
≤24	1.00		1.00		1.00	
25-34	2.08**	(1.42-3.03)	1.77**	(1.12-2.77)	1.83**	(1.17-2.88)
35-44	4.02**	(2.67-6.08)	3.70**	(2.23-6.12)	3.87**	(2.34-6.42)
45-54	4.28**	(2.70-6.78)	3.04**	(1.66-5.55)	3.26**	(1.78-5.98)
55-64	5.78**	(3.26-10.2)	4.91**	(2.35-10.2)	5.53**	(2.62-11.6)
65+	2.78**	(1.46-5.28)	2.74**	(1.15-6.50)	2.88**	(1.21-6.86)
Marital status						
Never-married	0.46**	(0.30-0.71)	0.68	(0.34-1.34)	0.66	(0.33-1.32)
Currently-married	1.39	(0.84-2.29)	1.34	(0.66-2.71)	1.28	(0.63-2.61)
Formerly-married	1.00		1.00		1.00	
Education						
Primary or less	1.00		1.00		1.00	
Secondary	0.69**	(0.53-0.91)	0.49**	(0.29-0.83)	1.70**	(1.11-2.61)
Tertiary or higher	0.96	(0.69-1.35)	0.87	(0.58-1.32)	1.99**	(1.16-3.38)
Residence						
Cities and towns	1.00		1.00		1.00	
Urban villages	1.13	(0.86-1.50)	1.06	(0.74-1.50)	1.07	(0.75-1.52)
Rural villages	1.44**	(1.04-1.99)	1.37	(0.88-2.12)	1.37	(0.88-2.12)
Work status						
Public sector	1.61	(0.89-2.93)	1.30	(0.64-2.65)	1.22	(0.59-2.50)
Private sector	0.86	(0.49-1.51)	1.20	(0.61-2.36)	1.19	(0.60-2.35)
self-employed	1.18	(0.65-2.13)	1.24	(0.61-2.51)	1.29	(0.63-2.62)
Not employed	1.25	(0.75-2.08)	1.30	(0.70-2.40)	1.30	(0.70-2.41)
Home-maker/student	0.61	(0.35-1.07)	0.96	(0.48-1.92)	0.96	(0.48-1.92)
Retired/other	1.00		1.00		1.00	
Wealth status						
Lowest	0.80	(0.55-1.17)	0.60	(0.34-1.05)	1.03	(0.64-1.68)
second	0.96	(0.66-1.40)	0.67	(0.40-1.11)	1.12	(0.68-1.85)
middle	0.88	(0.60-1.28)	0.73	(0.45-1.19)	1.20	(0.71-2.01)
Fourth	0.83	(0.57-1.21)	0.79	(0.50-1.23)	1.53	(0.87-2.69)

Highest	1.00	1.00	1.00
Smoking		N.I	
Yes	0.56* (0.37-0.83)		2.16** (1.22-3.83)
No	1.00		1.00
Poor physical activity		N.I	
Yes	1.08 (0.83-1.40)		1.46** (1.03-3.24)
No	1.00		1.00
Poor fruit/vegetable consumption		N.I	
Yes	0.82 (0.55-1.22)		0.80 (0.28-2.24)
no	1.00		1.00
Alcohol consumption		N.I	
Yes	0.66* (0.47-0.91)		1.23 (0.53-2.83)
No	1.00		1.00

N.I- not included, Model I: Crude model; Model II: socioeconomic variables included; Model III: socioeconomic + behavioural characteristics included. **statistically significant at $p < 0.05$.

Discussion

The prevalence of overweight/obesity in the study population was 41.3%. Botswana also seems to have joined with other Southern African countries where the national prevalence rates ranges between 30% and 60% among adults age 15 years and over^{10, 26-31}. Nutrition transition from high-fibre, low-calorie diets rich in fruits and vegetables to refined, energy-dense foods high in fat, calories, sweeteners and salt¹² and low physical activity coupled with transition from agrarian to a sedentary industrial society contributes to a greater risk of overweight/obesity in Botswana³².

Gender differences in overweight/obesity found in this study is a well-established finding and is consistent with most previous studies³³⁻³⁴. Congruent with most studies in developing countries women were found to have higher odds of overweight/obesity than men³⁵⁻³⁷. This is contrary to findings from some developed countries such as United States (US) and United Kingdom (UK) where it was found that adult men tend to be overweight/obese than adult women³⁸⁻⁴². In developing countries such as Botswana, adult women are generally noted to be overweight/obese than adult men at all ages^{26, 35, 39-40}.

In the context of Botswana, gender differences in overweight/obesity can be explained by sociocultural dynamics and complex sociocultural pathways that encourage weight gain among women^{26, 40}. For example, cultural values favour larger body size among women as a sign of fertility, healthfulness, or prosperity in Botswana²⁶. In this study, women reported lower levels of physical activity which may also contribute to higher prevalence of overweight/obesity among them.

1
2
3 Overweight/obesity was highest among individuals aged 55-64 years but declined at
4 ages 65+ years. This finding corroborates data from large scale surveys⁴³⁻⁴⁶ which
5 show that mean body weight and BMI gradually increases during most of adult life
6 and reach peak values at 50–59 years of age in both men and women and after the
7 age of 60 years, mean body weight and BMI tend to decrease⁴⁷⁻⁴⁹. Similarly in South
8 Africa it was found that overweight/obesity tends to increase with age²¹.
9
10
11
12
13
14
15
16
17
18
19

20 Education differences were also observed for overweight/obesity. Individuals who
21 had high educational attainment had higher odds of overweight/obesity than those
22 with low education. This is consistent with other studies that have shown that
23 education is associated with overweight/obesity via socioeconomic status, literacy
24 and health behaviours⁵⁰⁻⁵².
25
26
27
28
29
30
31
32
33

34 There was no significant association observed between overweight/obesity and work
35 status and for wealth status. Other studies in neighbouring countries with
36 comparable Human Development Index (HDI) as Botswana including Namibia and
37 South Africa²⁷⁻²⁸ have also shown no wealth or work status differences in
38 overweight/obesity. There are several reasons to such scenario. These reasons
39 include; rapid and disorganized urbanization, increasing sedentary lifestyles, easy
40 access to and consumption of unhealthy food and high energy drinks among both
41 the poor and non-poor in these countries^{21, 27-28}. Further to this a general observation
42 has been made by Ford et al.⁵² that overweight/obesity is no longer an issue of
43 affluence in majority of low and middle income countries.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Further, the study noted that the odds of overweight/obesity were two times greater
4 among smokers than non-smokers. Tuovinen *et al.*⁴¹ based on literature review
5 observed that heavy smokers tend to have greater body weight than do light
6 smokers or non-smokers, which likely reflect a clustering of other risky behaviours
7 (e.g. low level of physical activity and poor diet) that are conducive to weight gain²³⁻
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
24. On the other hand, some studies have shown that smoking tends to suppress
appetite leading to low BMI among smokers^{10, 21, 30}.

Other important finding in this study is that overweight/obesity was significantly
associated with poor physical activity. This is consistent with other studies which
have shown that a decline in physical activity leads to adiposity⁵². The interpretation
of this finding is that Botswana has undergone a transition from an agrarian to
industrial society due to urbanization subsequently leading to industrial modernity.
As a result there is an increase in poor physical activity in the general population²⁴.
Poor physical activity leads to overweight/obesity through energy imbalance between
calories consumed and calories expended⁵². Generally formal physical activity that is
planned, structured, repetitive and purposeful is not common in Botswana. Thus in
order to increase physical activity and reduce overweight/obesity there is need for a
population-based, multi-sectoral, multi-disciplinary and culturally relevant approach.

This study benefits from data collected from a large and randomly selected sample
of respondents. The data contained information on potential confounding factors,
with a low proportion of missing information making the study more comparable.
Given that the recall period for behavioural risk factors such as smoking, alcohol
consumption, poor physical activity and poor fruit and vegetable consumption was

1
2
3 30 days or less there were minimal chances of recall bias. The main limitation is that
4
5 since the study used a cross sectional design, it was not possible to establish the
6
7 causal relationship between explanatory variables and overweight/obesity.
8
9
10 Furthermore, the NCDs study sample was not designed to be representative of the
11
12 whole of Botswana. However, the findings of this study give an indication of
13
14 emerging patterns of overweight/obesity in the country.
15
16

17 **Conclusion**

18
19
20 In conclusion our findings indicate high prevalence of overweight/obesity in the adult
21
22 population of Botswana. Overweight/obesity was found to be significantly high
23
24 among women, older adults and people with higher education levels. Behavioural
25
26 factors such as smoking and poor physical activity were found to be key correlates of
27
28 overweight/obesity. These results demonstrate that socioeconomic and behavioural
29
30 factors play an important role in prevalence of overweight/obesity. Botswana's
31
32 multisectoral national strategic plan implementation framework for combating NCDs
33
34 should focus on developing a community based and people centered awareness
35
36 about the harmful impact of smoking, alcohol, physically inactivity and poor quality of
37
38 diet facilitated by comprehensive national monitoring and evaluation framework.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

We wish to thank University of Botswana for affording us the time and resources to undertake this study. This paper is a part of the larger study on “Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course (NCD study)” funded by the Office of Research of Development (ORD) at the University of Botswana. The authors wish to thank ORD for the financial support. We also wish to thank the study participants.

Contributors

This paper is a part of the larger NCD study which was designed, conceived and developed by the study team (MK, KN, GL, KB, SDR, LG, TM and RM). MK did the analysis and wrote the initial draft of this paper. KN, GL and SDR gave feedbacks for revising the first draft. MK, KN and GL worked on the reviewers comments. All authors have reviewed and approved the final version of the manuscript except KB who recently passed away. We dedicate this article to KB whom we miss very much.

Funding

The ‘Chronic Non communicable Diseases in Botswana: Chronic disease prevalence, Health Care Utilization, Health Expenditure and the Life course’ survey was supported by Office of Research and Development, through Round 28 Research Funding, University of Botswana.

Disclaimer

The funding body had no role in the design of the study, analyses, interpretation or decision to submit the manuscript for publication.

Competing Interests

Nil

Participant consent

Obtained

Ethics Approval

All ethical clearance formalities were completed before the start of the study. The study proposal along with the necessary documents was submitted to and approved by the Institutional Review Board of the University of Botswana (Ref #: UBR/RES/IRB/1583) and the Ministry of Health and Wellness (Ref #: HPDME: 13/18/1 Vol. X (130)). Privacy and confidentiality of highest standard were maintained by treating all respondents as anonymous and no names of respondents are mentioned or implied when presenting findings of the study.

Data sharing statement

The dataset used and/or analysed during the current study is available from the corresponding author (Mpho Keetile) on reasonable request to the Office of Research and Development, University of Botswana.

References

1. World Health Organization. The global status of overweight and obesity. *Fact Sheet*. Geneva. 2016.
2. Musaiger AO, Al-Hazzaa HM. Prevalence and risk factors associated with nutrition-related noncommunicable diseases in the Eastern Mediterranean region. *Int J Gen Med*. 2012;5:199-217.
3. Baldwin W, Kaneda T, Amato L, *et al*. Noncommunicable diseases and youth: a critical window of opportunity for Latin America and the Caribbean. Washington, DC: *Population Reference Bureau*; 2013.
4. Bhurosy T, Jeewon R. Overweight and obesity epidemic in developing countries: a problem with diet, physical activity, or socioeconomic status?. *ScientificWorldJournal*. 2014;964236. doi:10.1155/2014/964236.
5. Ford ND, Patel SA, Narayan KMV. Obesity in Low- and Middle-Income Countries: Burden, Drivers, and Emerging Challenges. *Annu. Rev. Public Health*. 2017; 38:145–64.
6. Amugsi DA, Dimbuene ZT, Mberu B, *et al*. Prevalence and time trends in overweight and obesity among urban women: an analysis of demographic and health surveys data from 24 African countries, 1991–2014. *BMJ*;7:e017344. doi:10.1136/bmjopen-2017-017344.
7. Hruby A, Hu FB. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics*. 2015;33(7):673-89.
8. Mamun AA , Finlay JE. Shifting of under-nutrition to over-nutrition and its determinants among women of reproductive ages in the 36 low to medium income countries. *Obes Res Clin Pract*. 2015; 9: 75-86.

- 1
2
3 9. Micklesfield LK, Lambert EV, Hume DJ, *et al.* Socio-cultural, environmental
4 and behavioural determinants of obesity in black South African women.
5
6 *Cardiovasc J Afr.* 2013;24(9-10):369-75.
7
8
9
10 10. Biadgilign S, Mgutshini T, Haile D, *et al.* Epidemiology of obesity and
11 overweight in sub-Saharan Africa: a protocol for a systematic review and
12 meta-analysis. *BMJ Open.* 2017;7(11):e017666.
13
14
15
16
17 11. Ball GD, Mushquash AR, Keaschuk RA, *et al.* Using intervention mapping to
18 develop the Parents as Agents of Change (PAC) intervention for managing
19 pediatric obesity. *BMC Res Notes.* 2017;10:43.
20
21
22
23
24 12. Kelly AS, Barlow SE, Rao G, *et al.* Severe obesity in children and
25 adolescents: identification, associated health risks, and treatment
26 approaches: a scientific statement from the American Heart Association.
27
28 *Circulation.* 2013;128:1689–712.
29
30
31
32
33 13. World Health Organization (WHO). Obesity and overweight. *World Health*
34 *Organization, Geneva.* 2016.
35
36
37
38
39 14. Levine JA. Poverty and obesity in the U.S. *Diabetes.* 2011; 60:2667-2668.
40
41
42
43 15. Kim J, Sharma SV, Park SK. Association between socioeconomic status and
44 obesity in adults: evidence from the 2001 to 2009 Korea national health and
45 nutrition examination survey. *J Prev Med Public Health.* 2014;47(2):94-
46
47
48
49 103.16.
50
51
52 16. Vyas S, Kumaranayake L, Constructing socio-economic status indices: how
53 to use principal components analysis, *Health Policy and Planning.* 2006;
54
55
56 21(6), 459–468. <https://doi.org/10.1093/heapol/czl029>.
57
58
59
60

- 1
2
3 17. Maffetone PB, Laursen PB. The Prevalence of Overfat Adults and Children in
4 the US. *Front Public Health*. 2017;5:290.
5
6
7
8
9 18. Lumsden MA, Hor K. Impact of obesity on the health of women in midlife. *The*
10 *Obstetrician & Gynaecologist*. 2015; 17 (1):201–8.
11
12
13 19. Kim TJ, von dem Knesebeck O. Income and obesity: what is the direction of
14 the relationship? A systematic review and meta-analysis. *BMJ Open*.
15 2018;8(1):e019862.
16
17
18
19 20. Hong I, Coker-Bolt P, Anderson KR, *et al*. Relationship Between Physical
20 Activity and Overweight and Obesity in Children: Findings From the 2012 National
21 Health and Nutrition Examination Survey National Youth Fitness Survey. *Am J*
22 *Occup Ther*. 2016; 70(5):7005180060p1-8.
23
24
25
26 21. Popkin BM, Slining MM. New dynamics in global obesity facing low- and
27 middle-income countries. *Obesity Reviews: An Official Journal of the International*
28 *Association for the Study of Obesity*. 2013; 14(02), 11–20.
29 <http://doi.org/10.1111/obr.12102>.
30
31
32
33 22. Dare S, Mackay DF, Pell JP. Relationship between Smoking and Obesity: A
34 Cross-Sectional Study of 499,504 Middle-Aged Adults in the UK General
35 Population. *PLoS ONE*. 2015; 10(4).
36
37
38
39 23. Tuovinen EL, Saarni SE, Männistö S. *et al*. Smoking status and abdominal
40 obesity among normal- and overweight/obese adults: Population-based FINRISK
41 study. *Preventive Medicine Reports*. 2016;4(1), 324–330.
42
43
44
45 24. Keetile M, Letamo G, Navaneetham K. Patterns and determinants of
46 hypertension in Botswana, *J Public Health*. 2015; 23(1),311–318.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 25. Republic of Botswana. Chronic Disease Risk Surveillance Report 2010-
4
5 Ministry of Health & WHO. *Government Printers*. 2010.
6
7
8 26. Letamo G. The prevalence of, and factors associated with, overweight and
9 obesity in Botswana. *J Biosoc Sci*. 2011; 43(1):75-84.
10
11
12
13 27. Peltzer K, Phaswana-Mafuya N. Fruit and vegetable intake and associated
14 factors in older adults in South Africa. *Global Health Action*, 2012; 5(1), 1-8.
15 doi:10.3402/gha.v5i0.18668.
16
17
18 28. Nnyepi MS, Gwisai N, Lekgoa M, *et al*. Evidence of nutrition transition in
19 Southern Africa. *Proc Nutr*. 2015;17 (1):1–9.doi: 10.1017/S0029665115000051.
20
21
22
23 29. World Health Organization (WHO). Obesity and overweight. *World Health*
24 *Organization*, Geneva. 2015.
25
26
27
28 30. GBD 2015 Risk Factors Collaborators. Global, regional, and national
29 comparative risk assessment of 79 behavioural, environmental and occupational,
30 and metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the
31 Global Burden of Disease Study 2015. *Lancet*. 2016;388 (1):1659–724.
32
33
34
35
36
37
38 31. Agymang C, Boatemaa S, Frempong-Agymang G, *et al*. Obesity in Sub
39 Saharan Africa. Metabolic Syndrome. *Springer International Publishing*,
40 Switzerland. Ed. Rexford Ahima. 2015.41-53.
41
42
43
44
45 32. Amoah AG. Socio-demographic variations in obesity among Ghanain adults.
46 *Public Health Nutr*. 2003;6 (1):751-757.
47
48
49
50 33. Biadgilign S, Mgutshini T, Haile D, *et al*. Epidemiology of obesity and
51 overweight in sub-Saharan Africa: a protocol for a systematic review and meta-
52 analysis. *BMJ Open*. 2017;7(11).
53
54
55
56
57 34. Cohen AK, Rai M, Rehkopf DH, *et al*. Educational attainment and obesity: A
58 systematic review. *Obesity Reviews: An Official Journal of the International*
59
60

1
2
3 *Association for the Study of Obesity*, 2013; 14(12), 989–1005.
4
5 <http://doi.org/10.1111/obr.12062>.
6
7

8 35. Kanter R, Caballero B. Global Gender Disparities in Obesity: A Review of
9 American Society for Nutrition. *Adv. Nutr.* 2012;3 (1): 491–498.
10
11 doi:10.3945/an.112.002063.
12
13

14 36. Li L, Gower BA, Shelton RC, *et al.* Gender-Specific Relationship between
15 Obesity and Major Depression. *Front Endocrinol (Lausanne)*. 2017;8 (1):292.
16
17

18 37. Pinto KA, Griep RH, Rotenberg L, *et al.* Correction: Gender, time use and
19 overweight and obesity in adults: Results of the Brazilian Longitudinal Study of
20 Adult Health (ELSA-Brasil). *PLOS ONE* 2018;13(8): e0198380.
21
22
23

24 38. Luppino FS, de Wit LM, Bouvy PF, *et al.* Overweight, obesity, and
25 depression: a systematic review and meta-analysis of longitudinal studies. *Arch*
26 *Gen Psychiatry*. 2010; 67(1):220–9.
27
28

29 39. The GBD 2015 Obesity Collaborators. Health Effects of Overweight and
30 Obesity in 195 Countries over 25 Years. *N Engl J Med* 2017; 377:13-27.
31
32

33 40. Kamadjeu RM, Edwards R, Atanga JS, *et al.* Anthropometry measures and
34 prevalence of obesity in the urban adult population of Cameroon: an update from
35 the Cameroon Burden of Diabetes Baseline Survey. *BMC Public Health*. 2006;6:
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
228.

41. Tuovinen EL, Saarni SE, Männistö S, *et al.* Smoking status and abdominal
obesity among normal- and overweight/obese adults: Population-based FINRISK
study. *Prev Med Rep.* 2016; 4:324–330.

- 1
2
3 42. Flegal KM, Carroll MD, Kuczmarski RJ, *et al.* Overweight and obesity in the
4 United States: prevalence and trends, 1960–1994. *Int J Obes Relat Metab*
5 *Disord*; 1998; 22 (1):39–47.
6
7
8
9
10 43. Villareal DT, Apovian CM, Kushner RF, *et al.* Obesity in older adults:
11 technical review and position statement of the American Society for Nutrition and
12 NAASO, The Obesity Society, *The American Journal of Clinical Nutrition*. 2005 82
13 (5), 923–934
14
15
16
17
18
19
20 44. Sperrin M, Marshall AD, Higgins V, *et al.* Body mass index relates weight to
21 height differently in women and older adults: serial cross-sectional surveys in
22 England (1992-2011). *J Public Health (Oxf)*. 2016;38(3):607-613.
23
24
25
26
27 45. Vazquez G, Duval S, Jacobs DR, *et al.* Comparison of Body Mass Index,
28 Waist Circumference, and Waist/Hip Ratio in Predicting Incident Diabetes: A
29 Meta-Analysis, *Epidemiologic Reviews*. 2007; 29(1).115–128.
30
31
32
33
34 46. Emerging Risk Factors Collaboration, Wormser D, Kaptoge S, *et al.*
35 Separate and combined associations of body-mass index and abdominal
36 adiposity with cardiovascular disease: collaborative analysis of 58 prospective
37 studies. *Lancet*. 2011;377(9771):1085-95.
38
39
40
41
42
43
44 47. Morio B, Barra V, Ritz P. *et al.* Benefit of endurance training in elderly people
45 over a short period is reversible. *Eur J Appl Physiol* 2000; 81(1): 329.
46
47
48
49 48. Fletcher J, Frisvold D. Higher education and health investments: Does more
50 schooling affect preventive health care use. *Journal Capital*. 2009; 3(2):144–176.
51
52
53
54 49. Cohen AK, Rai M, Rehkopf DH, *et al.* Educational attainment and obesity: a
55 systematic review. *Obes Rev*. 2013;14(12):989-1005.
56
57
58
59
60

1
2
3 50. Devaux M, Sassi F, Church J, *et al.* “Exploring the Relationship between
4 Education and Obesity”, *OECD Journal: Economic Studies*. 2011(1):121-159.
5
6

7
8 51. Traversy G, Jean-Philippe C. Alcohol Consumption and Obesity: An Update,
9
10 *Curr Obes Rep*. 2015; 4 (1):122–130.
11

12
13 52. Ford ND, Patel SA, Narayan KM. Obesity in Low- and Middle-Income
14 Countries: Burden, Drivers, and Emerging Challenges. *Annu Rev Public Health*.
15
16 2017;20(38):145-164.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7-9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	7-9
Bias	9	Describe any efforts to address potential sources of bias	10
Study size	10	Explain how the study size was arrived at	
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	n/a
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9
		(b) Describe any methods used to examine subgroups and interactions	n/a
		(c) Explain how missing data were addressed	10
		(d) If applicable, describe analytical methods taking account of sampling strategy	10
		(e) Describe any sensitivity analyses	n/a
Results			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	11
		(b) Give reasons for non-participation at each stage	-
		(c) Consider use of a flow diagram	-
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	11
		(b) Indicate number of participants with missing data for each variable of interest	11
Outcome data	15*	Report numbers of outcome events or summary measures	-
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	16-19
		(b) Report category boundaries when continuous variables were categorized	-
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	-
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	-
Discussion			
Key results	18	Summarise key results with reference to study objectives	20-22
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	22-23
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	23
Generalisability	21	Discuss the generalisability (external validity) of the study results	23
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	24

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.