

Table 1. Eye Characteristics at Enrollment by Treatment Group

Characteristic	Active Eyes			Placebo Eyes		
	N	Mean	(SD)	N	Mean	(SD)
Tear osmolarity, mOsms/L	494	303.9	(17.2)	260	300.6	(14.5)
Keratograph tear break-up time, secs	351	8.4	(5.7)	190	7.6	(5.5)
Tear meniscus height, mm	351	0.38	(0.17)	190	0.38	(0.14)
Bulbar conjunctival redness score	351	1.20	(0.49)	190	1.12	(0.49)
		<u>(N=351)</u>			<u>(N=190)</u>	
Upper lid meibography-gland loss, n (%)						
Missing value	46	(13%)		23	(12%)	
None	73	(21%)		35	(18%)	
≤25%	99	(28%)		61	(32%)	
>25-50%	76	(22%)		38	(20%)	
>50-75%	39	(11%)		26	(14%)	
>75%	18	(5%)		7	(4%)	
Lower lid meibography-gland loss, n (%)						
Missing value	12	(3%)		7	(4%)	
None	55	(16%)		37	(19%)	
≤25%	124	(35%)		70	(37%)	
>25-50%	78	(22%)		41	(22%)	
>50-75%	60	(17%)		19	(10%)	
>75%	22	(6%)		16	(8%)	
		<u>(N=665)</u>			<u>(N=357)</u>	
MMP-9 result, n (%)						
Negative	439	(67%)		247	(70%)	
Positive	218	(33%)		108	(30%)	

SD is standard deviation