

Table 3: Changes in Eye Characteristics from Baseline by Treatment Group, Stratified by Baseline Level of the Characteristic

Baseline subgroup	Active Eyes			Placebo Eyes			Between Group Difference		p
	N	Mean	(SD)	N	Mean	(SD)	Mean	(95% CI)	
Tear osmolarity, mOsm/L									
<308 (N=457)	288	5.6	(17.2)	169	7.3	(15.9)	-1.7	(-5.1, 1.8)	0.34
≥308 (N=179)	136	-14.1	(23.5)	43	-11.2	(19.9)	-2.9	(-10.1, 4.3)	0.44
Interaction									0.76
Tear break-up time, secs									
<4.6 (N=137)	86	2.6	(4.8)	51	2.5	(3.9)	0.0	(-1.6, 1.7)	0.97
≥4.6, <9.2 (N=148)	97	1.1	(5.1)	51	0.6	(4.7)	0.5	(-1.2, 2.2)	0.56
≥9.2 (N=154)	107	-4.4	(7.4)	47	-5.0	(6.8)	0.7	(-1.6, 2.9)	0.57
Interaction									0.88
Tear meniscus height, mm									
<0.30 (N=149)	113	0.04	(0.13)	36	0.05	(0.14)	-0.01	(-0.08, 0.05)	0.68
≥0.30, <0.41 (N=162)	95	0.01	(0.11)	67	0.01	(0.12)	0.00	(-0.04, 0.04)	0.87
≥0.41 (N=168)	109	-0.05	(0.20)	59	-0.06	(0.15)	0.01	(-0.05, 0.07)	0.77
Interaction									0.87
Bulbar conjunctival redness score									
<1.00 (N=154)	96	0.08	(0.20)	58	0.07	(0.26)	0.01	(-0.07, 0.10)	0.81
≥1.00, <1.40 (N=124)	81	0.07	(0.32)	43	-0.03	(0.31)	0.09	(-0.02, 0.21)	0.11
≥1.40 (N=140)	96	-0.13	(0.55)	44	-0.11	(0.44)	-0.01	(-0.20, 0.17)	0.90
Interaction									0.39

SD is standard deviation
CI is confidence interval