

Caveolin-1 gene therapy inhibits inflammasome activation to protect from bleomycin-induced pulmonary fibrosis


Xin Lin¹, Michael Barravecchia¹, R. Matthew Kottmann², Patricia Sime² & David A Dean^{1*}

¹Department of Pediatrics, School of Medicine and Dentistry, University of Rochester, Rochester, New York 14642, USA. ²Department of Medicine, School of Medicine and Dentistry, University of Rochester, Rochester, New York 14642, USA


*Correspondence and request for reprint should be addressed to D.D. (email: david_dean@urmc.rochester.edu). Department of Pediatrics, University of Rochester, 601 Elmwood Avenue BOX 850, Rochester, NY 14642, USA

Supplementary Table S1. Primers used for RT-PCR

Name	Sense	Antisense
Mouse Cav-1	5'-CATCTCTACACTGTTCCCATCC-3'	5'- TGTGCGCGTCATACACT T-3'
Mouse GAPDH	5'-TGTGATGGGTGTGAACCACGAGAA-3'	5'- GAGCCCTTCCACAATGC CAAAGTT-3'
Human Collagen I	5'- CCCTAACCAAGGATGCACTATG-3'	5'- CAGTTCTTGGCTGGGAT GTT-3'
Human Caspase-1	5'- GGATAAGACCCGAGCTTTGATT-3'	5'- CTGCCAGGTA ACTGTCT TCTTC-3'
Human IL-1 β	5'-GGATAAGACCCGAGCTTTGATT-3'	5'- CTGCCAGGTA ACTGTCT TCTTC-3'
Human Cav-1	5'-CTTCGCCATTCTCTCTTCCT-3'	5'- GGTGTGGACGTAGATG GAATAG-3'
Human GAPDH	5'-GGTGTGAACCATGAGAAGTATGA-3'	5'- GAGTCCTTCCACGATAC CAAAG -3'



Supplementary Figure S1. Original full-length blots used in Figure 5 are shown.



Supplementary Figure S2. Original full-length blots used in Figure 7 are shown.