

Supplementary information

The regional specific alterations in BBB permeability are relevant to the differential responses of 67-kDa LR expression in endothelial cells and astrocytes following status epilepticus

Hana Park^{1,2}, Tae-Cheon Kang^{1,2*}


¹Department of Anatomy and Neurobiology, College of Medicine, Hallym University, Chuncheon 24252,
South Korea

²Institute of Epilepsy Research, College of Medicine, Hallym University, Chuncheon 24252, South Korea

Running title: 67-kDa LR-mediated regional specific vasogenic edema


* Correspondence to: T. -C. Kang, Department of Anatomy and Neurobiology, College of Medicine, Hallym University, Chuncheon, Kangwon-Do 24252, South Korea; Tel: +82-33-248-2524; Fax: +82-33-248-2525; E-mail: tckang@hallym.ac.kr

Fig. 1C



Supplementary Fig. 1. Full-length gel images of western blot data in Fig. 1C.

Fig. 4A



Supplementary Fig. 2. Full-length gel images of western blot data in Fig. 4A.

Fig. 6A





Fig. 6C



Supplementary Fig. 3. Full-length gel images of western blot data in Figs. 6A and 6C.

Fig. 7A





Fig. 9A



Supplementary Fig. 4. Full-length gel images of western blot data in Figs. 7A and 9A.

Fig. 11A





Fig. 11C



Supplementary Fig. 5. Full-length gel images of western blot data in Fig. 11A and 11C.