


# **Lung Innate Lymphoid Cell Composition is Altered in Primary Graft Dysfunction**

Laurel A Monticelli, Joshua M Diamond, Steven A Saenz,  
Elia D Tait Wojno, Mary K Porteous, Edward Cantu,  
David Artis, Jason D Christie


Online Data Supplement

**Online Data Supplement Figure E1: Flowchart of lung transplant patient enrollment for allograft sample collection**



**Online Data Supplement Figure E2: Gating strategy for ILC identification.**

Representative flow cytometric plots of live CD45<sup>+</sup> cells (A) and live CD45<sup>-</sup> cells (B) in a lung allograft.



**Online Data Supplement Table E1: Lung transplant donor characteristics**

<b>Donor Characteristic</b>	<b>PGD (n=8)</b>	<b>Non-PGD (n=10)</b>	<b>p-value</b>
Female Gender, n (%)	3 (37.5)	4 (40)	0.9
Donor Smoking History, yes, n (%)	5 (66.7)	7 (70)	0.7
Donor Alcohol History, yes, n (%)	4 (50)	6 (60)	0.7
Highest Donor PaO2 (mmHg)	435.5	495	0.4
Lowest Donor PaO2 (mmHg)	270	224	0.7
Donor Days on Ventilator	2.5	3	0.8
Oxygen Challenge PaO2 (mmHg)	453	430.7	0.2