

Supplementary Table 1. Comparison of annual eGFR decline (%/year) according to various clinical parameters after adjusting for age

Parameter	Annual eGFR decline, %	P value
Sex		
Female	-2.1±2.5	<0.001 ^a
Male	-0.4±2.4	
Obesity/BMI, kg/m ²		
<25	-1.3±0.1	0.097
≥25	-1.0±0.1	
Diabetes duration, yr		
<10	-1.0±0.1	<0.001 ^a
≥10	-1.8±0.2	
HbA1c, %		
<7	-0.7±0.1	<0.001 ^a
≥7	-1.4±0.1	
Hypertension		
No	-0.9±0.1	0.014 ^a
Yes	-1.4±0.1	
CVD		
No	-1.1±0.1	0.269
Yes	-1.4±0.3	
Diabetic retinopathy		
No	-1.1±0.1	0.003 ^a
Yes	-1.8±0.2	
Carotid plaque		
No	-0.9±0.1	<0.001 ^a
Yes	-1.6±0.1	
Albuminuria		
No	-1.1±0.1	0.428
Yes	-1.3±0.1	
Use of ACEi/ARB		
No	-1.0±0.1	0.014 ^a
Yes	-1.5±0.2	

Values are presented as mean ± standard error of mean.
eGFR, estimated glomerular filtration rate; BMI, body mass index; HbA1c, glycosylated hemoglobin; CVD, cardiovascular disease; ACEi, angiotensin-converting-enzyme inhibitor; ARB, angiotensin II-receptor blocker.

^aStatistical significance.