

Supplemental material

Supplementary table S1. Overview of the three retrospective collections on NSCLCs used in this study

Cohort	Total number of tumors	Number of quantified tumors	Clinical annotation	Immunotherapy treated	Tumor genotype available
Cohort 1 (YTMA250)	288	258	Yes	No	No
Cohort 2 (YTMA310)	138	111	No	Unknown	Yes
Cohort 3 (YTMA404)	81	69	Yes	Yes	Yes
Total	507	438			

Supplementary table S2. Panel for multiplexed CMTM6 and PD-L1 immunofluorescence staining

	Cytokeratin	CD68	CMTM6	PDL1
Primary antibodies	Rabbit polyclonal (Agilent) 1/100, 1 hour incubation	Clone PG-M1 (mouse IgG3, Abcam), 1 ug/ml, 4°C overnight incubation	Clone RCT6 (mouse IgG1, Absea), 0.4 ug/ml, 4°C overnight incubation	Clone E1L3N (rabbit IgG, Cell Signaling), 1 ug/ml, 4° overnight incubation
Secondary antibodies	Anti-rabbit Alexa488 (Invitrogen), 1/100, 1 hour incubation, RT	Anti-mouse IgG3 (ab97260, Abcam), 1/1000, 1 hour incubation, RT	Anti-mouse IgG1 (M1-14D12, eBioscience), 1/100, 1 hour incubation, RT	Anti-rabbit EnVision (K4009, Agilent), 1 hour incubation, RT
Fluorescent reagents		Biotynilated tyramid/Streptavidine-Alexa750 conjugate (Perkin-Elmer), 1 hour incubation, RT	Cy3+-tyramide (Perkin-Elmer), 10 min incubation, RT	Cy5-tyramide (Perkin-Elmer), 10 min incubation, RT
Counterstain	DAPI, 1/1000, 5 min incubation, RT			

Supplementary table S3. Panel for multiplexed TILs immunofluorescence staining


	Cytokeratin	CD8	CD4	CD20
Primary antibodies	Clone Z0622 (rabbit IgG, Agilent) 1/100, 1 hour incubation	Clone C8/144B (mouse IgG1, Agilent), 1 ug/ml, 4°C overnight incubation	Clone SP35 (rabbit IgG, Spring Bioscience), 1 ug/ml, 4°C overnight incubation	Clone L26 (Mouse IgG2a, Agilent), 1 ug/ml, 4° overnight incubation
Secondary antibodies	Anti-rabbit Alexa456 (Molecular Probes), 1/100, 1 hour incubation, RT	Anti-mouse IgG1 (M1-14D12, eBioscience), 1/100, 1 hour incubation, RT	Anti-rabbit envision (K4009, Agilent), 1 hour incubation, RT	Anti-mouse IgG2 (, Abcam), 1/200, 1 hour incubation, RT
Fluorescent reagents		Fluorescein-tyramide (Perkin-Elmer), 10 min incubation, RT	Biotynilated tyramid/Streptavidine-Alexa750 conjugate (Perkin-Elmer), 1 hour incubation, RT	Cy5-tyramide (Perkin-Elmer), 10 min incubation, RT
Counterstain	DAPI, 1/1000, 5 min incubation, RT			

Supplementary table S4. CMTM6 expression in the tumor compartment and clinicopathological characteristics in two independent NSCLC cohorts


Characteristic	YTMA404 cohort			YTMA250 cohort		
	CMTM6 low	CMTM6 high	p value	CMTM6 low	CMTM6 high	p value
Gender						
Male	16	22	0.18	50	656	0.62
Female	18	13		66	65	
Age						
< 70 yo	15	20	0.27	65	67	0.97
>= 70 yo	19	15		51	53	
ECOG						
0	2	4	0.29			
1	26	28				
2	6	2				
3	0	1				
Smoking history						
Never smoker	6	7	0.78	23	15	0.24
Current smoker	8	8		27	35	
Former smoker	19	20		58	63	
Histology						
Adenocarcinoma	27	23	0.52	77	58	0.02
Squamous-cell carcinoma	6	9		24	39	
Large-cell carcinoma	1	2		3	9	
Others	0	1		11	13	
Stage						
I			0.52	75	72	0.46
II				22	23	
III	2	0		11	19	
IV (M1a)	8	10		6	4	
IV (M1b)	5	5				
IV (M1c)	19	20				
EGFR mutation status						
Wild type	22	22	0.36			
Mutant	6	3				
KRAS mutation status						
Wild type	15	17	0.08			
Mutant	13	5				
CNS metastasis						
No	26	24	0.58			
Yes	8	10				
Liver metastasis						
No	28	28	1.00			
Yes	6	6				
LIPI score						
Good	16	12	0.72			
Intermediate	12	14				
Poor	2	2				

Supplementary table S5. CMTM6 expression in the stromal compartment and clinical-pathological characteristics in two independent NSCLC cohorts


Characteristic	YTMA404 cohort			YTMA250 cohort		
	CMTM6 low	CMTM6 high	p value	CMTM6 low	CMTM6 high	p value
Gender						
Male	19	19	0.89	53	53	0.77
Female	15	16		63	68	
Age						
< 70 yo	16	19	0.54	60	72	0.15
>= 70 yo	18	16		57	47	
ECOG						
0	3	3	0.80			
1	27	27				
2	4	4				
3	0	1				
Smoking history						
Never smoker	5	8	0.63	21	17	0.52
Current smoker	8	8		27	35	
Former smoker	20	19		59	62	
Histology						
Adenocarcinoma	25	25	0.70	71	64	0.45
Squamous-cell carcinoma	8	7		28	35	
Large-cell carcinoma	1	2		4	8	
Others	0	1		13	11	
Stage						
I			0.38	77	70	0.50
II				19	26	
III	2	0		13	17	
IV (M1a)	9	9		6	4	
IV (M1b)	6	4				
IV (M1c)	17	22				
EGFR mutation status						
Wild type	22	22	0.76			
Mutant	4	5				
KRAS mutation status						
Wild type	14	18	0.42			
Mutant	10	8				
CNS metastasis						
No	28	22	0.04			
Yes	5	13				
Liver metastasis						
No	26	30	0.45			
Yes	7	5				
LIPI score						
Good	15	13	0.53			
Intermediate	10	16				
Poor	2	2				




Supplementary figure S1. Validation of anti-CMTM6 antibody clone RCT6. (a)-(c) Titration curves of three different anti-CMTM6 antibodies used for assay validation plotted at five different concentrations: polyclonal antibody (ab198284) (a), monoclonal antibody clone RCT6 (b), and monoclonal antibody clone KT174 (c); (d)-(f) Representative membranous CMTM6 staining pattern with anti-CMTM6 polyclonal antibody (ab198284) (d), anti-CMTM6 monoclonal antibody clone RCT6 (e) and anti-CMTM6 monoclonal antibody clone KT174 (f). (g)-(h) Cross validation of anti-CMTM6 antibody clone RCT6 to the anti-CMTM6 polyclonal antibody (ab198284) (g) and to a second anti-CMTM6 monoclonal antibody (clone KT174) targeting non-overlapping epitopes (h). (i) Reproducibility of CMTM6 measurement with clone RCT6 in two independent multiplexed experiments



Supplementary figure S2. CMTM6 expression levels in the tumor compartment (a) and the stromal compartment (b) in *EGFR*- and *KRAS*-mutant NSCLCs (YTMA310 and YTMA404 cohorts combined)




Supplementary figure S3. Clinical benefit and PFS under single-agent PD-1 axis blockade according to CMTM6 expression in the tumor compartment (a,c) and the stromal compartment (b,d)




Supplementary figure S4. Correlation between CMTM6 and PDL1 expression levels in the tumor compartment (a) and the stromal compartment (b) in the three tested cohorts separately

Immunotherapy treated cohort (YTMA404)




Supplementary figure S5. OS under PD-1 axis blockade in patients in the top 30th percentile of PD-L1 expression in the tumor compartment (a) and the stromal compartment (b)

DAPI/CMTM6/PD-L1




Supplementary figure S6. Representative images showing CMTM6 and PD-L1 co-expression phenotypes in NSCLC: (a) CMTM6 high / PD-L1 high; (b) CMTM6 low / PD-L1 high; (c) CMTM6 high / PD-L1 low; (d) CMTM6 low / PD-L1 low

Immunotherapy treated cohort (YTMA404)




Supplementary figure S7. PFS under PD-1 axis blockade in the four CMTM6/PD-L1 co-expression phenotypes


Immunotherapy treated cohort (YTMA404)



Supplementary figure S8. Indicative performance of high CMTM6 and PD-L1 in the full immunotherapy-treated patient cohort (n = 69). (a)-(b) OS in patients with high CMTM6 and PD-L1 co-expression in the tumor compartment (a) and the stromal compartment (b); (c)-(d) OS according to PD-L1 expression in the tumor compartment (c) and the stromal compartment (d); (e)-(f) OS according to CMTM6 expression in the tumor compartment (e) and the stromal compartment (f)



Supplementary figure S9. Percentage of CMTM6 and PD-L1 pixel co-localization in the tumor, stroma, and CD68 compartments



Supplementary figure S10. Percentage of CMTM6 and PD-L1 colocalization in the tumor, stroma and CD68 compartments in tumors with high expression of CMTM6 and PD-L1 compared to the rest of the three tumor phenotypes combined