

## Supplemental Data

**S1 Table. MLD<sub>50</sub> of Balb/c-passaged clones in Balb/c mice**

Virus	Titer (PFU/mL)	No. of dead mice	No. of survival mice	Log MLD <sub>50</sub> (PFU/mL)
SW293	$2.79 \times 10^7$	0	4	> 7.45
	$2.79 \times 10^6$	0	5	
	$2.79 \times 10^5$	0	5	
	$2.79 \times 10^4$	0	5	
	$2.79 \times 10^3$	0	5	
	$2.79 \times 10^2$	0	5	
B_P8_C2	$7.88 \times 10^7$	5	0	4.40
	$7.88 \times 10^6$	5	0	
	$7.88 \times 10^5$	5	0	
	$7.88 \times 10^4$	5	0	
	$7.88 \times 10^3$	0	5	
	$7.88 \times 10^2$	0	5	
B_P9_C2	$4.78 \times 10^7$	5	0	4.30
	$4.78 \times 10^6$	5	0	
	$4.78 \times 10^5$	4	0	
	$4.78 \times 10^4$	4	1	
	$4.78 \times 10^3$	0	5	
	$4.78 \times 10^2$	0	5	

Balb/c mice were inoculated with SW293, B\_P8\_C2, or B\_P9\_C2, and the live and dead mice were counted daily for nine days. The 50% mouse lethal dose (MLD<sub>50</sub>) was calculated using Reed-Muench method.