Supplemental Table S1. Fractional Inhibitory Concentration Index Values for Dose-Matrix Assays

| Organism | IC ₅₀ of CAS ^a | IC ₅₀ of CAS with PUUP | IC ₅₀ of PUUP | IC ₅₀ of PUUP with CAS | FICI ^b |
|---|--------------------------------------|-----------------------------------|--------------------------|--------------------------------------|-------------------|
| Cryptococcus neoformans (strain H99) | 30° | 1.5 | 0.6 | 0.2 | 0.38 |
| Candida glabrata (strain 102) | 17 | 2 | 0.72 | 0.26 | 0.48 |
| Candida albicans (strain DPL1010) | 20 | 6 | 3.8 | 0.35 | 0.39 |

 $^{{}^{}a}\text{IC}_{50}$ values are based on dose response curves.

 $^{{}^}b\mathsf{FICI} = [\mathsf{A}^*]/[\mathsf{A}] + [\mathsf{B}^*]/[\mathsf{B}],$ where $[\mathsf{A}^*]$ is IC_{50} of compound A in the presence of compound B, $[\mathsf{A}]$ is IC_{50} of compound A alone, $[\mathsf{B}^*]$ is IC_{50} of compound B in the presence of compound A, and $[\mathsf{B}]$ is IC_{50} of compound B alone.

^cAll values are in μg/ml.