

**The effect of NAMPT deletion in projection neurons on
the function and structure of neuromuscular junction (NMJ) in mice**

Samuel Lundt¹, Nannan Zhang², Xiaowan Wang¹, Luis Polo-Parada^{2,3}, Shinghua Ding^{1,2*}

¹Dept. of Biomedical, Biological and Chemical Engineering

²Dalton Cardiovascular Research Center


³Dept. of Medical Pharmacology and Physiology

University of Missouri, Columbia, MO 65211


*Corresponding author: Shinghua Ding, Dalton Cardiovascular Research Center, Dept. of Biomedical, Biological and Chemical Engineering, University of Missouri-Columbia, 134 Research Park Drive, Columbia, MO 65211. E-mail: dings@missouri.edu. Tel: 573 884 2489.

Supplementary Figures

A



B



Supplemental Figure 1. Full Western blotting images of NAMPT levels in semitendinosus muscle from *Nampt^{fl/fl}*, *Nampt^{-/-}*, and *Nampt^{-/-}* +NMN mice. A) Original image of film incubated with anti-NAMPT antibody (Exposure time=7.9 seconds). B) Original image of film incubated with anti- β -actin (Exposure time=1.0 seconds). Images are from different gels that were run with the same sample order at the same time.