
1	Disruption of CXCR6 Ameliorates Kidney Inflammation and Fibrosis in
2	Deoxycorticosterone Acetate/Salt Hypertension
3	Supplemental Material
4	Yuanbo Wu <sup>1,2</sup> , Changlong An <sup>3</sup> , Xiaogao Jin <sup>1</sup> , Zhaoyong Hu <sup>1</sup> , and Yanlin Wang <sup>1,3,4,5,6*</sup>
5	<sup>1</sup> Section of Nephrology, Department of Medicine, Baylor College of Medicine,
6	Houston, Texas, USA
7	<sup>2</sup> Department of Anesthesiology, Hubei Cancer Hospital, Hubei, Wuhan, China
8	<sup>3</sup> Division of Nephrology, Department of Medicine, University of Connecticut School of
9	Medicine, Farmington, Connecticut, USA
10	<sup>4</sup> Department of Cell Biology, University of Connecticut School of Medicine,
11	Farmington, Connecticut, USA
12	<sup>5</sup> Institute for Systems Genomics, University of Connecticut School of Medicine,
13	Farmington, Connecticut, USA
14	<sup>6</sup> Renal Section, Veterans Affairs Connecticut Healthcare System, West Haven,
15	Connecticut, USA
16	Running Title: CXCR6 in deoxycorticosterone acetate/salt hypertension
17	*Corresponding author:
18	Yanlin Wang, MD, PhD
19	Division of Nephrology/Department of Medicine
20	University of Connecticut School of Medicine
21	263 Farmington Avenue
22	Farmington, CT 06030-1405
23	Email: <u>yanlwang@uchc.edu</u>
24	

Figure S1. Disruption of CXCR6 reduces  $\alpha$ -SMA protein expression in the kidney. A. Representative Western blots show the protein levels of  $\alpha$ -SMA in the kidneys of WT and CXCR6 KO mice 4 weeks after DOCA/salt or vehicle treatment. **B.** Quantitative Western blot analysis of  $\alpha$ -SMA protein levels in the kidneys. \*\* *P*<0.01 vs WT controls, \* *P*<0.05 vs KO-DOCA, and # *P*<0.05 vs WT-DOCA. n=6 per group.



31

32

33